Clinical Policy: Human Growth Hormone (Somapacitan, Somatrogon, Somatropin, Lonapegsomatropin-tcgd)

Reference Number: PA.CHIP.PHAR.517
Effective Date: 01/2026
Last Review Date: 01/2026

Description

The following human growth hormone (hGH) formulations require prior authorization:

- hGH analogs: somapacitan-beco (Sogroya®), somatrogon-ghla (Ngenla™)
- Recombinant hGH (rhGH) formulations: somatropin (Genotropin®, Humatrope®, Norditropin®, Nutropin AQ® NuSpin®, Omnitrope®, Saizen®, Serostim®, Zomacton®), longpegsomatropin-tcgd (Skytrofa®)

Drugs	Children								Adults	
	GHD	PWS	TS	NS	SHOX	CKD	SGA	ISS	GHD	HIV
Sogroya	GF			GF			SS	SS	X	
Genotropin	GF	GF	GF				GF	GF	X	
Humatrope	GF		SS		SS/GF		SS	SS/GF	X	
Ngenla	GF									
Norditropin	GF	GF	SS	SS			SS	SS	X	
NutropinAQ NuSpin	GF		GF			GF		GF	X	
Omnitrope	GF	GF	GF				GF	GF	X	
Saizen	GF								X	
Serostim										X
Skytrofa	GF								X	
Zomacton	GF		SS		SS		SS	SS	X	

Abbreviations: CKD: chronic kidney disease, GF: growth failure, GHD: growth hormone deficiency, HIV: human immunodeficiency virus, ISS: idiopathic short stature, NS: Noonan syndrome, PWS: Prader-Willi syndrome, SGA: small for gestational age, SHOX: short stature homeobox-containing gene, SS: short stature, TS: Turner syndrome

FDA Approved Indication(s)

hGH Analogs:

Sogroya is indicated for:

- Treatment of pediatric patients aged 2.5 years and older with: GF due to inadequate secretion of endogenous GH, SS born SGA and with no catch-up growth by 2 years of age, GF associated with NS, and ISS.
- Replacement of endogenous GH in adults with GHD

Ngenla is indicated for:

- Treatment of pediatric patients aged 3 years and older who have GF due to inadequate secretion of endogenous GH

rhGH Formulations:

Genotropin is indicated for treatment of:

- Children with GF due to GHD, PWS, SGA, TS, and ISS.
- Adults with either childhood-onset (CO) or adult-onset (AO) GHD.

Humatrope is indicated for treatment of:

- Pediatric patients: GF due to inadequate secretion of endogenous GH; SS associated with

TS; ISS, high standard deviation score (SDS) <- 2.25, and associated with growth rates unlikely to permit attainment of adult height in the normal range; SS or GF in SHOX deficiency; SS born small for SGA with no catch-up growth by 2 years to 4 years of age.
- Replacement of endogenous GH in adults with GHD.

Norditropin FlexPro is indicated for the treatment of:
- Children with GF due to GHD, SS associated with NS, SS associated with TS, SS born SGA with no catch-up growth by age 2 to 4 years, ISS, and GF due to PWS.
- Replacement of endogenous GH in adults with GHD.

Nutropin AQ NuSpin is indicated for the treatment of:
- Children with GF due to GHD, ISS, TS, and CKD up to the time of renal transplantation.
- Adults with either CO or AO GHD.

Omnitrope is indicated for the treatment of:
- Children with GF due to GHD, PWS, SGA, TS, and ISS.
- Adults with either CO or AO GHD.

Saizen is indicated for:
- Children with GF due to GHD.
- Adults with either CO or AO GHD.

Serostim is indicated for the treatment of:
- HIV patients with wasting or cachexia to increase lean body mass and body weight, and improve physical endurance.

Skytrofa is indicated for treatment of:
- Pediatric patients 1 year and older who weigh at least 11.5 kg and have GF due to inadequate secretion of endogenous GH.
- Replacement of endogenous GH in adults with GHD.

Zomacton is indicated for:
- Treatment of pediatric patients who have GF due to inadequate secretion of endogenous GH, SS associated with TS, ISS, SS or GF in SHOX deficiency, and SS born SGA with no catch-up growth by 2 years to 4 years.
- Replacement of endogenous GH in adults with GHD.

Policy/Criteria
Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of PA Health & Wellness® that Skytrofa, Sogroya, Ngenla, and somatropin are **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Growth Hormone Deficiency with Neonatal Hypoglycemia (off-label) (must meet all)**:**
1. Diagnosis of neonatal hypoglycemia due to GHD;
2. Request is for a somatropin formulation;
3. Prescribed by or in consultation with a pediatric endocrinologist;

4. Age ≤ 1 month;
5. Serum GH concentration ≤ 5 µg/L;
6. Member meets one of the following (a or b):
   a. Imaging shows hypothalamic-pituitary abnormality;
   b. Deficiency of ≥ 1 anterior pituitary hormone other than GH (e.g., ACTH, TSH, LH, FSH, prolactin);
7. The requested product is not prescribed concurrently with Increlex® (mecasermin);
8. If request is NOT for Norditropin or Omnitrope cartridge, one of the following (a, b, or c):
   a. Member must use Norditropin* and Omnitrope cartridge*;
   b. If both Norditropin and Omnitrope cartridge are not available (e.g., due to drug shortages) member must use Omnitrope* vial, unless contraindicated or clinically significant adverse effects are experienced;
   c. Norditropin and Omnitrope cartridge are all contraindicated or clinically significant adverse effects are experienced;

   **Prior authorization may be required for Norditropin and Omnitrope*
9. Dose does not exceed 0.30 mg/kg per week.

Approval duration: 12 months

B. Growth Hormone Deficiency with Short Stature/Growth Failure - Children *(open epiphyses)* (must meet all)**:**

1. Diagnosis of GHD;
2. Prescribed by or in consultation with a pediatric endocrinologist;
3. Age < 18 years;
4. If request is for Skytrofa, age ≥ 1 years and weight ≥ 11.5 kg;
5. If request is for Sogroya, age ≥ 2.5 years;
6. If request is for Ngenla, age ≥ 3 years;
7. If age > 10 years, open epiphysis on x-ray;
8. Member meets one of the following (a or b):
   a. Low insulin-like growth factor (IGF)-I serum level;
   b. Low insulin-like growth factor binding protein (IGFBP)-3 serum level;
9. Member meets one of the following (a, b, c, d, or e):
   a. Two GH stimulation tests with peak serum levels ≤ 10 µg/mL (e.g., stimulants: arginine, clonidine, glucagon);
   b. Deficiency of ≥ 3 pituitary hormones (i.e., ACTH, TSH, LH, FSH, prolactin);
   c. Prior surgery or radiotherapy to the hypothalamic-pituitary region;
   d. Imaging shows hypothalamic-pituitary abnormality;
   e. GHD-specific mutation (e.g., POU1F1, PROP1, LHX3, LHX4, HESX1, OTX2, TBX19, SOX2, SOX3, GLI2, GHRHR, GH1);
10. Member meets one of the following (a or b):
    a. SS: height is > 2 SD below the mean for age and sex (SD, height, date, and age in months within the last 90 days are required);
    b. GF: one of the following (i, ii, or iii):
       i. Height deceleration across two growth chart percentiles representing > 1 SD below the mean for age and sex (SD and 2 heights, dates, and ages in months at least 6 months apart within the last year are required);
       ii. Growth velocity > 2 SD below the mean for age and sex over 1 year (SD and 2 heights, dates, and ages in months at least 1 year apart within the last year are required);

iii. Growth velocity > 1.5 SD below the mean for age and sex sustained over 2 years (SD and 2 heights, dates, and ages in months at least 2 years apart within the last two years are required);

11. The requested product is not prescribed concurrently with Increlex (mecasermin);
12. If request is for a long acting growth hormone, (e.g.Ngenla, Skytrofa, Sogroya) (a or b);
    a. Member must use Sogroya* and Skytrofa*;
    b. Sogroya and Skytrofa are both contraindicated or clinically significant adverse effects are experienced;

**Prior authorization may be required*

13. If request is NOT for Norditropin*, Omnitrope cartridge*, Sogroya*, Ngenla*, or Skytrofa*, one of the following (a, b, or c):
    a. Member must use Norditropin* and Omnitrope cartridge*;
    b. If both Norditropin and Omnitrope cartridge are not available (e.g., due to drug shortages) member must use Omnitrope* vial, unless contraindicated or clinically significant adverse effects are experienced;
    c. Norditropin and Omnitrope cartridge are all contraindicated or clinically significant adverse effects are experienced;

    **Prior authorization may be required*

14. Dose does not exceed one of the following (a, b, c, or d):
    a. For Ngenla: 0.66 mg/kg once weekly;
    b. For Skytrofa: 0.24 mg/kg once weekly;
    c. For Sogroya: 0.16 mg/kg once weekly;
    d. For somatropin agents: 0.30 mg/kg per week.

Approval duration: 12 months

C. Genetic Disorders with Short Stature/Growth Failure - Children (must meet all)**:**

1. Diagnosis of PWS, TS, NS, or SHOX deficiency confirmed by a genetic test;
2. Request is for one of the following (a or b):
    a. Somatropin formulation;
    b. Somapacitan formulation, and member has diagnosis of NS;
3. Prescribed by or in consultation with a pediatric endocrinologist;
4. Age < 18 years;
5. If age > 10 years, open epiphysis on x-ray;
6. Member meets one of the following (a or b):
    a. SS: height is > 2 SD below the mean for age and sex (> 1.5 SD if TS) (SD, height, date, and age in months within the last 90 days are required);
    b. GF: one of the following (i, ii, or iii):
        i. Height deceleration across two growth chart percentiles representing > 1 SD below the mean for age and sex (SD and 2 heights, dates, and ages in months at least 6 months apart within the last year are required);
        ii. Growth velocity > 2 SD below the mean for age and sex over 1 year (SD and 2 heights, dates, and ages in months at least 1 year apart within the last year are required);
        iii. Growth velocity > 1.5 SD below the mean for age and sex sustained over 2 years (SD and 2 heights, dates, and ages in months at least 2 years apart within the last two years are required);
7. The requested product is not prescribed concurrently with Increlex (mecasermin);
8. If request is NOT for Norditropin or Omnitrope cartridge, one of the following (a, b, or c):

a. Member must use Norditropin* and Omnitrope cartridge*;

b. If both Norditropin and Omnitrope cartridge are not available (e.g., due to drug shortages) member must use Omnitrope* vial, unless contraindicated or clinically significant adverse effects are experienced;

c. Norditropin, Omnitrope cartridge are all contraindicated or clinically significant adverse effects are experienced;

**Prior authorization may be required for Norditropin and Omnitrope*

9. Request meets one of the following (a or b):
   a. For somatropin agents, one of the following (i, ii, or iii):
      i. PWS: Dose does not exceed 0.24 mg/kg per week;
      ii. TS, NS: Dose does not exceed 0.5 mg/kg per week;
      iii. SHOX deficiency: Dose does not exceed 0.35 mg/kg per week.
   b. Sogroya for NS: Dose does not exceed 0.24 mg/kg once weekly.

Approval duration: 12 months

D. Chronic Kidney Disease with Growth Failure – Children (must meet all)**:**

1. Diagnosis of CKD;
2. Request is for a somatropin formulation;
3. Prescribed by or in consultation with a pediatric endocrinologist or nephrologist;
4. Age < 18 years;
5. If age > 10 years, open epiphysis on x-ray;
6. Member meets one of the following (a, b, c, or d):
   a. GFR < 60 mL/min per 1.73 m^2 for ≥ 3 months;
   b. Dialysis dependent;
   c. Diagnosis of nephropathic cystinosis;
   d. History of kidney transplant ≥ 1 year ago;
7. Member meets one of the following (a or b):
   a. SS: height is > 2 SD below the mean for age and sex (SD, height, date, and age in months within the last 90 days are required);
   b. GF: one of the following (i, ii, or iii):
      i. Height deceleration across two growth chart percentiles representing > 1 SD below the mean for age and sex (SD and 2 heights, dates, and ages in months at least 6 months apart within the last year are required);
      ii. Growth velocity > 2 SD below the mean for age and sex over 1 year (SD and 2 heights, dates, and ages in months at least 1 year apart within the last year are required);
      iii. Growth velocity > 1.5 SD below the mean for age and sex sustained over 2 years (SD and 2 heights, dates, and ages in months at least 2 years apart within the last two years are required);
8. The requested product is not prescribed concurrently with Increlex (mecasermin);
9. If request is NOT for Norditropin or Omnitrope cartridge one of the following (a, b, or c):
   a. Member must use Norditropin* and Omnitrope cartridge*;
   b. If both Norditropin and Omnitrope cartridge are not available (e.g., due to drug shortages) member must use Omnitrope* vial, unless contraindicated or clinically significant adverse effects are experienced;
   c. Norditropin and Omnitrope cartridge are all contraindicated or clinically significant adverse effects are experienced;

**Prior authorization may be required for Norditropin and Omnitrope*

10. Dose does not exceed 0.35 mg/kg per week.

Approval duration: 12 months

E. Born Small for Gestational Age with Short Stature/Growth Failure - Children (must meet all)**:**

1. Diagnosis of SGA:
2. Request is for a somatropin or somapacitan formulation;
3. Prescribed by or in consultation with a pediatric endocrinologist;
4. Age ≥ 2 years and < 18 years;
5. If age > 10 years, open epiphysis on x-ray;
6. Birth weight or length > 2 SD below the mean for gestational age (SD, birth weight or length, and gestational age are required);
7. Current height > 2 SD below the mean for age and sex measured within the last year at ≥ 2 years of age (SD, height, date, and age in months are required);
8. The requested product is not prescribed concurrently with Increlex (mecasermin);
9. If request is NOT for Norditropin or Omnitrope cartridge, one of the following (a, b, or c):
   a. Member must use Norditropin* and Omnitrope cartridge*;
   b. If both Norditropin and Omnitrope cartridges are not available (e.g., due to drug shortages) member must use Omnitrope* vial, unless contraindicated or clinically significant adverse effects are experienced;
   c. Norditropin and Omnitrope cartridge are all contraindicated or clinically significant adverse effects are experienced;

   **Prior authorization may be required for Norditropin and Omnitrope*
10. Request meets one of the following (a or b):
    a. For somatropin agents: Dose does not exceed 0.48 mg/kg per week;
    b. For Sogroya: Dose does not exceed 0.24 mg/kg once weekly.

Approval duration: 12 months

F. Idiopathic Short Stature:

1. Use of Growth Hormones for the treatment of Idiopathic Short Stature is a benefit exclusion and will not be authorized because it is considered cosmetic in nature.

Approval duration: Not applicable

G. Growth Hormone Deficiency – Adults and Transition Patients *(closed epiphyses)* (must meet all):

1. Diagnosis of GHD;
2. Request is for a somatropin or somapacitan formulation;
3. Prescribed by or in consultation with an endocrinologist;
4. Age ≥ 18 years OR closed epiphysis on x-ray;
5. Member has NOT received somatropin therapy for ≥ 1 month prior to GH/IGF-I testing as outlined below;
6. Member meets one of the following (a, b, or c):
   a. Two fasting a.m. GH stimulation tests with peak serum levels ≤ 5 µg/mL (accepted stimulants: Macrilen™ [macimorelin] or combination of 2 stimulants such as arginine + glucagon);
   b. Both of the following (i and ii):
      i. One fasting a.m. GH stimulation test with peak serum level ≤ 5 µg/ml (accepted stimulants: Macrilen [macimorelin] or combination of 2 stimulants

such as arginine + glucagon);
ii. One low IGF-I serum level;

c. One low IGF-I serum level and one of the following (i, ii, or iii):
i. Imaging shows hypothalamic-pituitary abnormality;
ii. Deficiency of ≥ 3 pituitary hormones (i.e., ACTH, TSH, LH, FSH, prolactin);
iii. GHD-specific mutation (e.g., POU1F1, PROP1, LHX3, LHX4, HESX1, OTX2, TBX19, SOX2, SOX3, GLI2, GHRHR, GH1);

7. The requested product is not prescribed concurrently with Increlex (mecasermin);
8. One of the following (a or b):
a. Request is for a long-acting formulation (Skytrofa*, Sogroya*);
b. Request is NOT for Norditropin, Omnitrope cartridge, Skytrofa or Sogroya and one of the following (i, ii, or iii):
i. Member must use Norditropin* and Omnitrope cartridge*;
ii. If both Norditropin and Omnitrope cartridge are not available (e.g., due to drug shortages) member must use Omnitrope* vial, unless contraindicated or clinically significant adverse effects are experienced;
iii. Norditropin and Omnitrope cartridge are all contraindicated or clinically significant adverse effects are experienced;

**Prior authorization may be required for Norditropin, Omnitrope, Skytrofa, and Sogroya*

9. Dose does not exceed one of the following (a, b or c):
a. For Sogroya: 8 mg once weekly;
b. For somatropin agents: 0.4 mg/day (may adjust by up to 0.2 mg/day every 4 weeks to maintain normal IGF-1 serum levels; doses > 1.6 mg/day would be uncommon);
c. For Skytrofa: 6.3 mg once weekly.

Approval duration: 12 months

H. HIV-Associated Wasting or Cachexia (must meet all):
1. Diagnosis of HIV;
2. Request is for a somatropin formulation;
3. Prescribed by or in consultation with a physician specializing in HIV management;
4. Age ≥ 18 years;
5. Member meets one of the following (a, b, or c):
a. Unintentional weight loss of ≥ 10% in the last 12 months occurring while on antiretroviral therapy;
b. Unintentional weight loss of ≥ 5% in the last 6 months occurring while on antiretroviral therapy;
c. Body mass index (BMI) ≤ 20 kg/m^2;
6. Failure of at least 2 pharmacologic therapies from two separate drug classes *(Appendix B)* unless contraindicated or clinically adverse effects are experienced;
7. Member is currently on antiretroviral therapy;
8. If request is NOT for Norditropin or Omnitrope cartridge, one of the following (a, b, or c):
a. Member must use Norditropin* and Omnitrope cartridge*;
b. If both Norditropin and Omnitrope cartridge are not available (e.g., due to drug shortages) member must use Omnitrope* vial, unless contraindicated or clinically significant adverse effects are experienced;
c. Norditropin and Omnitrope cartridge are all contraindicated or clinically

significant adverse effects are experienced;
**Prior authorization may be required for Norditropin and Omnitrope*

9. Requested duration of therapy does not exceed 12 months;
10. Dose does not exceed 6 mg per day.

Approval duration: 12 months (up to 12 months total)

I. Other diagnoses/indications

1. If request is NOT for Norditropin or Omnitrope cartridge one of the following (a, b, or c):
   a. Member must use Norditropin* and Omnitrope cartridge*;
   b. If both Norditropin and Omnitrope cartridge are not available (e.g., due to drug shortages) member must use Omnitrope* vial, unless contraindicated or clinically significant adverse effects are experienced;
   c. Norditropin and Omnitrope cartridge are all contraindicated or clinically significant adverse effects are experienced;
   **Prior authorization may be required for Norditropin and Omnitrope*
2. Member meets one of the following (a or b):
   a. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (i or ii):
      i. For drugs on the, the no coverage criteria policy: CP.PMN.255; or
      ii. For drugs NOT on the formulary, the non-formulary: CP.PMN.16; or
   b. Refer to the off-label use policy if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.PMN.53

II. Continued Therapy

A. All Pediatric Indications *(open epiphyses)* (must meet all)**:**

1. Currently receiving medication via PA Health & Wellness benefit and documentation supports positive response to therapy or the Continuity of Care policy (PA.PHARM.01) applies;
2. Age < 18 years OR open epiphysis on x-ray;
3. Member meets one of the following (a or b):
   a. For diagnosis of neonatal hypoglycemia, when member has received somatropin therapy for ≥ 2 years, member's height has increased ≥ 2 cm in the last year as documented by 2 height measurements taken no more than 1 year apart (dates and height measurements required);
   b. For all other pediatric diagnoses, member's height has increased ≥ 2 cm in the last year as documented by 2 height measurements taken no more than 1 year apart (dates and height measurements required);
4. If request is for a dose increase, request meets one of the following (a, b, c or d):
   a. For Ngenla for GHD (without neonatal hypoglycemia): New dose does not exceed 0.66 mg/kg once weekly;
   b. For Skytrofa for GHD (without neonatal hypoglycemia): New dose does not exceed 0.24 mg/kg once weekly;
   c. For Sogroya, one of the following (i or ii):
      i. GHD (without neonatal hypoglycemia): New dose does not exceed 0.16 mg/kg per week;
      ii. Born SGA, NS: New dose does not exceed 0.24 mg/kg once weekly;
   d. For somatropin agents, one of the following (i, ii, iii, iv, or v):
      i. GHD (with or without neonatal hypoglycemia): New dose does not exceed

0.30 mg/kg per week;
ii. PWS: New dose does not exceed 0.24 mg/kg per week;
iii. TS, NS: New dose does not exceed 0.5 mg/kg per week;
iv. SHOX deficiency, CKD: New dose does not exceed 0.35 mg/kg per week;
v. Born SGA: New dose does not exceed 0.48 mg/kg per week.

Approval duration: 12 months

B. Growth Hormone Deficiency – Adults and Transition Patients *(closed epiphyses)* (must meet all)**:**

1. Currently receiving medication via PA Health & Wellness benefit and documentation supports positive response to therapy or the Continuity of Care policy (PA.PHARM.01) applies;
2. Member is responding positively to therapy;
3. For IGF-1 test results and dosing (test conducted within the last 90 days), one of the following (a, b, or c):
   a. Low IGF-1 serum level (i, ii ot iii):
      i. For Sogroya: 8 mg once weekly;
      ii. For somatropin formulations: If request is for a dose increase, new dose does not exceed an incremental increase of more than 0.2 mg/day and a total dose of 1.6 mg/day;
      iii. For Skytrofa: 6.3 mg once weekly;
   b. Normal IGF-1 serum level: Requested dose is for the same or lower dose;
   c. Elevated IGF-1 serum level: Requested dose has been titrated downward.

Approval duration: 12 months

C. HIV-Associated Wasting/Cachexia - Adults (must meet all):

1. Currently receiving medication via PA Health & Wellness benefit and documentation supports positive response to therapy or the Continuity of Care policy (PA.PHARM.01) applies.
2. Member is responding positively to therapy;
3. Member has not received ≥ 12 months of therapy;
4. If request is for a dose increase, new dose does not exceed 6 mg per day.

Approval duration: 12 months (up to 12 months total)

D. Idiopathic Short Stature:

1. Use of Growth Hormones for the treatment of Idiopathic Short Stature is a benefit exclusion and will not be authorized because it is considered cosmetic in nature.

Approval duration: Not applicable

E. Other diagnoses/indications (must meet 1 or 2)**:**

1. If request is NOT for Norditropin or Omnitrope cartridge one of the following (a, b, or c):
   a. Member must use Norditropin* and Omnitrope cartridge*;
   b. If both Norditropin and Omnitrope cartridge are not available (e.g., due to drug shortages) member must use Omnitrope* vial, unless contraindicated or clinically significant adverse effects are experienced;
   c. Norditropin and Omnitrope cartridge are all contraindicated or clinically significant adverse effects are experienced;

**Prior authorization may be required for Norditropin and Omnitrope*

2. Member meets one of the following (a or b):
   a. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (i or ii):
      i. For drugs on the, the no coverage criteria policy: CP.PMN.255; or
      ii. For drugs NOT on the formulary, the non-formulary: CP.PMN.16; or
   b. Refer to the off-label use policy if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.PMN.53

III. Diagnoses/Indications for which coverage is NOT authorized:

A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.PMN.53
B. Idiopathic short stature (ISS);
C. Constitutional delay of growth and puberty (i.e., constitutional growth delay; the member's growth rate is delayed compared to chronological age but appropriate for bone age as determined by x-ray);
D. Familial (genetic) short stature (i.e., height velocity and bone age, as determined by x- ray, are within the normal range and one or both parents are short);
E. Adult short stature or altered body habitus associated with antiviral therapy (other than HIV-associated wasting or cachexia);
F. Obesity treatment or enhancement of body mass/strength for non-medical reasons (e.g., athletic gains).

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

AO: adult-onset
CKD: chronic kidney disease
CO: childhood-onset
FDA: Food and Drug Administration
GF: growth failure
GFR: glomerular filtration rate
GH: growth hormone
GHD: growth hormone deficiency
hGH: human growth hormone
HIV: human immunodeficiency virus
IGF-1: insulin-like growth factor-1
IGFBP-3: insulin-like growth factor binding protein-3
ISS: idiopathic short stature
NS: Noonan syndrome
PWS: Prader-Willi syndrome
rhGH: recombinant human growth hormone
SD: standard deviation
SGA: small for gestational age
SHOX: short stature homeobox-containing gene
SS: short stature
TS: Turner syndrome

Appendix B: Therapeutic Alternatives
This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent and may require prior authorization.

Drug*	Dosing Regimen	Dose Limit/Maximum Dose
Appetite Stimulants		
megestrol (Megace®, Syndros®)	400 - 800 mg PO daily (10 – 20 ml/day)	800 mg/day
dronabinol (Marinol®)	2.5 mg PO BID	20 mg/day
Testosterone Replacement Products		
testosterone enanthate or cypionate (various brands)	50 - 400 mg IM Q2 – 4 wks	400 mg Q 2 wks
Androderm® (testosterone transdermal patch)	2.5 – 7.5 mg patch applied topically QD	7.5 mg/day
testosterone transdermal gel (Androgel®, Testim®)	5 - 10 gm gel (delivers 50 – 100 mg testosterone) applied topically QD	10 gm/day gel (100 mg/day testosterone)
Anabolic Steroids		
oxandrolone (Oxandrin®)	2.5 – 20 mg PO /day	20 mg/day
Nausea/Vomiting Treatments		
chlorpormazine	10 to 25 mg PO q4 to 6 hours prn	2,000 mg/day
perphenazine	8 to 16 mg/day PO in divided doses	64 mg/day
prochlorperazine	5 to 10 mg PO TID or QID	40 mg/day
promethazine	12.5 to 25 mg PO q4 to 6 hours prn	50 mg/dose; 100 mg/day
trimethobenzamide	300 mg PO TID or QID prn	1,200 mg/day

Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.
**Preferred status may be formulary specific.*

Appendix C: Contraindications/Boxed Warnings

- Contraindications:
  - Acute critical illness
  - Children with PWS who are severely obese, have history of upper airway obstruction or sleep apnea, or have severe respiratory impairment due to risk of sudden death
  - Active malignancy
  - Hypersensitivity to product or any of the excipients
  - Active proliferative or severe non-proliferative diabetic retinopathy
  - Children with closed epiphyses
- Boxed warning(s): none reported

Appendix D: Short Stature and Growth Failure

- For SS, the policy follows the World Health Organization (WHO) definition of > 2 SD below the mean for age and sex.[1]
- For GF, the policy follows:
  - Haymond et al (2013) and Rogol et al (2014) for height deceleration across two major

percentiles representing a change of > 1 SD corrected for age and sex[2,3] and

- the Growth Hormone Research Society (2000) for height velocity in the absence of SS that would prompt further investigation, namely, a height velocity > 2 SD below the mean over 1 year or > 1.5 SD below the mean sustained over 2 years for age and sex.[4]

- The Centers for Disease Control and Prevention (CDC) recommend WHO growth charts for infants and children age 0 to < 2 years and CDC growth charts for children age 2 years to < 20 years in the U.S.[5]
  - Based on CDC recommended growth chart data, SD approximations of major height percentiles falling below the mean are listed below:
    - 2nd percentile: 2 SD below the mean
    - 5th percentile: 1.5 SD below the mean
    - 15th percentile: 1 SD below the mean
    - 30th percentile: 0.5 SD below the mean
    - 50th percentile: 0 SD mean
  - CDC recommended growth charts, data tables, and related information that may be helpful in assessing length, height and growth are available at the following link: https://www.cdc.gov/growthcharts/index.htm.

1. WHO Child Growth Standards: Length/Height-for-Age, Weight-for-Age, Weight-for-Length, Weight-for-Height and Body Mass Index-for-Age: Methods and Development. Geneva, Switzerland: World Health Organization; 2006. As cited in CDC. Division of Nutrition, Physical Activity, and Obesity. Growth Chart Training: Using the WHO Growth Charts. Page last reviewed January 13, 2022. Available at https://www.who.int/publications/i/item/924154693X. Accessed December 3, 2024.
2. Haymond M, Kappelgaard AM, Czernichow P, et al. Early recognition of growth abnormalities permitting early intervention. Acta Pædiatrica ISSN 0803-5253. April 2013. DOI:10.1111/apa.12266.
3. Rogol AD, Hayden GF. Etiologies ad early diagnosis of short stature and growth failure in children and adolescents. J Pediatr. 2014 May;164(5 Suppl):S1-14.e6. doi: 10.1016/j.jpeds.2014.02.027.
4. Consensus guidelines for the diagnosis and treatment of growth hormone (GH) deficiency in childhood and adolescence: summary statement of the GH Research Society. JCEM. 2000; 85(11): 3990-3993.
5. Centers for Disease Control and Prevention, National Center for Health Statistics. CDC growth charts: United States. http://www.cdc.gov/growthcharts/. Accessed November 27. 2023.

V. Dosage and Administration

Drug Name	Indication	Dosing Regimen	Maximum Dose
Pediatric Indications (Subcutaneous administration; weekly doses should be divided [except Skytrofa, Sogroya, and Ngenla])			
Genotropin, Humatrope, Norditropin, Nutropin, Omnitrope, Saizen, Zomacton	GHD	G, O: 0.16 to 0.24 mg/kg/week H, Z: 0.18 to 0.30 mg/kg/week N: 0.17 to 0.24 mg/kg/week Nu: to 0.30 mg/kg/week S: 0.18 mg/kg/week	See dosing regimens
Genotropin, Norditropin, Omnitrope	PWS	G, N, O: 0.24 mg/kg/week	0.24 mg/kg/week
Genotropin, Humatrope, Norditropin, Omnitrope, Zomacton	SGA	G, O: to 0.48 mg/kg/week H, N, Z: to 0.47 mg/kg/week	0.48 mg/kg/week

Genotropin, Humatrope, Norditropin, Nutropin, Omnitrope, Zomacton	TS	G, O: 0.33 mg/kg/week H, Nu, Z: to 0.375 mg/kg/week N: to 0.47 mg/kg/week	See dosing regimens
Humatrope, Zomacton	SHOX	H, Z: 0.35 mg/kg/week	0.35 mg/kg/week
Norditropin	NS	0.46 mg/kg/week	0.46 mg/kg/week
Nutropin	CKD	0.35 mg/kg/week	0.35 mg/kg/week
Skytrofa	GHD	0.24 mg/kg/once weekly	0.24 mg/kg/week
Sogroya	DGH	0.16 mg/kg once weekly	0.16 mg/kg/week
	SGA, NS, ISS	0.24 mg/kg once weekly	0.24 mg/kg/week
Ngenla	GHD	0.66 mg/kg once weekly	0.66 mg/kg/week
Adult Indications (Subcutaneous administration)			
Genotropin, Humatrope, Norditropin, Nutropin, Omnitrope, Saizen, Zomacton	GHD	0.4 mg/day - may adjust by increments up to 0.2 mg/day every 6 weeks to maintain normal IGF-1 serum levels.* _____ **Dosing regimen from Endocrine Society guidelines (Fleseriu, et al., 2016).* Adult GHD dosing should be substantially lower than that prescribed for children. Adult doses beyond 1.6 mg/day would be uncommon.	See dosing regimen
Serostim	HIV-associated wasting	0.1 mg/kg QOD or QD to 6 mg QD	6 mg/day up to 24 weeks
Skytrofa	GHD	• 1.4 mg once weekly for adults 30 to 60 years old, with no oral estrogen intake • 2.1 mg once weekly for adults < 30 years old, or adults of any age intaking oral estrogen • 0.7 mg once weekly for adults > 60 years old, with no oral estrogen intake Increase the dose monthly to a higher strength cartridge based on the clinical response and/or IGF-1 concentration	6.3 mg/week

Sogroya	GHD	1.5 mg once weekly – increase by increments of 0.5-1.5 mg every 2-4 weeks based on clinical response and serum IGF-1 concentrations	8 mg/week

V. Product Availability

Drug	Availability*
hGH Analogs	
Sogroya	MD pens: 5 mg/1.5 mL, 10 mg/1.5 mL, 15 mg/1.5 mL
rhGH Formulations	
Genotropin lyophilized powder	MD dual-chamber syringes: 5 mg, 12 mg
Genotropin Miniquick	SD pen cartridges: 0.2 mg, 0.4 mg, 0.6 mg, 0.8 mg, 1.0 mg, 1.2 mg, 1.4 mg, 1.6 mg. 1.8 mg, 2.0 mg
Humatrope	MD pen cartridges: 6 mg, 12 mg, 24 mg MD vial: 5mg
Ngenla	MD pens: 24 mg/1.2 mL, 60 mg/1.2 mL
Norditropin Flexpro	MD pens: 5 mg/1.5 mL, 10 mg/1.5 mL, 15 mg/1.5 mL, 30 mg/3 mL
Nutropin AQ NuSpin	MD: 5 mg/2 mL, 10 mg/2 mL, 20 mg/2 mL MD pen cartridges: 10 mg/2 mL, 20 mg/2 mL
Omnitrope	MD pen cartridges: 5 mg/1.5 mL, 10 mg/1.5 mL MD vial: 5.8 mg
Saizen	MD pen cartridges: 8.8 mg MD vials: 5 mg, 8.8 mg
Serostim	MD vial: 4 mg SD vials: 5 mg, 6 mg
Skytrofa	SD prefilled cartridges: 0.7 mg, 1.4 mg, 1.8 mg, 2.1 mg, 2.5 mg, 3 mg, 3.6 mg, 4.3 mg, 5.2 mg, 6.3 mg, 7.6 mg, 9.1 mg, 11 mg, 13.3 mg
Zomacton	MD vials: 5 mg, 10 mg

SD: single-dose, MD: multidose

VI. References

FDA Labels

1. Genotropin Prescribing Information. NY, NY: Pfizer, Inc.; July 2025. Available at www.genotropin.com. Accessed November 6, 2025.
2. Humatrope Prescribing Information. Indianapolis, IN: Eli Lilly; July 2025. Available at: www.humatrope.com. Accessed November 6, 2025.
3. Ngenla Prescribing Information. New York, NY: Pfizer Labs; June 2023. Available at: https://www.accessdata.fda.gov/drugsatfda_docs/label/2023/761184Orig1s000Corrected_lbl.pdf. Accessed November 6, 2025.
4. Norditropin Prescribing Information. Plainsboro, NJ: Novo Nordisk; July 2025. Available at: www.norditropin.com. Accessed November 6, 2025.
5. Nutropin AQ. Prescribing Information. South San Francisco, CA: Genentech; December 2016. Available at: www.nutropin.com. Accessed November 6, 2025.
6. Omnitrope Prescribing Information. Princeton, NJ: Sandoz; July 2025. Available at: www.omnitrope.com. Accessed November 6, 2025.
7. Saizen Prescribing Information. Rockland, MA: Serono; February 2020. Available at:

https://www.emdserono.com/us-en/pi/saizen-ce-pi.pdf. Accessed November 6, 2025.
8. Serostim Prescribing Information. Rockland, MA: EMD Serono Inc.; June 2019. Available at: https://serostim.com/. Accessed November 6, 2025.
9. Skytrofa Prescribing Information. Princeton, New Jersey: Ascendis Pharma Endocrinology Inc., May 2024. Available at: https://skytrofa.com/. Accessed November 1, 2024.
10. Sogroya Prescribing Information. Plainsboro, NJ: NovoNordisk Health Care AG;. Available at: https://www.novo-pi.com/sogroya.pdf. Accessed March 3, 2026.
11. Zomacton Prescribing information. Parsippany, NJ: Ferring Pharmaceuticals Inc., July 2025. Available at: www.zomacton.com. Accessed November 6, 2025.

Compendia

12. DRUGDEX® System [Internet database]. Greenwood Village, Colo: Thomson Healthcare. Updated periodically.
13. Clinical Pharmacology [database online]. Tampa, FL: Gold Standard, Inc.; 2020. Available at https://www.clinicalkey.com/pharmacology/.

Somatropin Therapy - Children

14. Grimberg A, DiVall SA, Polychronakos C, et al. Guidelines for growth hormone and insulin-like growth factor-I treatment in children and adolescents: growth hormone deficiency, idiopathic short stature, and primary insulin-like growth factor-I deficiency. Horm Res Paediatr 2016; 86:361-397.
15. Rose SR, Cook DM, Fine MJ. Growth hormone therapy guidelines: Clinical and managed care perspectives. Am J Pharm Benefits. 2014;6(5):e134-e146.
16. Drube J, Wan M, Bonthuis M. Consensus statement: Clinical practice recommendations for growth hormone treatment in children with chronic kidney disease. Nephrology. September 2019; (15):S77-89.
17. National Kidney Foundation. KDOQI Clinical Practice Guideline for Nutrition in Children with CKD: 2008 Update. Am J Kidney Dis 53: S1-S124, 2009 (suppl 2).

GHD - Adults and Transition Patients

18. Yuen Keven CJ, Biller BMK, Radovick S, et al. American Association of Clinical Endocrinologists and American College of Endocrinology (AACE) guidelines for management of growth hormone deficiency in adults and patients transitioning from pediatric to adult care: 2019 AACE Growth Hormone Task Force. Endocrine Practice, November 2019; 25(11):1191-1232.
19. Fleseriu M, Hashim IA, Karavitaki N, et al. Hormonal replacement in hypopituitarism in adults: An Endocrine Society clinical practice guideline. J Clin Endocrinol Metab, November 2016, 101(11):3888 –3921 doi: 10.1210/jc.2016-2118.
20. Cook DM, Rose SR. A review of guidelines for use of growth hormone in pediatric and transition patients. Pituitary. September 2012, Volume 15, Issue 3, pp 301–310.
21. Molitch ME, Clemmons DR, Malozowski S, et al. Evaluation and treatment of adult growth hormone deficiency: an Endocrine Society clinical practice guideline. J Clin Endocrinol Metab. 2011; 96: 1587-1609.

HIV-Associated Wasting

22. Badowski ME, Perez SE. Clinical utility of dronabinol in the treatment of weight loss associated with HIV and AIDS. HIV AIDS (Auckl). 2016 Feb 10;8:37-45. doi: 10.2147/HIV.S81420. eCollection 2016.
23. Nemechek PM, Polsky B, Gottlieb M. Treatment guidelines for HIV-associated wasting. Subspecialty clinics: Infectious Diseases 2000;74(4):p386-394.
24. Wanke C, Kotler D, and the HIV Wasting Collaborative Consensus Committee. Collaborative recommendation: The approach to diagnosis and treatment of HIV wasting. Journal of Acquired Immune Deficiency Syndromes 2004:37;pS284-S288.

25. Bedimo R, Hardy D, Lee D, Palella F, and Wohl D. Expert consensus statement on an updated definition of unintended weight loss among persons with human immunodeficiency virus in the modern treatment era. CID 2024;79:S63-75.

Somatropin Product Comparative Data

26. Romer T, Zabransky M, Walczak M, Szalecki M, and Balser S. Effect of switching recombinant human growth hormone: comparative analysis of phase 3 clinical data. Biol Ther 2011; 1(2):005. DOI 10.1007/s13554-011-0004-8

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J2941	Injection, somatropin, 1 mg
C9399	Unclassified drugs or biologics
J3590	Unclassified biologics

Reviews, Revisions, and Approvals	Date
Policy created	09/2025
1Q 2026 annual review: added Sogroya as a co-preferred long-acting growth hormone alongside Skytrofa; RT4: for Skytrofa, added new indication for replacement of endogenous GH in adults with GHD and added new cartridge strengths (0.7 mg, 1.4 mg, 1.8 mg, 2.1 mg, 2.5 mg); removed Zorbtive from policy due to market discontinuation; removed criteria for short bowel syndrome due to lack of support by AGA; for HIV-associated wasting, added option for unintentional weight loss of ≥ 5% in the last 6 months while on antiretroviral and removed use of ideal body weight criteria per update 2024 consensus expert statement; extended initial approval duration from 6 to 12 months for this maintenance medication for a chronic condition; references reviewed and updated.	01/2026