

Clinical Policy: Omalizumab (Xolair), Omalizumab-igec (Omlyclo)

Reference Number: PA.CHIP.PHAR.01

Effective Date: 01/2026

Last Review Date: 09/2025

Description

Omalizumab (Xolair®) and its biosimilar omalizumab-igec (Omlyclo®) are anti-immunoglobulin E (IgE) antibodies.

FDA Approved Indication(s)

Xolair and Omlyclo are indicated for:

- Moderate to severe persistent asthma in adults and pediatric patients 6 years of age and older with a positive skin test or in vitro reactivity to a perennial aeroallergen and symptoms that are inadequately controlled with inhaled corticosteroids
- Chronic rhinosinusitis with nasal polyps (CRSwNP) in adult patients 18 years of age and older with inadequate response to nasal corticosteroids, as add-on maintenance treatment
- Chronic spontaneous urticaria (CSU) in adults and adolescents 12 years of age and older who remain symptomatic despite H1 antihistamine treatment
- Reduction of allergic reactions (Type I), including anaphylaxis, that may occur with accidental exposure to one or more foods in adult and pediatric patients aged 1 year and older with IgE-mediated food allergy. Xolair and Omlyclo are to be used in conjunction with food allergen avoidance

Limitation(s) of use: Xolair and Omlyclo are not indicated for the relief of acute bronchospasm or status asthmaticus, treatment of other forms of urticaria, or emergency treatment of allergic reactions including anaphylaxis.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of PA Health & Wellness® that Xolair and Omlyclo are **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Moderate to Severe Persistent Asthma (must meet all):

1. Diagnosis of asthma;
2. Age \geq 6 years;
3. Member has experienced \geq 1 exacerbation within the last 12 months, requiring any of the following despite adherent use of controller therapy (i.e., medium- to high-dose inhaled corticosteroid [ICS] plus a long-acting beta₂ agonist [LABA] or ICS plus one additional asthma controller medication):
 - a. Oral/systemic corticosteroid treatment (or increase in dose if already on

oral corticosteroid);

- b. Urgent care/emergency room (ER) visit or hospital admission;
4. Positive skin test or in vitro reactivity to a perennial aeroallergen (*see Appendix D*);
5. IgE level ≥ 30 IU/mL;
6. Xolair/Omlyclo is prescribed concurrently with an ICS plus either a LABA or one additional asthma controller medication;
7. Xolair/Omlyclo is not prescribed concurrently with Cinqair[®], Fasenra[®], Nucala[®], Dupixent[®], or Tezspire[®];
8. Dose does not exceed 375 mg administered every 2 weeks (*see Appendix E and F for dosing based on pre-treatment IgE level, weight, and age*).

Approval duration: 6 months

B. Chronic Spontaneous Urticaria (must meet all):

1. Diagnosis of CSU (formerly known as chronic idiopathic urticaria [CIU]);
2. Age ≥ 12 years;
3. Failure of one antihistamine at maximum indicated doses used for ≥ 2 weeks, unless clinically significant adverse effects are experienced or all are contraindicated;
4. Xolair/Omlyclo is not prescribed concurrently with Cinqair, Fasenra, Nucala, Dupixent, or Tezspire;
5. Dose does not exceed 300 mg every 4 weeks.

Approval duration: 6 months

C. Chronic Rhinosinusitis with Nasal Polypsis (must meet all):

1. Diagnosis of CRSwNP with documentation of all of the following (a, b, and c):
 - a. Presence of nasal polyps;
 - b. Member has experienced signs and symptoms (e.g., nasal congestion/blockage/ obstruction, loss of smell, rhinorrhea) for ≥ 12 weeks;
2. Prescribed by or in consultation with an allergist, immunologist, or otolaryngologist;
3. Age ≥ 18 years;
4. Member has required the use of systemic corticosteroids for symptom control within the last 2 years, unless contraindicated or clinically significant adverse effects are experienced (*see Appendix B for examples*);
5. Failure of maintenance therapy with one intranasal corticosteroid used for ≥ 4 weeks, unless contraindicated or clinically significant adverse effects are experienced (*see Appendix B for examples*);
6. Xolair/Omlyclo is prescribed concurrently with an intranasal corticosteroid, unless contraindicated or clinically significant adverse effects are experienced (*see Appendix B for examples*);
7. Xolair/Omlyclo is not prescribed concurrently with Cinqair, Fasenra, Nucala, Dupixent, or Tezspire;
8. Dose does not exceed 600 mg every 2 weeks (*see Appendix G for dosing based on pre-treatment IgE level and weight*).

Approval duration: 6 months

D. IgE-Mediated Food Allergy (must meet all):

1. Diagnosis of IgE-mediated food allergy;
2. Prescribed by or in consultation with an allergist or immunologist;
3. Age \geq 1 year;
4. Confirmation of one of the following (a, b, or c):
 - a. Positive skin prick test with wheal diameter \geq 4 mm greater than control;
 - b. Food-specific serum IgE \geq 6 kUA/L;
 - c. Positive oral food challenge test;
5. Member has history of significant allergic reaction(s) to the food (e.g., hives, swelling, wheezing, hypotension, gastrointestinal symptoms) that meets both of the following (a and b):
 - a. Prescriber deemed past allergic reaction to the food significant enough to require a prescription for injectable epinephrine;
 - b. Xolair/Omlyclo is prescribed concurrently with injectable epinephrine;
6. Medical justification supports necessity for Xolair/Omlyclo despite food allergen avoidance (e.g., member lacks sufficient mental capacity to effectively avoid food allergens);
7. Xolair/Omlyclo is not prescribed concurrently with Palforzia™, Cinqair, Fasenra, Nucala, Dupixent, or Tezspire;
8. Dose does not exceed 600 mg every 2 weeks (*see Appendix H for dosing based on pre-treatment IgE level and weight*).

Approval duration: 6 months

E. NCCN Compendium Indications (off-label) (must meet all):

1. Diagnosis of one of the following (a or b):
 - a. Systemic mastocytosis;
 - b. Immune checkpoint inhibitor-related severe (G3; *see Appendix I*) pruritus;
2. Prescribed by or in consultation with an oncologist;
3. For systemic mastocytosis, prescribed in one of the following settings (a, b, c, or d):
 - a. As stepwise prophylactic treatment for chronic mast cell mediator-related cardiovascular and pulmonary symptoms when the member has tried both of the following, unless clinically significant adverse effects are experienced or all are contraindicated (i and ii):
 - i. Antihistamine (i.e., H1 blocker, H2 blocker);
 - ii. Corticosteroid;
 - b. For prevention of unprovoked anaphylaxis;
 - c. For prevention of hymenoptera (e.g., bees, wasps, hornets) or food-induced anaphylaxis, and one of the following (i or ii):
 - i. Member has negative specific IgE;
 - ii. Member has negative skin test;
 - d. To improve tolerability of immunotherapy;
4. For immune checkpoint inhibitor-related severe pruritis, all of the following (a, b, and c):
 - a. Pruritus is refractory;
 - b. Member has an increased IgE level;
 - c. Member has not responded to a gabapentinoid (e.g., gabapentin, pregabalin) after 1 month of therapy;

5. Xolair/Omlyclo is not prescribed concurrently with Cinqair, Fasenra, Nucala, Dupixent, or Tezspire;
6. Dose is within FDA maximum limit for any FDA-approved indication or is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).*

**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration: 6 months

F. Other diagnoses/indications

1. Refer to the off-label use policy if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.PMN.53

II. Continued Therapy

A. Moderate to Severe Persistent Asthma (must meet all):

1. Currently receiving medication via PA Health & Wellness benefit and documentation supports positive response to therapy or the Continuity of Care policy (PA.PHARM.01) applies.
2. Demonstrated adherence to asthma controller therapy (an ICS plus either a LABA or LTRA) as evidenced by proportion of days covered (PDC) of 0.8 in the last 6 months (i.e., member has received asthma controller therapy for at least 5 of the last 6 months);
3. Member is responding positively to therapy (examples may include but are not limited to: reduction in exacerbations or corticosteroid dose, improvement in forced expiratory volume over one second since baseline, reduction in the use of rescue therapy);
4. Xolair/Omlyclo is not prescribed concurrently with Cinqair, Fasenra, Nucala, Dupixent, or Tezspire;
5. If request is for a dose increase, new dose does not exceed 375 mg every 2 weeks (*see Appendix E and F for dosing based on pre-treatment IgE level, weight, and age*).

Approval duration: 12 months

B. Chronic Spontaneous Urticaria (must meet all):

1. Currently receiving medication via PA Health & Wellness benefit and documentation supports positive response to therapy or the Continuity of Care policy (PA.PHARM.01) applies.
2. Member is responding positively to therapy;
3. Xolair/Omlyclo is not prescribed concurrently with Cinqair, Fasenra, Nucala, Dupixent, or Tezspire;
4. If request is for a dose increase, new dose does not exceed 300 mg every 4 weeks.

Approval duration: 12 months

C. Chronic Rhinosinusitis with Nasal Polypsis (must meet all):

1. Currently receiving medication via PA Health & Wellness benefit and documentation supports positive response to therapy or the Continuity of Care policy (PA.PHARM.01) applies.
2. Demonstrated adherence to an intranasal corticosteroid, unless contraindicated or clinically significant adverse effects are experienced;

3. Member is responding positively to therapy (examples may include but are not limited to: reduced nasal polyp size, reduced need for systemic corticosteroids, improved sense of smell, improved quality of life);
4. Xolair/Omlyclo is not prescribed concurrently with Cinqair, Fasenra, Nucala, Dupixent, or Tezspire;
5. If request is for a dose increase, new dose does not exceed 600 mg every 2 weeks (*see Appendix G for dosing based on pre-treatment IgE level and weight*).

Approval duration: 12 months

D. IgE-Mediated Food Allergy (must meet all):

1. Currently receiving medication via PA Health & Wellness benefit and documentation supports positive response to therapy or the Continuity of Care policy (PA.PHARM.01) applies.
2. Member is responding positively to therapy;
3. Xolair/Omlyclo is prescribed concurrently with injectable epinephrine;
4. Xolair/Omlyclo is not prescribed concurrently with Palforzia, Cinqair, Fasenra, Nucala, Dupixent, or Tezspire;
5. If request is for a dose increase, new dose does not exceed 600 mg every 2 weeks (*see Appendix H for dosing based on pre-treatment IgE level and weight*).

Approval duration: 12 months

E. NCCN Compendium Indications (off-label) (must meet all):

1. Currently receiving medication via PA Health & Wellness benefit and documentation supports positive response to therapy or the Continuity of Care policy (PA.PHARM.01) applies.
2. Member is responding positively to therapy (examples may include but are not limited to: improvement in itching or skin pain, reduction in number of nodules);
3. Dupixent is not prescribed concurrently with another biologic immunomodulator (e.g., Adbry, Cinqair, Fasenra, Nucala, Tezspire, Xolair) or a JAK inhibitor (e.g., Olumiant, Rinvoq, Cibinquo, Opzelura);
4. If request is for a dose increase, new dose does not exceed 300 mg every other week.

Approval duration: 12 months

F. Chronic Obstructive Pulmonary Disease (must meet all):

1. Currently receiving medication via PA Health & Wellness benefit and documentation supports positive response to therapy or the Continuity of Care policy (PA.PHARM.01) applies.
2. Member is responding positively to therapy;
3. If request is for a dose increase, new dose is within FDA maximum limit for any FDA-approved indication or is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).*

**Prescribed regimen must be FDA-approved or recommended by NCCN.*

Approval duration: 6 months

G. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via PA Health & Wellness benefit and documentation supports positive response to therapy or the Continuity of Care policy (PA.PHARM.01) applies.

Approval duration: Duration of request or 12 months (whichever is less); or

2. Refer to the off-label use policy if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.PMN.53

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies CP.PMN.53
- B. Acute bronchospasm or status asthmaticus.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

AAAAI: American Academy of Allergy, Asthma, and Immunology

ADL: activity of daily living

CIU: chronic idiopathic urticaria

CRSwNP: chronic rhinosinusitis with nasal polyps

CSU: chronic spontaneous urticaria

EAACI: European Academy of Allergy and Clinical Immunology

EDF: European Dermatology Forum

EPR3: Expert Panel Report 3

FDA: Food and Drug Administration

GA2LEN: Global Allergy and Asthma European Network

GINA: Global Initiative for Asthma

ICS: inhaled corticosteroids

IgE: immunoglobulin E

kUA/L: kilounits of allergen-specific IgE per liter

LABA: long-acting beta-agonist

LTRA: leukotriene modifier

PDC: proportion of days covered

WAO: World Allergy Organization

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
ASTHMA		
ICS (medium – high dose)		
Qvar® (beclomethasone)	> 100 mcg/day 40 mcg, 80 mcg per actuation 1-4 actuations BID	4 actuations BID
budesonide (Pulmicort®)	> 200 mcg/day 90 mcg, 180 mcg per actuation 2-4 actuations BID	2 actuations BID
Alvesco® (ciclesonide)	> 80 mcg/day 80 mcg, 160 mcg per actuation 1-2 actuations BID	2 actuations BID

CLINICAL POLICY
Omalizumab, Omalizumab-igec

fluticasone propionate (Flovent®)	> 100 mcg/day 44-250 mcg per actuation 2-4 actuations BID	2 actuations BID
Arnuity Ellipta® (fluticasone furoate)	≥ 50 mcg/day 100 mcg, 200 mcg per actuation 1 actuation QD	1 actuation QD
Asmanex® (mometasone)	> 100 mcg/day HFA: 100 mcg, 200 mcg per actuation Twisthaler: 110 mcg, 220 mcg per actuation 1-2 actuations QD to BID	2 inhalations BID
LABA		
Serevent® (salmeterol)	50 mcg per dose 1 inhalation BID	1 inhalation BID
Combination Products (ICS + LABA)		
Dulera® (mometasone/ formoterol)	100/5 mcg, 200/5 mcg per actuation 2 actuations BID	4 actuations per day
Breo Ellipta® (fluticasone/vilanterol)	100/25 mcg, 200/25 mcg per actuation 1 actuation QD	1 actuation QD
fluticasone/ salmeterol (Advair®)	Diskus: 100/50 mcg, 250/50 mcg, 500/50 mcg per actuation HFA: 45/21 mcg, 115/21 mcg, 230/21 mcg per actuation 1 actuation BID	1 actuation BID
fluticasone/salmeterol (Airello RespiClick®)	55/13 mcg, 113/14 mcg, 232/14 mcg per actuation 1 actuation BID	1 actuation BID
budesonide/ formoterol (Symbicort®)	80 mcg/4.5 mcg, 160 mcg/4.5 mcg per actuation 2 actuations BID	2 actuations BID
LTRA		
montelukast (Singulair®)	4 to 10 mg PO QD	10 mg per day
zafirlukast (Accolate®)	10 to 20 mg PO BID	40 mg per day
zileuton ER (Zyflo® CR)	1,200 mg PO BID	2,400 mg per day
Zyflo® (zileuton)	600 mg PO QID	2,400 mg per day
Oral Corticosteroids		
dexamethasone (Decadron®)	0.75 to 9 mg/day PO in 2 to 4 divided doses	Varies
methylprednisolone (Medrol®)	40 to 80 mg PO in 1 to 2 divided doses	Varies
prednisolone (Millipred®, Orapred ODT®)	40 to 80 mg PO in 1 to 2 divided doses	Varies
prednisone (Deltasone®)	40 to 80 mg PO in 1 to 2 divided doses	Varies
CSU		

CLINICAL POLICY
Omalizumab, Omalizumab-igec



hydroxyzine (Vistaril®)	Adult: 25 mg PO TID to QID Age ≥ 6 years: 50 mg-100 mg/day in divided doses	Adult: Will vary according to condition Age ≥ 6 years: 50 mg- 100 mg/day in divided doses
diphenhydramine (Benadryl®)	Adult: 25 mg to 50 mg PO TID to QID Pediatric: 12.5 mg to 25 mg PO TID to QID or 5 mg/kg/day or 150 mg/m ² /day	Adult: Will vary according to condition Children: 300 mg/day
chlorpheniramine (Aller- Chlor®)	Immediate Release: 4 mg PO every 4 to 6 hours Extended Release: 12 mg PO every 12 hours	Do not exceed 24 mg/day
cetirizine (Zyrtec®)	5 to 10 mg PO QD	10 mg/day
levocetirizine (Xyzal®)	2.5 mg to 5 mg PO QD	5 mg/day
loratadine (Claritin®)	10 mg PO QD	10 mg/day
desloratadine (Clarinex®)	5 mg PO QD	Will vary according to condition
fexofenadine (Allegra®)	60 mg PO BID or 180 mg QD	180 mg/day
CRSwNP		
Intranasal Corticosteroids		
beclomethasone (Beconase AQ®, Qnasl®)	1-2 sprays IN BID	2 sprays/nostril BID
budesonide (Rhinocort® Aqua, Rhinocort®)	128 mcg IN QD or 200 mcg IN BID	1-2 inhalations/nostril/day
flunisolide	2 sprays IN BID	2 sprays/nostril TID
fluticasone propionate (Flonase®)	1-2 sprays IN BID	2 sprays/nostril BID
mometasone (Nasonex®)	2 sprays IN BID	2 sprays/nostril BID
Omnaris®, Zetonna® (ciclesonide)	Omnaris: 2 sprays IN QD Zetonna: 1 spray IN QD	Omnaris: 2 sprays/nostril/day Zetonna: 2 sprays/nostril/day
triamcinolone (Nasacort®)	2 sprays IN QD	2 sprays/ nostril/day
Xhance™ (fluticasone propionate)	1 to 2 sprays (93 mcg/spray) to nostril IN BID	744 mcg/day
Oral Corticosteroids		
dexamethasone (Decadron®)	0.75 to 9 mg/day PO in 2 to 4 divided doses	Varies
methylprednisolone (Medrol®)	4 to 48 mg PO in 1 to 2 divided doses	Varies
prednisolone (Millipred®, Orapred ODT®)	5 to 60 mg PO in 1 to 2 divided doses	Varies

prednisone (Deltasone®)	5 to 60 mg PO in 1 to 2 divided doses	Varies
Systemic mastocytosis, Immunotherapy-related pruritus		
antihistamines, H1 blockers: examples – diphenhydramine, chlorpheniramine, hydroxyzine, cetirizine, loratadine, fexofenadine	Varies	Varies
antihistamines, H2 blockers: examples – cimetidine, famotidine,	Varies	Varies
corticosteroids: examples – betamethasone, dexamethasone, methylprednisolone, prednisolone, prednisone	Varies	Varies

Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): hypersensitivity
- Boxed warning(s): anaphylaxis

Appendix D: General Information

- Allergic asthma:
 - The definition of moderate to severe allergy varied among the clinical trials. The definition most often used was a patient who required oral systemic steroid bursts or unscheduled physician office visits for “uncontrolled” asthma exacerbations despite maintenance inhaled steroid use. Patients in the clinical trials most often were required to have an FEV1 between 40% and 80% of predicted. No patients were enrolled with an FEV1 greater than 80% of predicted.
 - Omalizumab has been shown to be marginally effective in decreasing the incidence of asthma exacerbations in patients who have met all the criteria described above.
 - Omalizumab provides little therapeutic benefit over existing therapies. Use in

patients on inhaled corticosteroids or chronic oral steroids plus or minus a second controller agent decreased asthma exacerbation by 0.5 to 1 per year. Use of rescue beta-agonists declined by 1 inhalation per day. Small changes in pulmonary function tests were also seen. An analysis of unpublished data indicated that hospital admissions declined by 3 per hundred patient years, emergency department (ED) visits by 2 per hundred patient years, and unscheduled physician office visits by 14 per one hundred patient years.

- The 2007 National Heart, Lung and Blood Institute's Expert Panel Report 3 (EPR3) Guidelines for the Diagnosis and Management of Asthma recommend Omalizumab may be considered as adjunct therapy for patients 12 years and older with allergies and Step 5 or 6 (severe) asthma whose symptoms have not been controlled by ICS and LABA.
- The Global Initiative for Asthma (GINA) guidelines recommend Omalizumab be considered as adjunct therapy for patients 6 years of age and older with exacerbations or poor symptom control despite taking at least high dose ICS/LABA and who have allergic biomarkers or need maintenance oral corticosteroids.
- The four perennial aeroallergens most commonly tested for in the clinical trials were dog dander, cat dander, cockroach, and house dust mite.
- Serious and life-threatening allergic reactions (anaphylaxis) in patients after treatment with Omalizumab have been reported. Usually these reactions occur within two hours of receiving a Omalizumab subcutaneous injection. However, these new reports include patients who had delayed anaphylaxis—with onset two to 24 hours or even longer—after receiving Omalizumab treatment. Anaphylaxis may occur after any dose of Omalizumab (including the first dose), even if the patient had no allergic reaction to the first dose.
- Patients could potentially meet asthma criteria for both Omalizumab and Nucala, though there is insufficient data to support the combination use of multiple asthma biologics. The combination has not been studied. Approximately 30% of patients in the Nucala MENSA study also were candidates for therapy with Omalizumab.
- PDC is a measure of adherence. PDC is calculated as the sum of days covered in a time frame divided by the number of days in the time frame. To achieve a PDC of 0.8, a member must have received their asthma controller therapy for 144 days out of the last 180 days, or approximately 5 months of the last 6 months.
- CSU:
 - CSU is classified as spontaneous onset of wheals, angioedema, or both, for more than 6 weeks due to an unknown cause.
 - Clinical studies have shown that Omalizumab 150 mg and 300 mg significantly improved the signs and symptoms of chronic idiopathic urticaria compared to placebo in patients who had remained symptomatic despite the use of approved dose of H1-antihistamine.
 - The Joint Task Force on Practice Parameters representing various American allergy organizations include Omalizumab in combination with H1-antihistamines as a fourth line treatment option following a stepwise approach starting with a second generation antihistamine. This is followed by one or more of the following: a dose increase of the second generation antihistamine, or the addition of another second generation antihistamine, H2-antagonist, LTRA, or first generation antihistamine. Treatment with hydroxyzine or doxepin can be considered in patients whose symptoms remain poorly controlled.

CLINICAL POLICY

Omalizumab, Omalizumab-igec



- The EAACI/GA2LEN/EDF/AAAAI/WAO Guideline for the Management of Urticaria include Omalizumab combination with H₁-antihistamines as a third line treatment option in patients who have failed to respond to higher doses of H₁- antihistamines.
- Omalizumab is the first medicine in its class approved for CSU since non-sedating antihistamines.
- The use of over-the-counter H₁ antihistamines may not be a benefit to the treatment of CIU. Credit will be given for their use, but will not be covered under plan.
- Anaphylaxis has occurred as early as after the first dose of Omalizumab, but also occurred beyond 1 year after beginning regularly administered treatment.
- Idiopathic anaphylaxis: A randomized, double-blind, placebo-controlled study in 19 patients with frequent episodes ($\geq 6/\text{year}$) of idiopathic anaphylaxis found Omalizumab to have no significant difference compared to placebo in the number of anaphylactic episodes at 6 months (Carter MC et al).
- Atopic dermatitis: A double-blind, placebo-controlled study in 62 pediatric patients with severe atopic dermatitis found Omalizumab to have a statistically significant difference compared to placebo in the Scoring Atopic Dermatitis [SCORAD] index at 24 weeks; however, the clinical significance of this is unknown (Chan S et al). Another randomized, double-blind, placebo-controlled study found that while Omalizumab reduced serum levels of free IgE and decreased surface-bound IgE, it did not significantly alter several measures of clinical disease activity (i.e., atopy patch test results in single patients) (Heil PM et al). The 2023 American Academy of Dermatology atopic dermatitis guidelines state that there are insufficient data to make a recommendation regarding the use of Omalizumab.

Appendix E: Age ≥ 12 Years: Asthma Dosing Based on Pre-treatment IgE and Body Weight[†]

Pre-treatment serum IgE IU/mL	Dosing Frequency	Body Weight			
		30-60 kg	> 60-70 kg	> 70-90 kg	> 90-15 kg
$\geq 30-100$	Q 4 weeks	150 mg	150 mg	150 mg	300 mg
> 100-200		300 mg	300 mg	300 mg	225 mg
> 200-300		300 mg	225 mg	225 mg	300 mg
> 300-400	Q 2 weeks	225 mg	225 mg	300 mg	
> 400-500		300 mg	300 mg	375 mg	
> 500-600		300 mg	375 mg	Insufficient Data to Recommend a Dose	
> 600-700		375 mg			

[†]The manufacturer recommends dose adjustments for significant body weight changes during treatment.

Appendix F: Age 6 to < 12 Years: Asthma Dosing Based on Pre-treatment IgE and Body Weight[†]

Pre-treatment serum IgE IU/mL	Dosing Frequency	Body Weight									
		20-25 kg	> 25-30 kg	> 30-40 kg	> 40-50 kg	> 50-60 kg	> 60-70 kg	> 70-80 kg	> 80-90 kg	> 90-125 kg	> 125-150 kg
$\geq 30-100$	Q 4 weeks	75	75	75	150	150	150	150	150	300	300
> 100-200		150	150	150	300	300	300	300	300	225	300
> 200-300		150	150	225	300	300	225	225	225	300	375
> 300-400		225	225	300	225	225	225	300	300		
> 400-500		225	300	225	225	300	300	375	375		

CLINICAL POLICY
Omalizumab, Omalizumab-igec



> 500-600		300	300	225	300	300	375	
> 600-700		300	225	225	300	375		
> 700-800	Q 2 weeks	225	225	300	375			
> 800-900		225	225	300	375			
> 900-1,000		225	300	375				
> 1,000-1,100		225	300	375				
> 1,100-1,200		300	300					
> 1,200-1,300		300	375					

Insufficient Data to Recommend a Dose

[†]The manufacturer recommends dose adjustments for significant body weight changes during treatment.

Appendix G: Age \geq 18 Years: CRSwNP Dosing Based on Pre-treatment IgE and Body Weight[†]

Pre-treatment serum IgE IU/mL	Dosing Frequency	Body Weight							
		> 30-40 kg	> 40-50 kg	> 50-60 kg	> 60-70 kg	> 70-80 kg	> 80-90 kg	> 90-125 kg	> 125-150 kg
≥ 30-100	Q 4 weeks	75	150	150	150	150	150	300	300
> 100-200		150	300	300	300	300	300	450	600
> 200-300		225	300	300	450	450	450	600	375
> 300-400		300	450	450	450	600	600	450	525
> 400-500		450	450	600	600	375	375	525	600
> 500-600		450	600	600	375	450	450	600	
> 600-700		450	600	375	450	450	525		
> 700-800	Q 2 weeks	300	375	450	450	525	600		
> 800-900		300	375	450	525	600			
> 900-1,000		375	450	525	600				
> 1,000-1,100		375	450	600					
> 1,100-1,200		450	525	600					
> 1,200-1,300		450	525						
> 1,300-1,500		525	600						

Insufficient Data to Recommend a Dose

[†]The manufacturer recommends dose adjustments for significant body weight changes during treatment.

Appendix H: Age \geq 1 Year: IgE-Mediated Food Allergy Dosing Based on Pre-treatment IgE and Body Weight[†]

Pre-treatment serum IgE IU/mL	Dosing Frequency	Body Weight (continued on next table)						
		≥ 10-12 kg	> 12-15 kg	> 15-20 kg	> 20-25 kg	> 25-30 kg	> 30-40 kg	> 40-50 kg
≥ 30-100	Q 4 weeks	75	75	75	75	75	75	150
> 100-200		75	75	75	150	150	150	300
> 200-300		75	75	150	150	150	225	300
> 300-400		150	150	150	225	225	300	450
> 400-500		150	150	225	225	300	450	450
> 500-600		150	150	225	300	300	450	600
> 600-700		150	150	225	300	225	450	600
> 700-800	Q 2 weeks	150	150	150	225	225	300	375
> 800-900		150	150	150	225	225	300	375
> 900-1,000		150	150	225	225	300	375	450
> 1,000-1,100		150	150	225	225	300	375	450
> 1,100-1,200		150	150	225	300	300	450	525
> 1,200-1,300		150	225	225	300	375	450	525
> 1,300-1,500		150	225	300	300	375	525	600
> 1,500-1,850		*	225	300	375	450	600	*

[†]The manufacturer recommends dose adjustments for significant body weight changes during treatment.

* Insufficient data to recommend a dose

Pre-treatment serum IgE IU/mL	Dosing Frequency	Body Weight (continued from previous table)					
		> 50- 60 kg	> 60- 70 kg	> 70-80 kg	> 80- 90 kg	> 90- 125 kg	> 125- 150 kg
≥ 30-100	Q 4 weeks	150	150	150	150	300	300
> 100-200		300	300	300	300	450	600
> 200-300		300	450	450	450	600	375
> 300-400		450	450	600	600	450	525
> 400-500		600	600	375	375	525	600
> 500-600		600	375	450	450	600	
> 600-700		375	450	450	525		
> 700-800	Q 2 weeks	450	450	525	600		
> 800-900		450	525	600			
> 900-1,000		525	600				
> 1,000-1,100		600					
> 1,100-1,200		600					
> 1,200-1,300							
> 1,300-1,500							
> 1,500-1,850							

[†]The manufacturer recommends dose adjustments for significant body weight changes during treatment.

Appendix I: Immunotherapy-related Pruritus

- Immunotherapy refers to immune checkpoint inhibitors. Immune checkpoint inhibitors comprise a class of agents that target immune cell checkpoints, such as programmed cell death-1 (PD-1; e.g., Opdivo®, Keytruda®) and PD-1 ligand (PD-L1; e.g., Tecentriq®, Bavencio®, Imfinzi®), as well as cytotoxic T-lymphocyte-associated antigen 4 (e.g., Yervoy®, Imjudo®).
- NCCN grading of pruritus
 - G1: Mild or localized
 - G2: Moderate. Intense or widespread; intermittent; skin changes from scratching (e.g., edema, papulation, excoriations, lichenification, oozing/crusts); limiting instrumental activities of daily living (ADLs)
 - G3: Severe. Intense or widespread; constant; limiting self-care ADLs or sleep

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
Asthma*	75 to 375 mg SC every 2 or 4 weeks based on serum total IgE level (IU/mL) measured before the start of treatment, and body weight (kg). Adjust doses for significant changes in body weight during treatment Xolair and Omlyclo are not approved for use in patients weighing more than 150 kg (see Appendix E and F) Do not administer more than 150 mg (contents of one vial) per injection site. Divide doses of more than 150 mg amongst two or more injection sites	375 mg/2 weeks

Indication	Dosing Regimen	Maximum Dose
CSU	150 mg or 300 mg SC every 4 weeks	300 mg/4 weeks
CRSwNP*	75 to 600 mg SC every 2 or 4 weeks based on serum total IgE level (IU/mL) measured before the start of treatment, and body weight (kg). Adjust doses for significant changes in body weight during treatment	600 mg/2 weeks
IgE-mediated food allergy*	75 mg to 600 mg SC every 2 or 4 weeks based on serum total IgE level (IU/mL) measured before the start of treatment and body weight (kg). Adjust doses for significant changes in body weight during treatment	600 mg/2 weeks

**For patients with a combination of either asthma, CRSwNP, and/or IgE-mediated food allergy, dosing determination should be based on the primary diagnosis for which Xolair/Omlyclo is being prescribed.*

VI. Product Availability*

Drug Name	Availability
Omalizumab (Xolair)	<ul style="list-style-type: none"> Single-dose vial: 150 mg Single-dose prefilled syringes: 75 mg/0.5 mL, 150 mg/mL, 300 mg/2 mL Single-dose prefilled autoinjectors: 75 mg/0.5 mL, 150 mg/mL, 300 mg/2 mL
Omalizumab-igec (Omlyclo)	Single-dose prefilled syringes: 75 mg/0.5 mL, 150 mg/mL

VII. References

1. Xolair Prescribing Information. Irvine, CA: Spectrum Pharmaceuticals, Inc.; February 2024. Available at: https://www.gene.com/download/pdf/xolair_prescribing.pdf. Accessed March 25, 2025.
2. Omlyclo Prescribing Information. Jersey City, NJ: Celltrion USA, Inc.; March 2025. Available at: https://www.accessdata.fda.gov/drugsatfda_docs/label/2025/761399s000lbl.pdf. Accessed March 25, 2025.
3. National Asthma Education and Prevention Program: Expert panel report III: Guidelines for the diagnosis and management of asthma. Bethesda, MD: National Heart, Lung, and Blood Institute, 2007. (NIH publication no. 08-4051). Available at <http://www.nhlbi.nih.gov/health-pro/guidelines/current/asthma-guidelines>. Accessed November 14, 2024.
4. Cloutier MM, Dixon AE, Krishnan JA, et al. Managing asthma in adolescents and adults 2020: asthma guideline update from the National Asthma Education and Prevention Program. JAMA. 2020; 324: 2301-2317.
5. Bernstein JA, Lang DM, Khan DA, et al. The diagnosis and management of acute and chronic urticaria: 2014 update. J Allergy Clin Immunol. 2014; 133(5): 1270-1277.
6. Zuberbier T, Aberer W, Asero R, et al. The EAACI/GA(2) LEN/EDF/WAO guideline for the definition, classification, diagnosis, and management of urticarial (2018 revision). Allergy. 2018; 73: 1393-1414.
7. Fine LM, Bernstein JA. Guideline of chronic urticaria beyond. Allergy Asthma Immunol Res. 2016 September; 8(5): 396-403.
8. Micromedex® Healthcare Series [Internet database]. Greenwood Village, CO: Thomson Healthcare. Updated periodically. Accessed November 14, 2024.

9. Global Initiative for Asthma. Global strategy for asthma management and prevention (2024 update). Available from: www.ginasthma.org. Accessed November 14, 2024.
10. Global Initiative for Asthma. Difficult-to-treat and severe asthma in adolescent and adult patients – diagnosis and management, v5.0 November 2024. Available at: www.ginasthma.org. Accessed November 14, 2024.
11. Rosenfeld RM, Piccirillo JF, Chandrasekhar SS, et al. Clinical practice guideline (update): adult sinusitis. *Otolaryngology–Head and Neck Surgery* 2015, Vol. 152(2S) S1–S39.
12. Peters AT, Spector S, Hsu J, et al. Diagnosis and management of rhinosinusitis: a practice parameter update. *Ann Allergy Asthma Immunol* 2014; 113:347-85.
13. Fokkens WJ, Lund V, Bachert C, et al. EUFOREA consensus on biologics for CRSwNP with or without asthma. doi: 10.1111/all.13875.
14. ClinicalTrials.gov. A clinical trial of omalizumab in participants with chronic rhinosinusitis with nasal polyps (POLYP 1). Available at: <https://clinicaltrials.gov/ct2/show/NCT03280550>. Accessed November 14, 2024.
15. ClinicalTrials.gov. A clinical trial of omalizumab in participants with chronic rhinosinusitis with nasal polyps (POLYP 2). Available at: <https://clinicaltrials.gov/ct2/show/NCT03280537>. Accessed November 14, 2024.
16. Carter MC, Maric I, Brittain EH, et al. A randomized double-blind, placebo-controlled study of omalizumab for idiopathic anaphylaxis. *J Allergy Clin Immunol*. 2021; 147(3): 1004-1010.e2.
17. Han JK, Bosson JV, Cho SH, et al. Multidisciplinary consensus on a stepwise treatment algorithm for management of chronic rhinosinusitis with nasal polyps. *Int Forum Allergy Rhinol*. 2021;1-10. Available at: <https://onlinelibrary.wiley.com/doi/10.1002/alr.22851>. Accessed November 14, 2024.
18. Rank MA, Chu DK, Bognanni A, et al. The Joint Task Force on practice parameters GRADE guidelines for the medical management of chronic rhinosinusitis with nasal polyposis. *J Allergy Clin Immunol*. 2023;151(2):386-398.
19. Wood RA, Togias A, Sicherer SH, et al. Omalizumab for the treatment of multiple food allergies. *N Engl J Med*. 2024;390(10):889-899.
20. NIAID-Sponsored Expert Panel, Boyce JA, Assa'ad A, et al. Guidelines for the diagnosis and management of food allergy in the United States: report of the NIAID-sponsored expert panel. *J Allergy Clin Immunol*. 2010;126(6 Suppl):S1-S58.
21. Sampson HA, Aceves S, Bock SA, et al. Food allergy: a practice parameter update-2014. *J Allergy Clin Immunol*. 2014;134(5):1016-25.
22. Santos AF, Riggioni C, Agache I, et al. EAACI guidelines on the diagnosis of IgE-mediated food allergy. *Allergy*. 2023;78(12):3057-3076.
23. Anagnostou A, Bird JA, Chinthurajah S, et al. The use and implementation of omalizumab as food allergy treatment: Consensus-based guidance and Work Group Report of the Adverse Reactions to Foods Committee of the American Academy of Allergy, Asthma & Immunology. *J Allergy Clin Immunol*. 2024. Available at: [https://www.jacionline.org/article/S0091-6749\(24\)01177-1/abstract](https://www.jacionline.org/article/S0091-6749(24)01177-1/abstract). Accessed November 26, 2024.
24. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at www.nccn.org. Accessed March 25, 2025.
25. National Comprehensive Cancer Network. Systemic Mastocytosis Version 3.2024. Available at https://www.nccn.org/professionals/physician_gls/pdf/mastocytosis.pdf.

Accessed March 25, 2025.

26. National Comprehensive Cancer Network. Management of Immunotherapy-Related Toxicities Version 2.2024. Available at https://www.nccn.org/professionals/physician_gls/pdf/immunotherapy.pdf. Accessed March 25, 2025.
27. Cardet JC, Akin C, Lee MJ. Mastocytosis: update on pharmacotherapy and future directions. *Expert Opin Pharmacother.* 2013;14(15):2033-2045.
28. Heil PM, Maurer D, Klein B, et al. Omalizumab therapy in atopic dermatitis: depletion of IgE does not improve the clinical course - a randomized, placebo-controlled, and double-blind pilot study. *J Dtsch Dermatol Ges.* 2010;8(12):990.
29. Chan S, Cornelius V, Cro S, et al. Treatment effect of omalizumab on severe pediatric atopic dermatitis: the ADAPT randomized clinical trial. *JAMA Pediatr.* 2019;174(1):29-37.
30. Davis DMR, Drucker AM, Alikhan A, et al. Guidelines of care for the management of atopic dermatitis in adults with phototherapy and systemic therapies. *J Am Acad Dermatol.* 2024; e43-e56.

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to- date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J2357	Injection, omalizumab, 5 mg

Reviews, Revisions, and Approvals	Date
Policy created	09/2025