

Clinical Policy: Holter Monitors

Reference Number: CP.MP.113

Effective Date: 05/18

Last Review Date: 04/18

[Coding Implications](#)

[Revision Log](#)

Description

Ambulatory electrocardiogram (ECG) monitoring provides a view of cardiac activity over an extended period of time. Holter monitoring, or continuous ambulatory ECG monitoring, for 24 to 48 hours is most practical as the initial monitor for members with daily or near daily symptoms, as well as for assessing the efficacy of medication and other treatments for cardiac arrhythmias.

Policy/Criteria

- I. It is the policy of PA Health & Wellness that Holter monitoring is **medically necessary** for adult members who require 24 to 48 hours of cardiac activity monitoring with any of the following symptoms or indications:
 - A. Evaluation of any of these unexplained indications: syncope, near-syncope, episodic dizziness, recurrent palpitations, episodic shortness of breath or chest pain;
 - B. Evaluation of neurological events when transient atrial fibrillation or flutter is suspected;
 - C. Evaluation of syncope, near-syncope, episodic dizziness, or palpitation in whom a probable cause other than an arrhythmia has been identified but in whom symptoms persist despite treatment of this other cause;
 - D. Evaluation of patients with cardiomyopathy, or a first-degree relative with arrhythmogenic right ventricular cardiomyopathy;
 - E. Evaluation of possible or documented prolonged QT syndromes;
 - F. To screen for asymptomatic arrhythmia in a patient with Brugada syndrome;
 - G. Assessment of efficacy of medication for arrhythmia treatment when baseline arrhythmia frequency is reproducible and of sufficient frequency to permit analysis;
 - H. Detection of proarrhythmic responses to antiarrhythmic therapy in patients at high risk;
 - I. Assessment of the function of pacemakers or implantable cardioverter defibrillators (ICD) with frequent palpitations, syncope, or near-syncope, and to assist in programming of enhanced features;
 - J. Evaluation of suspected pacemaker or ICD component failure or malfunction when device interrogation is inconclusive;
 - K. Assessment of efficacy of adjunctive medications in patients receiving frequent ICD therapy;
 - L. Assessment of suspected variant angina.

- II. It is the policy of PA Health & Wellness that Holter monitoring is **medically necessary** for pediatric members ≤ 18 years old who require 24 to 48 hours of cardiac activity monitoring with any of the following symptoms or indications:
 - A. Evaluation of syncope, near-syncope, or dizziness in members with identified cardiac disease, previously documented arrhythmia, or pacemaker dependency;
 - B. Evaluation of syncope or near-syncope associated with exertion when cause is not established;
 - C. Evaluation of unexplained syncope, near-syncope, or sustained palpitation when there is no overt clinical evidence of heart disease;

CLINICAL POLICY

Holter Monitors

- D. Assessment of efficacy of medications for arrhythmia following initiation of treatment or during rapid somatic growth;
- E. Evaluation of patients with cardiomyopathy, with or a first-degree relative with arrhythmogenic right ventricular cardiomyopathy;
- F. Evaluation of possible or documented prolonged QT syndromes;
- G. Evaluation of palpitation in a member with prior surgery for congenital heart disease and significant residual hemodynamic abnormalities;
- H. Evaluation of asymptomatic congenital complete atrioventricular (AV) block, non-paced;
- I. Evaluation of cardiac rhythm after transient AV block associated with heart surgery or catheter ablation;
- J. Evaluation of rate-responsive or physiological pacing function in symptomatic patients.

III. It is the policy of PA Health & Wellness that Holter monitoring for any other indication not included in this policy is **not medically necessary** because efficacy has not been established.

Background

The most common use of ambulatory ECG monitoring is the evaluation and diagnosis of cardiac arrhythmias or conduction abnormalities. The device continuously monitors the heart's electrical activity for a period of 24 to 48 hours. The member has a self-activated event marker which identifies when they are experiencing symptoms such as palpitations, syncope/near-syncope, dizziness, shortness of breath, chest pain, or episodic fatigue. This is especially helpful in members who experience symptoms too infrequent to be caught on a standard ECG.

The recorded data are analyzed with the event markers to determine if the symptoms are related to an arrhythmia. There are four outcomes this analysis could provide. Useful findings include the simultaneous documentation of a cardiac arrhythmia capable of producing the noted symptoms, which can lead to directed therapy for the arrhythmia; and symptoms that occur without arrhythmia, demonstrating symptoms are not related to an arrhythmia. Of equivocal value, the findings may show that a cardiac arrhythmia is present but no symptoms were present during the recording, indicating the arrhythmia may or may not be related to the symptoms. Lastly, if there were no symptoms during the recording and there were no arrhythmias identified, the recording is not useful.

Ambulatory ECG is also helpful in assessing the efficacy of antiarrhythmic therapy. It is noninvasive, provides quantitative data, and permits correlation of symptoms with ECG phenomena. It does have some limitations in regard to its use as a therapeutic guide, which should be taken into consideration. Additionally, ambulatory ECG monitoring is useful in assessing pacemakers and ICDs, as it can evaluate symptoms of palpitations, syncope, or near-syncope to assess device function; assist in the programming of enhanced features; evaluate suspected component failure or a malfunctioning device; and assess concomitant pharmacological therapy for members receiving frequent ICD therapy.

Due to the advancement of technological capabilities in ambulatory ECG assessment, it can provide accurate and clinically meaningful information about myocardial ischemia in patients with coronary disease. The most commonly encountered ambulatory ECG sign of ischemia is

CLINICAL POLICY

Holter Monitors

ST-segment depression and, while this is an important finding, it is important to note that ST-segment changes and other repolarization abnormalities can occur for reasons other than ischemia. These conditions must be considered when evaluating the predictive value of ST-segment changes in each specific member. Furthermore, ambulatory ECG can be beneficial in members suspected of having variant angina. Periods of ST-segment elevation indicative of transmural ischemia can be identified in those with variant angina or high-grade proximal stenosis.

In the pediatric population, ambulatory ECG can be used for the same indications as for adults, in addition to a number of pediatric-specific concerns. Monitoring in children with heart disease, with or without symptoms, is used to observe the evolution of disease processes, identify medication dose changes required due to growth, and identify the progressive onset of late arrhythmias after surgery for congenital heart defects. Likewise, this monitoring is beneficial in pediatric members with hypertrophic or dilated cardiomyopathies or known or suspected prolonged QT syndromes. Ambulatory ECG can also be used to evaluate asymptomatic pediatric members with congenital complete AV block in order to identify those at increased risk for sudden arrhythmic events who may benefit from prophylactic pacemaker implantation.

Coding Implications

This clinical policy references Current Procedural Terminology (CPT®). CPT® is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2017, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

CPT® Codes	Description
93224	External electrocardiographic recording up to 48 hours by continuous rhythm recording and storage; includes recording, scanning analysis with report, physician review and interpretation
93225	External electrocardiographic recording up to 48 hours by continuous rhythm recording and storage; recording (includes connection, recording, and disconnection)
93226	External electrocardiographic recording up to 48 hours by continuous rhythm recording and storage; scanning analysis with report
93227	External electrocardiographic recording up to 48 hours by continuous rhythm recording and storage; review and interpretation by a physician or other qualified health care professional

ICD-10-CM Diagnosis Codes that Support Coverage Criteria

ICD-10-CM Code	Description
G45.9	Transient cerebral ischemic attack, unspecified

CLINICAL POLICY

Holter Monitors

ICD-10-CM Code	Description
G71.0	Muscular dystrophy
G99.0	Autonomic neuropathy in diseases classified elsewhere
I20.0-I20.9	Angina pectoris
I24.0-I24.9	Other acute ischemic heart diseases
I25.10	Atherosclerotic heart disease of native coronary artery without angina pectoris
I34.0-I34.9	Nonrheumatic mitral valve disorders
I35.0-I35.9	Nonrheumatic aortic valve disorders
I36.0-I36.9	Nonrheumatic tricuspid valve disorders
I37.0-I37.9	Nonrheumatic pulmonary valve disorders
I42.0	Dilated cardiomyopathy
I42.1	Obstructive hypertrophic cardiomyopathy
I42.2	Other hypertrophic cardiomyopathy
I42.8	Other cardiomyopathies
I42.9	Cardiomyopathy, unspecified
I44.0-I44.7	Atrioventricular and left bundle-branch block
I45.0-I45.9	Other conduction disorders
I46.2-I46.9	Cardiac arrest
I47.0-I47.9	Paroxysmal tachycardia
I48.0-I48.92	Atrial fibrillation and flutter
I49.0-I49.9	Other cardiac arrhythmias
I50.1-I50.9	Heart failure
I51.7	Cardiomegaly
I63.00-I63.9	Cerebral infarction
I67.841-I67.848	Cerebral vasospasm and vasoconstriction
Q20.0-Q20.9	Congenital malformations of cardiac chambers and connections
Q21.0-Q21.9	Congenital malformations of cardiac septa
Q22.0-Q22.9	Congenital malformations of pulmonary and tricuspid valves
Q23.0-Q23.9	Congenital malformations of aortic and mitral valves
Q24.0-Q24.9	Other congenital malformations of heart
Q25.0-Q25.9	Congenital malformations of great arteries
R00.0-R00.9	Abnormalities of heart beat
R06.00-R06.09	Shortness of breath
R07.2	Precordial pain
R07.89	Other chest pain
R07.9	Chest pain, unspecified
R42	Dizziness and giddiness
R53.81-R53.83	Malaise and fatigue
R55	Syncope and collapse
R94.31	Abnormal electrocardiogram
Z48.812	Encounter for surgical aftercare following surgery on the circulatory system

CLINICAL POLICY

Holter Monitors

ICD-10-CM Code	Description
Z82.41	Family history of sudden cardiac death
Z87.74	Personal history of (corrected) congenital malformations of heart and circulatory systems
Z94.1	Heart transplant status
Z95.0	Presence of cardiac pacemaker
Z95.810	Presence of automatic (implantable) cardiac defibrillator

Reviews, Revisions, and Approvals	Date	Approval Date
Policy developed	04/18	06/18

References

1. Crawford MH, Bernstein SJ, Deedwania PC, DiMarco JP, Ferrick KJ, Garson A Jr, Green LA, Greene HL, Silka MJ, Stone PH, Tracy CM. ACC/AHA guidelines for ambulatory electrocardiography: executive summary and recommendations: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee to Revise the Guidelines for Ambulatory Electrocardiography). *Circulation*. 1999;100:886-893
2. Gersh BJ, Maron BJ, Bonow RO, et al. 2011 ACCF/AHA guideline for the diagnosis and treatment of hypertrophic cardiomyopathy: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *Circulation*. 2011;124(24):e783.
3. Gray B, Kirby A, Kabunga P, et al. Twelve-lead ambulatory electrocardiographic monitoring in Brugada syndrome: Potential diagnostic and prognostic implications. *Heart Rhythm*. 2017 Jun;14(6):866-874. doi: 10.1016/j.hrthm.2017.02.026.
4. Groeneweg JA, Bhonsale A, James CA, et al. Clinical Presentation, Long-Term Follow-Up, and Outcomes of 1001 Arrhythmogenic Right Ventricular Dysplasia/Cardiomyopathy Patients and Family Members. *Circ Cardiovasc Genet* 2015; 8:437.
5. Podrid PJ. Ambulatory ECG monitoring. In: UpToDate, Zimetbaum PJ (Ed), UpToDate, Waltham, MA. Accessed 07/21/16.
6. Sen-Chowdhry S, Lowe MD, Sporton SC, McKenna WJ. Arrhythmogenic right ventricular cardiomyopathy: clinical presentation, diagnosis, and management. *Am J Med* 2004; 117:685.