

Clinical Policy: Balloon Sinus Ostial Dilation for Treatment of Chronic Sinusitis

Reference Number: PA.CP.MP.119

Effective Date: 01/18

Last Review Date: 07/17

[Coding Implications](#)

[Revision Log](#)

Description

Sinuplasty, also known as balloon catheter sinusotomy and balloon sinus ostial dilation, is a minimally invasive technique intended to dilate the sinus ostia in patients with chronic sinusitis. The Relieva Balloon Sinuplasty System by Acclarant Inc. received FDA approval in April of 2005. It is a set of single-use, endoscopic, catheter-based instruments for minimally invasive sinus surgery. Per the FDA, the Relieva Sinus Balloon Dilation Catheter is intended to provide a means to dilate the sinus ostia and spaces within the paranasal sinus cavities for diagnostic and therapeutic procedures to open passages and to restore normal drainage. Balloon sinuplasty is proposed to treat patients with chronic sinusitis who have exhausted less aggressive treatment options.

Policy/Criteria

- I. It is the policy of Pennsylvania Health and Wellness[®] (PHW) that balloon sinuplasty is **medically necessary** for individuals with chronic rhinosinusitis (CRS) in order to relieve obstruction of the maxillary, sphenoid, and frontal sinus ostia, either alone or in combination with standard endoscopic sinus surgery techniques, when all of the following are met:
 - A. Documentation that the inflammation of the paranasal sinuses has persisted for 12 weeks or longer;
 - B. If > 18 years of age, meets both of the following (1 and 2):
 1. Has at least one of the following:
 - a. Anterior or posterior mucopurulent nasal discharge;
 - b. Nasal obstruction;
 - c. Facial-pain-pressure-fullness;
 - d. Decreased or lost sense of smell;
 2. Has at least one finding of chronic sinusitis by computed tomography (CT) scan (a or b):
 - a. Polyps in nasal cavity or the middle meatus, and/or opacification;
 - b. Radiographic imaging showing inflammation of the paranasal sinuses.
 - C. If ≤ 18 years of age, meets both of the following (1 and 2):
 1. Has at least two of the following:
 - a. Purulent rhinorrhea;
 - b. Nasal obstruction;
 - c. Facial pressure/pain;
 - d. Cough;
 2. Mucosal changes within the osteomeatal complex and/or sinuses, by CT scan;
 - D. Continued symptoms after medical therapy consisting of both of the following (1 and 2):
 1. Antibiotic therapy meeting one of the following (a or b):
 - a. Antibiotic therapy guided by culture and sensitivity for ≥ 3 weeks;
 - b. Beta-lactamase resistant antibiotic for ≥ 3 weeks (e.g., amoxicillin [recommended], amoxicillin-clavulanate, trimethoprim-sulfisoxazole, cefuroxime).
 2. Intranasal corticosteroids for ≥ 4 weeks.

- II.** It is the policy of PHW that balloon sinuplasty is **not medically necessary** in any of the following situations:
- A. For the treatment of ethmoid disease;
 - B. Extensive previous surgery with significant osteoneogenesis.

Background

Chronic rhinosinusitis (CRS) is defined as an inflammatory condition involving the paranasal sinuses and linings of the nasal passages, which persists for 12 weeks or longer. Symptoms of CRS include anterior and/or posterior mucopurulent drainage, nasal obstruction, facial pain, pressure, and/or fullness and decreased sense of smell. The goal of medical therapy (e.g., antibiotics, nasal irrigation, topical corticosteroids) is directed toward facilitating the drainage of sinus secretions and treatment to eradicate the offending pathogens. Surgical intervention may be indicated when the patient requires more than three courses of antibiotics for sinusitis within a 12-month period along with evidence of abnormalities of the sinuses or ostiomeatal complex (OMC) on nasal endoscopy or CT imaging. The goal of functional endoscopic sinus surgery or FESS is to restore physiologic sinus ventilation and drainage, which allows for the gradual resolution of mucosal disease. Balloon dilation is a less invasive alternative to endoscopic sinus surgery in the management of chronic sinusitis.

The goal of balloon sinuplasty is to restore normal sinus drainage by enlarging passages of the sinus ostia and spaces within the paranasal sinus cavities, without cutting bone or removing tissue. Per the manufacturer of the Relieva Sinus Balloon Dilation Catheter, the procedure is performed under fluoroscopic guidance using endoscopic technique, by an otolaryngologist trained in the use of the Balloon Sinuplasty System. The initial sinus access is achieved by the introduction of a guide catheter into the target sinus. A flexible guidewire is then introduced through the guide catheter and gently advanced into the target sinus. The balloon catheter tracks smoothly over the guide wire and positioned across the blocked ostium. After the position of the balloon catheter is confirmed, it is gradually inflated to gently restructure the blocked ostium. The system is removed leaving the ostium open and allowing the return of normal sinus drainage and function with little to no disruption to the mucosal lining. Balloon sinuplasty may be performed in conjunction with endoscopic sinus surgery and used as an assistive procedure for sinus tissue biopsy or culturing, sinus lavage, drainage, or antibiotic irrigation.

Studies evaluating balloon sinuplasty are limited and include a prospective randomized trial, cohort studies, case series, observational and retrospective studies. Most studies were small and long term studies are lacking. However, the available studies suggest that balloon sinuplasty for chronic sinusitis refractory to medical therapy is safe and efficacious. The data show that balloon sinuplasty can successfully dilate the sinus ostia and relieve symptoms of chronic sinusitis. In addition, the use of balloon sinuplasty is supported as a treatment option by the professional societies noted below.

There is limited evidence regarding balloon sinuplasty in the pediatric population. However, two small studies have found positive effects of balloon sinuplasty in pediatric patients with CRS failing to respond to medical therapy^{16,18}. Additionally, one study found that balloon sinuplasty led to improved outcomes after failure to respond adequately to adenoidectomy¹⁴.

Guideline Recommendations

Both the American Academy of Otolaryngology (AAO)-Head and Neck Surgery and the American Rhinologic Society (ARS) position statements on dilation of sinuses, any method (e.g., balloon, etc.), state “sinus ostial dilation (e.g., balloon ostial dilation) is an appropriate therapeutic option for selected patients with sinusitis. This approach may be used alone to dilate a sinus ostium (frontal, maxillary, or sphenoid) or in conjunction with other instruments (e.g., microdebrider, forceps). The final decision regarding use of techniques or instrumentation for sinus surgery is the responsibility of the attending surgeon.” For pediatric patients, the AAO did not reach consensus on whether balloon sinuplasty should be recommended for the treatment of CRS; however, near consensus was reached regarding the safety of balloon sinuplasty.

Coding Implications

This clinical policy references Current Procedural Terminology (CPT®). CPT® is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2017, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

CPT® Codes	Description
31295	Nasal/sinus endoscopy, surgical; with dilation of maxillary sinus ostium (e.g., balloon dilation), transnasal or via canine fossa
31296	Nasal/sinus endoscopy, surgical; with dilation of frontal sinus ostium (e.g., balloon dilation)
31297	Nasal/sinus endoscopy, surgical; with dilation of sphenoid sinus ostium (e.g., balloon dilation)

HCPCS Codes	Description
N/A	

ICD-10-CM Diagnosis Codes that Support Coverage Criteria

ICD-10-CM Code	Description
J32.0	Chronic maxillary sinusitis
J32.1	Chronic frontal sinusitis
J32.3	Chronic sphenoidal sinusitis
J32.8	Other chronic sinusitis
J32.9	Chronic sinusitis, unspecified

Reviews, Revisions, and Approvals	Date	Approval Date
Changed 2 nd listing of CPT code 31296 to 31297 in order to correctly match the code description.	07/17	

References

1. Abreu CB, Balsalobre L, Pascoto GR, et al. Effectiveness of balloon sinuplasty in patients with chronic rhinosinusitis without polyposis. *Braz J Otorhinolaryngol.* 2014 Nov-Dec;80(6):470-5.
2. American Academy of Otolaryngology-Head and Neck Surgery. Position statement on Dilation of sinuses, any method (e.g., balloon, etc.). Last updated 1/2014. Available at: <http://www.entnet.org/content/position-statement-dilation-sinuses-any-method-eg-balloon-etc>
3. American Rhinologic Society (ARS). Ostial Balloon Dilation Position Statement. Revised 1/2015. Available at: http://www.american-rhinologic.org/position_balloon_dilation
4. Bizaki AJ, Taulu R, Numminen J, Rautiainen M. Quality of life after endoscopic sinus surgery or balloon sinuplasty: a randomized clinical study. *Rhinology.* 2014 Dec;52(4):300-5.
5. Bolger WE, Brown CL, Church CA, et al.: Safety & Outcomes of Balloon Catheter Sinusotomy: A Multicenter 24 week Analysis in 115 Patients (The CLEAR Study). *Otolaryngology-Head & Neck Surgery.* 2007 Jul;137(1):10-20.
6. Brietzke SE, Shin JJ, Choi S et al. Clinical consensus statement: pediatric chronic rhinosinusitis by the American Academy of Otolaryngology- Head and Neck Surgery Foundation. *Otolaryngology– Head and Neck Surgery.* 2014, Vol. 151(4) 542–553
7. Brown CL, Bolger WE. Safety and feasibility of balloon catheter dilation of paranasal sinus ostia: a preliminary investigation. *Ann Otol Rhinol Laryngol.* 2006 Apr;115(4):293-9;
8. ElBadawey MR, Alwaa A, ElTaher M, Carrie S. Quality of life benefit after endoscopic frontal sinus surgery. *Am J Rhinol Allergy.* 2014 Sep-Oct;28(5):428-32.
9. Hayes Health Technology Brief. Relieva Balloon Sinuplasty (Acclarent Inc.) for Chronic Sinusitis in Adults. Oct 2012. Archived Nov. 2015
10. Kim E, Cutler JL. Balloon Dilatation of the Paranasal Sinuses: A Tool in Sinus Surgery. *Otolaryngologic Clinics of North America - Volume 42, Issue 5 (October 2009).*
11. Koskinen A, Myller J, Mattila P, et al. Long-term follow-up after ESS and balloon sinuplasty: Comparison of symptom reduction and patient satisfaction. *Acta Otolaryngol.* 2016 May;136(5):532-6.
12. Orlandi RR, Kingdom TT, Hwang PH, et al. International Consensus Statement on Allergy and Rhinology: Rhinosinusitis. *Int Forum Allergy Rhinol.* 2016 Feb; 6 Suppl 1: S22-209.
13. Ramadan HH. Safety and feasibility of balloon sinuplasty for treatment of chronic rhinosinusitis in children. *Ann Otol Rhinol Laryngol,* 01-MAR-2009; 118(3): 161-5.
14. Ramadan HH, Terrell AM. Balloon catheter sinuplasty and adenoidectomy in children with chronic rhinosinusitis. *Ann Otol Rhinol Laryngol.* 2010; 119:578-582.
15. Rosenfeld RM, Piccirillo JF, Chandrasekhar SS, et al. Clinical Practice Guideline (Update): Adult Sinusitis by the American Academy of Otolaryngology- Head and Neck Surgery Foundation. *Otolaryngology– Head and Neck Surgery.* 2015 152(2S) S1-S39.
16. Setzen G, Ferguson BJ, Han JK, et al. Clinical Consensus Statement: Appropriate Use of Computed Tomography for Paranasal Sinus Disease by the American Academy of

- Otolaryngology- Head and Neck Surgery Foundation. Otolaryngology– Head and Neck Surgery. 2012 Nov;147(5):808-16.
17. Thottam PJ, Metz CM, Kieu MC, et al. Functional Endoscopic Sinus Surgery Versus Balloon Sinuplasty with Ethmoidectomy: A 2-year Analysis in Pediatric Chronic Rhinosinusitis. *Indian J Otolaryngol Head Neck Surg.* 2016 Sep;68(3):300-6.
 18. Tomazic PV, Stammberger H, Braun H, et al. Feasibility of balloon sinuplasty in patients with chronic rhinosinusitis: the Graz experience. *Rhinology.* 2013 Jun;51(2):120-7.
 19. Wang F, Song Y, Zhang X, Tan G. Sinus balloon catheter dilation in pediatric chronic rhinosinusitis resistant to medical therapy. *JAMA Otolaryngol Head Neck Surg.* 2015 Jun;141(6):526-31