

## Clinical Policy: Nonmyeloablative Allogeneic Stem Cell Transplants

Reference Number: PA.CP.MP.141

Effective Date: 01/18

Date of Last Revision: 03/2024

[Coding Implications](#)

[Revision Log](#)

### Description

Allogeneic hematopoietic stem cell transplants that do not destroy all of the hematopoietic cells in the bone marrow are termed reduced-intensity or nonmyeloablative conditioning regimens. Although there are no clear definitions, reduced-intensity conditioning (RIC) generally destroys more hematopoietic cells and is more toxic than nonmyeloablative conditioning, but less so than myeloablative conditioning. Both nonmyeloablative and RIC regimens are categorized as non-fully ablative regimens and are used interchangeably in this policy, unless otherwise noted. RIC/nonmyeloablative approaches can circumvent the need for high-dose conditioning regimens that are associated with organ toxicity and mortality depending on graft vs. tumor and immunosuppressive mechanisms.<sup>1</sup>

*Note: Please refer to PA.CP.MP.108 for requests for Allogeneic Hematopoietic Cell Transplants for Sickle Cell Anemia and  $\beta$ -Thalassemia.*

*Please refer to PA.CP.MP.162 Tandem Transplant if request is for an allogeneic reduced conditioning transplant for multiple myeloma in a tandem transplant.*

### Policy/Criteria

- I. It is the policy of Pennsylvania Health and Wellness<sup>®</sup> (PHW) that nonmyeloablative/reduced-intensity conditioning (RIC) allogeneic transplants are **medically necessary** for members/enrollees who meet all of the following criteria:
  - A. Candidate for allogeneic stem cell transplantation for any of the following diagnoses:
    1. Acute lymphoblastic leukemia;
    2. Acute myelogenous leukemia;
    3. Acquired bone marrow failure such as severe aplastic anemia;
    4. Familial bone marrow failure syndromes such as, but not limited to, one of the following:
      - a. Dyskeratosis congenita;
      - b. Shwachman-Diamond syndrome;
      - c. Diamond-Blackfan anemia;
      - d. Kostmann syndrome;
      - e. Fanconi anemia;
    5. Paroxysmal nocturnal hemoglobinuria;
    6. Chronic lymphocytic leukemias;
    7. Chronic myelogenous leukemia;
    8. Congenital immunodeficiency syndromes;
    9. Non-Hodgkin's lymphoma, any of the following:
      - a. Primary refractory or relapsed, including those who have relapsed after having an autologous bone marrow transplant (excluding diffuse large B-cell lymphoma);
      - b. Follicular lymphomas;
      - c. Mantle cell lymphoma;

## CLINICAL POLICY

### Nonmyeloablative Allogeneic Stem Cell Transplants

- d. Diffuse large B-cell lymphoma that is in remission following second-line therapy for relapsed or refractory disease;
  - 10. Myelodysplastic syndromes;
  - 11. Lysosomal storage disorders types IH/IS (Hurler/Hurler-Scheie), VI (maroteaux), VII (Sly);
  - 12. Macrophage discords such as hemophagocytic lymphohistiocytosis (HLH);
  - 13. Myeloproliferative neoplasms such as, but not limited to:
    - a. Chronic myeloid leukemia;
    - b. Juvenile myelomonocytic leukemia;
    - c. Primary myelofibrosis;
    - d. Essential thrombocytosis;
    - e. Polycythemia vera;
- B.** Unsuitable for conventional high-dose myeloablative allografting because of untreatable significant dysfunction of another major organ system and/or severe comorbidities, including, but not limited to, any of the following:
- 1. Bilirubin  $> 2$  mg/dL;
  - 2. Hemostasis: international normalized ratio (INR)  $> 1.6$  (unless on oral anticoagulants);
  - 3. Cardiac function: multigated acquisition (MUGA) scan or echocardiogram with ejection fraction (EF)  $< 45\%$ ;
  - 4. Pulmonary function, one of the following:
    - a. Forced expiratory volume in 1 second (FEV1)  $\leq 50\%$  of predicted value;
    - b. Diffusing capacity of the lung for carbon monoxide (DLCO)  $\leq 60\%$  of predicted value;
  - 5. Performance scale index, one of the following:
    - a. Karnofsky or Lansky score  $< 70\%$ ;
    - b. Eastern Cooperative Oncology Group (ECOG) performance score  $> 2$ .
- II.** It is the policy of PHW that current evidence does not support the use of nonmyeloablative/RIC allogeneic transplants for any of the following indications:
- A.** Solid tumors including, but not limited to:
- 1. Brain tumors;
  - 2. Ovarian epithelia and mixed epithelial/germ cell cancers;
  - 3. Primitive neuroectodermal tumors (PNET), including medulloblastoma and ependymoma;
  - 4. Renal cell carcinoma;
  - 5. Testicular cancer;
  - 6. Wilms tumor;
  - 7. Ewing sarcoma;
  - 8. Melanoma;
  - 9. Osteosarcoma;
  - 10. Rhabdomyosarcoma;
  - 11. Retinoblastoma;
  - 12. Germ cell tumors;
  - 13. Neuroblastoma;
  - 14. Multiple myeloma (except in tandem transplant- refer to CP.MP.162);

## CLINICAL POLICY

### Nonmyeloablative Allogeneic Stem Cell Transplants

- B. Autoimmune disorders including, but not limited to:
  - 1. Multiple sclerosis;
  - 2. Rheumatoid arthritis;
  - 3. Juvenile idiopathic arthritis;
  - 4. Systemic lupus erythematosus;
  - 5. Systemic sclerosis;
  - 6. Dermatomyositis;
  - 7. Polymyositis;
  - 8. Scleroderma;
- C. Hemoglobinopathies including, but not limited to:
  - 1. Thalassemias;
  - 2. Sickle cell anemia.

#### Background

Allogeneic hematopoietic cell transplantation (HCT) has been used as a treatment for cancer and diseases of the blood system for decades. For this treatment, stem cells are collected from either related or unrelated donors.<sup>1</sup> During the conditioning phase, high doses of chemotherapy (HDC), with or without radiation therapy, are used to eradicate the disease, and this is followed by infusion of stem cells to rescue bone marrow and restore normal immune function. Major limitations of this technique include the increased risk of high morbidity and mortality related to increased age, relapsed or refractory disease or disease with an elevated risk of relapse following HCT, a history of aggressive chemotherapy, and comorbidities.<sup>2</sup> All stem cell transplants (SCTs) preparative regimens have the potential for extensive toxicity. Loss of appetite and energy, alopecia, and nausea/vomiting occur frequently and contribute to poor physical and emotional tolerance of the transplant procedure. In addition, mucositis, diarrhea, and transient pancytopenia are inevitable side effects of most preparative regimens, and these complications are synergistic in dramatically increasing the risk of infections during and post-transplant.<sup>3</sup> Any decrease in toxicity, without concomitant loss of efficacy, would be desirable.

Myeloablative means that the treatment kills (ablates) the stem cells in the bone marrow; the cells that produce new blood cells. Several less intense conditioning regimens have been developed and rely more on immuno-suppression than cytotoxic effects to permit engraftment of donor cells. These regimens are collectively termed nonmyeloablative. Studies have shown that donor allogeneic stem cells can engraft in recipients using less-intensive conditioning regimens that are sufficiently immunosuppressive to permit graft-host tolerance. This manifests as a stable mixed donor-host hematopoietic chimerism, a term which means coexistence of donor and recipient cells. Nonmyeloablative allogeneic transplants, also referred to as “mini-transplant” or “transplant lite”, are thought to be potentially as effective as conventional HDC followed by an allogeneic stem cell transplantation, but with decreased morbidity and mortality related to the less intense, nonmyeloablative chemotherapy conditioning regimen.<sup>1,4</sup>

#### Coding Implications

This clinical policy references Current Procedural Terminology (CPT®). CPT® is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2023, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are

## CLINICAL POLICY

### Nonmyeloablative Allogeneic Stem Cell Transplants

included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

CPT® Codes	Description
38204	Management of recipient hematopoietic progenitor cell donor search and cell acquisition
38205	Blood-derived hematopoietic progenitor cell harvesting for transplantation, per collection; allogeneic
38207	Transplant preparation of hematopoietic progenitor cells; cryopreservation and storage
38208	Transplant preparation of hematopoietic progenitor cells; thawing of previously frozen harvest, without washing, per donor
38209	Transplant preparation of hematopoietic progenitor cells; thawing of previously frozen harvest, with washing, per donor
38210	Transplant preparation of hematopoietic progenitor cells; specific cell deletion within harvest. T-cell depletion
38211	Transplant preparation of hematopoietic progenitor cells; tumor depletion
38212	Transplant preparation of hematopoietic progenitor cells; red blood cell removal
38213	Transplant preparation of hematopoietic progenitor cells; platelet depletion
38214	Transplant preparation of hematopoietic progenitor cells; plasma (volume) depletion
38215	Transplant preparation of hematopoietic progenitor cells; cell concentration in plasma, mononuclear, or buffy coat layer
38230	Bone marrow harvesting for transplantation; allogeneic
38240	Hematopoietic progenitor cell (HPC), allogeneic transplantation per donor

HCPCS Codes	Description
S2142	Cord blood-derived stem cell transplantation, allogeneic
S2150	Bone marrow or blood-derived peripheral stem cells (peripheral or umbilical), allogeneic or autologous, harvesting, transplantation, and related complications; including pheresis and cell preparation/storage, marrow ablative therapy, drugs, supplies, hospitalization with outpatient follow-up, medical/surgical, diagnostic, emergency, and rehabilitative services, and the number of days of pre- and post-transplant care in the global definition

Reviews, Revisions, and Approvals	Revision Date	Approval Date
Updated description. Moved beta thalassemia and sickle cell anemia from the list of approved indications to the list of E/I indications. Removed age restriction from myelodysplastic syndromes. Added to the multiple myeloma indication that an RIC/NMA approach is appropriate post –autologous or fully	03/19	

## CLINICAL POLICY

### Nonmyeloablative Allogeneic Stem Cell Transplants

Reviews, Revisions, and Approvals	Revision Date	Approval Date
<p>myeloablative stem cell transplant. Removed diffuse large b-cell lymphoma from E/I list. Clarified that diffuse large cell lymphoma is diffuse large b-cell lymphoma, and added requirement that the patient is in remission following second-line therapy for relapsed or refractory disease. Specialist reviewed. References reviewed and updated.</p>		
<p>Added note to refer to CP.MP.108 for requests for Allogeneic HCT's for Sickle Cell Anemia and <math>\beta</math>-Thalassemia and CP.MP.162 if request is for a tandem transplant for multiple myeloma. Clarified in I.A.8. that Hodgkin's disease is now referred to as Hodgkin's lymphoma. Moved multiple myeloma and neuroblastoma from list of approved indications to the list of E/I indications. Removed sickle cell anemia from list of E/I indications. Removed CPT 38206 as code is for autologous transplant. Added ICD-10 Codes: D59.5, D75.81 Specialist reviewed. References reviewed and updated.</p>	10/2020	12/2020
<p>Annual review completed. References reviewed. Codes checked. Changed "members/enrollee" to members/enrollee." Specialty review completed with no updates.</p>	7/2021	
<p>Annual review. Rephrased criteria I.A.3. from "aplastic anemia" to "acquired bone marrow failure such as severe aplastic anemia." Added new indication I.A.4., "Familial bone marrow syndromes such as..." Removed "molecular remissions induced by Gleevec" from I.A.8." Added criteria points 13. and 14. to criteria I.A. "Experimental/investigational" verbiage in criteria II. replaced with descriptive language. Sorted list of non-supported indications in criteria II. into 3 subcategories, solid tumors, autoimmune disorders and hemoglobinopathies. In criteria I.C., combined and rephrased contraindications 2. and 3. and updated verbiage regarding substance abuse and dependence in 4. Minor rewording in description and background with no impact on criteria. Removed ICD-10 codes D57.00-D57.819 for sickle-cell disorders from ICD-10 table of codes to support coverage. References reviewed and updated. Changed "review date" in the header to "date of last revision" and "date" in the revision log header to "revision date." Reviewed by specialist.</p>	2/21/2023	
<p>Annual review completed. Background updated; minor rewording with no clinical significance. ICD-10 diagnosis code table removed. Removed Hodgkin's lymphoma from Criteria I.A.9. per updated National Comprehensive Cancer Network (NCCN) recommendations. Added Criteria I.A.13.e. to include polycythemia vera. Updated Criteria I.B.4.b. from diffusing capacity of the lung for carbon monoxide (DLCO) <math>\leq</math> 50% of predicted value to DLCO <math>\leq</math> 60% of predicted value. Removed</p>	03/2024	

**CLINICAL POLICY**  
**Nonmyeloablative Allogeneic Stem Cell Transplants**



Reviews, Revisions, and Approvals	Revision Date	Approval Date
absolute contraindications in Criteria I.C. References reviewed and updated. Reviewed by internal specialist and reviewed by external specialist.		

**References**

- American Cancer Society. Types of Stem Cell Transplants for Cancer Treatment. <https://www.cancer.org/treatment/treatments-and-side-effects/treatment-types/stem-cell-transplant/types-of-transplants.html>. Published March 20, 2020. Accessed January 16, 2024.
- Deeg HJ, Sandmaier BM. Determining eligibility for allogeneic hematopoietic cell transplantation. UpToDate. [www.uptodate.com](http://www.uptodate.com). Updated February 21, 2022. Accessed January 16, 2024.
- Negrin RS. Early complications of hematopoietic cell transplantation. UpToDate. [www.uptodate.com](http://www.uptodate.com). Updated June 30, 2022. Accessed January 24, 2024.
- Negrin RS. Preparative regimens for hematopoietic cell transplantation. UpToDate. [www.uptodate.com](http://www.uptodate.com). Updated February 4, 2022. Accessed January 24, 2024.
- Brodsky RA. Treatment and prognosis of paroxysmal nocturnal hemoglobinuria. UpToDate. [www.uptodate.com](http://www.uptodate.com). Updated December 3, 2021. Accessed January 16, 2024.
- National coverage determination: Stem cell transplantation (110.23). Centers for Medicare and Medicaid Services Web site. <http://www.cms.hhs.gov/mcd/search.asp>. Published January 27, 2016. Accessed January 16, 2024.
- Angelucci E, Benz EJ. Hematopoietic cell transplantation for transfusion-dependent thalassemia. UpToDate. [www.uptodate.com](http://www.uptodate.com). Updated February 27, 2023. Accessed January 17, 2024.
- Djulgovic B, Seidenfeld J, Bonnell C, Kumar A. Nonmyeloablative allogeneic stem-cell transplantation for hematologic malignancies: a systematic review. *Cancer Control*. 2003;10(1):17 to 41. doi:10.1177/107327480301000104
- Negrin RS. Treatment of high or very high risk myelodysplastic syndromes. UpToDate. [www.uptodate.com](http://www.uptodate.com). Updated July 28, 2020. Accessed January 22, 2024.
- Kröger N, Holler E, Kobbe G, et al. Allogeneic stem cell transplantation after reduced-intensity conditioning in patients with myelofibrosis: a prospective, multicenter study of the Chronic Leukemia Working Party of the European Group for Blood and Marrow Transplantation. *Blood*. 2009;114(26):5264 to 5270. doi:10.1182/blood-2009-07-234880
- Lee SE, Park SS, Jeon YW, et al. Outcomes of allogeneic stem cell transplantation in patients with paroxysmal nocturnal hemoglobinuria with or without aplastic anemia. *Eur J Haematol*. 2017;99(4):336 to 343. doi:10.1111/ejh.12922
- Moskowitz C, Alencar AJ. Hematopoietic cell transplantation in classic Hodgkin lymphoma. UpToDate. [www.uptodate.com](http://www.uptodate.com). Updated March 17, 2022. Accessed January 17, 2024.
- Castagna L, Crocchiolo R, Furst S, et al. Bone marrow compared with peripheral blood stem cells for haploidentical transplantation with a nonmyeloablative conditioning regimen and post-transplantation cyclophosphamide. *Biol Blood Marrow Transplant*. 2014;20(5):724 to 729. doi:10.1016/j.bbmt.2014.02.001
- National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology: B-cell lymphomas. Version 1.2024. [https://www.nccn.org/professionals/physician\\_gls/pdf/b-cell.pdf](https://www.nccn.org/professionals/physician_gls/pdf/b-cell.pdf). Published January 18, 2024. Accessed January 22, 2024.

## CLINICAL POLICY

### Nonmyeloablative Allogeneic Stem Cell Transplants

15. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology: Chronic lymphocytic leukemia/small lymphocytic lymphoma. Version 1.2024. [https://www.nccn.org/professionals/physician\\_gls/pdf/cll.pdf](https://www.nccn.org/professionals/physician_gls/pdf/cll.pdf). Published November 03, 2023. Accessed January 23, 2024.
16. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology: Chronic myeloid leukemia. Version 2.2024. [https://www.nccn.org/professionals/physician\\_gls/pdf/cml.pdf](https://www.nccn.org/professionals/physician_gls/pdf/cml.pdf). Published December 05, 2023. Accessed January 22, 2024.
17. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology: Hodgkin Lymphoma. Version 1.2024. [https://www.nccn.org/professionals/physician\\_gls/pdf/hodgkins.pdf](https://www.nccn.org/professionals/physician_gls/pdf/hodgkins.pdf). Published October 12, 2023. Accessed January 17, 2024.
18. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology: Multiple Myeloma. Version 2.2024. [https://www.nccn.org/professionals/physician\\_gls/pdf/myeloma.pdf](https://www.nccn.org/professionals/physician_gls/pdf/myeloma.pdf). Published November 01, 2023. Accessed January 24, 2024.
19. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology: Myelodysplastic syndromes. Version 1.2023. [https://www.nccn.org/professionals/physician\\_gls/pdf/mds.pdf](https://www.nccn.org/professionals/physician_gls/pdf/mds.pdf). Published November 10, 2023. Accessed January 17, 2024.
20. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology: Myeloproliferative Neoplasms. Version 1.2024. [https://www.nccn.org/professionals/physician\\_gls/pdf/mpn.pdf](https://www.nccn.org/professionals/physician_gls/pdf/mpn.pdf). Published December 21, 2023. Accessed January 22, 2024.
21. Gagelmann N, Kröger N. Dose intensity for conditioning in allogeneic hematopoietic cell transplantation: can we recommend "when and for whom" in 2021?. *Haematologica*. 2021;106(7):1794 to 1804. Published 2021 Jul 1. doi:10.3324/haematol.2020.268839
22. Brodsky RA. Paroxysmal nocturnal hemoglobinuria. *Blood*. 2014;124(18):2804 to 2811. doi:10.1182/blood-2014-02-522128
23. Oliansky DM, Czuczman M, Fisher RI, et al. The role of cytotoxic therapy with hematopoietic stem cell transplantation in the treatment of diffuse large B cell lymphoma: update of the 2001 evidence-based review. *Biol Blood Marrow Transplant*. 2011;17(1):20 to 47.e30. doi:10.1016/j.bbmt.2010.07.008
24. Pantin J, Tian X, Geller N, et al. Long-term outcome of fludarabine-based reduced-intensity allogeneic hematopoietic cell transplantation for debilitating paroxysmal nocturnal hemoglobinuria. *Biol Blood Marrow Transplant*. 2014;20(9):1435 to 1439. doi:10.1016/j.bbmt.2014.05.012
25. Pophali PA, Klotz JK, Ito S, et al. Male survivors of allogeneic hematopoietic stem cell transplantation have a long term persisting risk of cardiovascular events. *Exp Hematol*. 2014;42(2):83 to 89. doi:10.1016/j.exphem.2013.07.003
26. Samuelson S, Sandmaier BM, Heslop HE, et al. Allogeneic haematopoietic cell transplantation for myelofibrosis in 30 patients 60 through 78 years of age. *Br J Haematol*. 2011;153(1):76 to 82. doi:10.1111/j.1365-2141.2011.08582.x
27. Sieff CA. Overview of hematopoietic stem cells. UpToDate. [www.uptodate.com](http://www.uptodate.com). Updated March 9, 2022. Accessed January 25, 2024.
28. Storb R, Sandmeier BM. Nonmyeloablative allogeneic hematopoietic cell transplantation.

## CLINICAL POLICY

### Nonmyeloablative Allogeneic Stem Cell Transplants

- Haematologica*. 2016;101(5): 521 to 530. doi:10.3324/haematol.2015.132860
29. Tefferi A. Management of primary myelofibrosis. UpToDate. [www.uptodate.com](http://www.uptodate.com). Updated October 7, 2022. Accessed January 25, 2024.
  30. Velázquez-Sánchez-de-Cima S, Zamora-Ortiz G, Hernández-Reyes J, et al. Oral versus intravenous fludarabine as part of a reduced-intensity conditioning for allogeneic stem cell transplantation. *Acta Haematol*. 2014;132(1):125 to 128. doi:10.1159/000357108
  31. Rajkumar SV. Multiple myeloma: Use of allogeneic hematopoietic cell transplantation. UpToDate. [www.uptodate.com](http://www.uptodate.com). Updated March 28, 2022. Accessed January 25, 2024.
  32. Mikhael J, Ismaila N, Cheung MC, et al. Treatment of Multiple Myeloma: ASCO and CCO Joint Clinical Practice Guideline [published correction appears in *J Clin Oncol*. 2020 Jul 20;38(21):2469]. *J Clin Oncol*. 2019;37(14):1228 to 1263. doi:10.1200/JCO.18.02096
  33. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology: Primary cutaneous lymphomas. Version 1.2024. [https://www.nccn.org/professionals/physician\\_gls/pdf/primary\\_cutaneous.pdf](https://www.nccn.org/professionals/physician_gls/pdf/primary_cutaneous.pdf). Published December 21, 2023. Accessed January 16, 2024.
  34. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology: T-cell lymphomas. Version 1.2024. [https://www.nccn.org/professionals/physician\\_gls/pdf/t-cell.pdf](https://www.nccn.org/professionals/physician_gls/pdf/t-cell.pdf). Published December 21, 2023. Accessed January 16, 2024.
  35. Rogers ZR, Myers KC. Shwachman-Diamond syndrome. UpToDate. [www.uptodate.com](http://www.uptodate.com). Updated August 12, 2022. Accessed January 24, 2024.
  36. Khan S, Myers KC. Hematopoietic cell transplantation (HCT) for inherited bone marrow failure syndromes (IBMFS). UpToDate. [www.uptodate.com](http://www.uptodate.com). Updated November 23, 2021. Accessed January 24, 2024.
  37. Chien JW, Sullivan KM. Carbon monoxide diffusion capacity: how low can you go for hematopoietic cell transplantation eligibility?. *Biol Blood Marrow Transplant*. 2009;15(4):447 to 453. doi:10.1016/j.bbmt.2008.12.509