

Clinical Policy: Trigger Point Injections for Pain Management

Reference Number: PA.CP.MP.169

Effective Date: 09/18

Last Review Date: 09/18

[Coding Implications](#)

[Revision Log](#)

Description

Trigger points cause pain at their physical location and also referred pain to other areas in a specific pattern. Trigger point injections consist of an injection of a local anesthetic, with or without steroid medication, into a painful portion of the muscle containing the trigger point.

Policy/Criteria

It is the policy of Pennsylvania Health and Wellness® that invasive pain management procedures performed by a physician are **medically necessary** when *the relevant criteria are met and the patient receives only one procedure per visit, with or without radiographic guidance.*

- I. Trigger point injections are **medically necessary** for the following indications:
 - A. *Diagnosis/stabilization* of trigger points with injections of corticosteroids and/or local anesthetics at the trigger point, all of the following:
 1. The member has local pain symptoms in the neck, shoulder and/or back that have persisted for more than 3 months causing tenderness and/or weakness, restricting motion and/or causing referred pain when compressed;
 2. The member has failed ≥ 3 weeks of conventional multidisciplinary medical therapy including all of the following:
 - a. Chiropractic, physical therapy, or prescribed home exercise program or the member is unable to tolerate such therapy and the injection is intended as a bridge to therapy;
 - b. NSAID unless contraindicated or not tolerated;
 - c. Activity modification;
 3. Trigger points have been identified by palpation;
 4. Trigger points are located in a *few* discrete areas and are not associated with widespread areas of muscle tenderness (as with fibromyalgia);
 5. Injections are not used as sole method of treatment, rather are intended for pain relief to facilitate mobilization to allow non-invasive modalities, e.g., physical therapy and other alternate therapies that address muscle strengthening, flexibility, and functional restoration.
 - B. *Additional trigger point injections*, all of the following:
 1. Prior injections resulted in $\geq 50\%$ improvement for ≥ 6 weeks;
 2. There was a return of pain and/or deterioration following 6 weeks of improvement;

Up to 2 sets of injections at least 7 days apart may be given for diagnosis and stabilization for the same trigger point. When a given body region is injected, it will be considered as one injection service no matter how many injections are given.

CLINICAL POLICY

Trigger Point Injections

3. Injections are given in the neck, shoulder, and/or back;
4. Injections are given at least 2 months apart for up to 12 months (maximum of 6 sessions);
5. Injections are not used as sole method of treatment, but rather are intended for pain relief to facilitate mobilization to allow for non-invasive modalities, e.g., physical therapy and other alternate therapies that address muscle strengthening, flexibility, and functional restoration.

When a given body region is injected, it will be considered as one injection service no matter how many injections are given.

II. *Trigger point therapies* for the following indications are considered **not medically necessary**, because although there are ongoing studies, there is little scientifically based data that their use results in improved patient outcomes in the medical literature:

- A. Dry needle stimulation of trigger points;
- B. Trigger point injection with saline or glucose;
- C. The use of Botox during trigger point injections.

Background

A trigger point is a discrete, hyperirritable focus found in a palpable taut band occurring in any skeletal muscle and/or muscle fascia on the body that is particularly sensitive to touch and, when compressed, gives rise to characteristic referral pain patterns, tenderness and autonomic phenomena. Trigger points are thought to result from repetitive strain produced by acute or chronic overload or a degenerative and/or inflammatory problem, such as arthritis.

Trigger point injections of local anesthetic and/or steroids are a common intervention for back and neck pain, although evidence is mixed. A Cochrane review of injections for subacute and chronic back pain found no clear advantage of local or trigger point injections with a local anesthetic, with or without a corticosteroid, and control interventions for short-term pain relief across 3 trials.¹⁻³ Another systematic review found that intramuscular injection of lidocaine more effectively relieved neck pain in the short term than placebo.⁴

A systematic review of trigger point injections with botulinum toxin concluded that a statistically or clinically significant benefit could not be confirmed from the use of botulinum toxin-A used alone for chronic neck pain in the short term.⁶ Secondary outcomes such as pain, disability, and quality of life were also investigated without confirmed benefit of botulinum injections.⁶

There is preliminary evidence that dry needling of trigger points is effective for short-term pain relief, and to improve quality of life and range of motion when compared to a placebo, but further studies of high quality and with a standardized needling procedure are needed.⁷

Coding Implications

This clinical policy references Current Procedural Terminology (CPT®). CPT® is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2018, American Medical Association. All rights reserved. CPT codes and CPT descriptions are

CLINICAL POLICY

Trigger Point Injections

from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

CPT® Codes	Description
20552	Injection(s); single or multiple trigger point(s), 1 or 2 muscle(s)
20553	Injection(s); single or multiple trigger point(s), 3 or more muscles

HCPCS Codes	Description
N/A	

ICD-10-CM Diagnosis Codes that Support Coverage Criteria

+ Indicates a code requiring an additional character

ICD-10-CM Code	Description
M25.511- M25.519	Pain in shoulder
M54.5	Low back pain
M54.9	Dorsalgia, unspecified
M79.1	Myalgia

Reviews, Revisions, and Approvals	Date	Approval Date
New policy split from retired CP.MP.118 Injections for Pain Management. No criteria changes.	09/18	

References

1. Staal JB et al. Injection therapy for subacute and chronic low-back pain. *Cochrane Database of Systematic Reviews* 2008, Issue 3. Art. No. CD001824. DOI: 10.1002/14651858.CD001824.pub3.
2. Chou R. Subacute and chronic low back pain: Nonsurgical interventional treatment. In: UpToDate, Atlas SJ (Ed), UpToDate, Waltham, MA. Accessed 8/15/18.
3. Scott NA, Guo B, Barton PM, Gerwin RD. Trigger point injections for chronic non-malignant musculoskeletal pain: A systematic review. *Pain Med.* 2009;10(1):54-69.
4. Peloso P, Gross A, Haines T, et al. Medicinal and injection therapies for mechanical neck disorders. *Cochrane Database Syst Rev.* 2007;(3):CD000319.
5. Cummings TM, White AR. Needling therapies in the management of myofascial trigger point pain: A systematic review. *Archiv Phys Med Rehab*, 2001;82(7):986-992.
6. Langevin P, Lowcock J, Weber J. Botulinum toxin intramuscular injections for neck pain: a systematic review and metaanalysis. *J Rheumatol.* 2011;38(2):203.

CLINICAL POLICY

Trigger Point Injections

7. Espejo-Antúnez L, Tejeda JF, Albornoz-Cabello M, et al. Dry needling in the management of myofascial trigger points: A systematic review of randomized controlled trials. *Complement Ther Med*. 2017 Aug;33:46-57. doi: 10.1016/j.ctim.2017.06.003. Epub 2017 Jun 15.
8. Chou R, et al. Diagnosis and treatment of low back pain: a joint clinical practice guideline from the American College of Physicians and the American Pain Society. *Ann Intern Med*. 2007;147:478-491.
9. Gerwin R. Botulinum toxin treatment of myofascial pain: a critical review of the literature. *Curr Pain Headache Rep*. 2012 Oct;16(5):413-22. doi: 10.1007/s11916-012-0287-6.
10. Han SC, Harrison P. Myofascial pain syndrome and trigger-point management. *Reg Anesth*. 1997 Jan-Feb;22(1):89-101.
11. Isaac Z. Treatment of neck pain. Atlas SJ (Ed). Updated Sep 21, 2017. In: UpToDate Waltham, MA: UpToDate. Accessed 8/16/18.