

Clinical Policy: Bariatric Surgery

Reference Number: PA.CP.MP.37

Effective Date: 01/2018

Date of Last Revision: 04/2024

[Coding Implications](#)

[Revision Log](#)

Description

There are two categories of bariatric surgery: restrictive procedures and malabsorptive procedures. Gastric restrictive procedures include procedures where a small pouch is created in the stomach to restrict the amount of food that can be eaten, resulting in weight loss. The laparoscopic adjustable gastric banding (LAGB) and laparoscopic sleeve gastrectomy (LSG) are examples of restrictive procedures. Malabsorptive procedures bypass portions of the stomach and intestines causing incomplete digestion and absorption of food. Duodenal switch is an example of a malabsorptive procedure. Roux-en-y gastric bypass (RYGB), biliopancreatic diversion with duodenal switch (BPD-DS), and biliopancreatic diversion with gastric reduction duodenal switch (BPD-GRDS) are examples of restrictive and malabsorptive procedures.¹

LAGB devices are currently not FDA approved for adolescents less than 18 years and are being used less for adolescents in favor of sleeve gastrectomy (SG).²

Policy/Criteria

- I. It is the policy of Pennsylvania Health and Wellness[®] (PHW) that bariatric surgery is **medically necessary** when the following criteria under **section A, B, and C** are met:
 - A. Medical history, meets all of the following:
 1. Age and body mass index (BMI) (meet criteria in a *or* b):
 - a. Age ≥ 18 and one of the following (i or ii):
 - i. BMI ≥ 35 kg/m² or ≥ 32.5 kg/m² for South Asian, Southeast Asian, and East Asian adults when laparoscopic adjustable gastric banding (LAGB), laparoscopic sleeve gastrectomy (LSG), laparoscopic Roux-en-y gastric bypass (RYGB) or laparoscopic biliopancreatic diversion with duodenal switch (BPD-DS)/biliopancreatic diversion with gastric reduction duodenal switch (BPD-GRDS) is requested;
 - ii. BMI ≥ 30 and < 35 kg/m², or ≥ 27.5 and < 32.5 kg/m² for South Asian, Southeast Asian, and East Asian adults, when LAGB, LSG, laparoscopic RYGB or BPD-DS/BPD-GRDS is requested and at least one of the following:
 - a) Obesity has continued despite previous weight loss attempts using nonsurgical methods;
 - b) Type 2 diabetes mellitus (DM);
 - c) One of the following obesity related co-morbidities has not improved despite using nonsurgical weight loss methods:
 - i. Hypertension;
 - ii. Dyslipidemia;
 - iii. Obstructive sleep apnea;
 - iv. Obesity-hypoventilation syndrome/Pickwickian syndrome;
 - v. Nonalcoholic fatty liver disease or nonalcoholic steatohepatitis;
 - vi. Pseudotumor cerebri;
 - vii. Coronary artery disease;
 - viii. Gastroesophageal reflux disease;

- ix. Asthma;
 - x. Venous stasis disease;
 - xi. Bone and joint diseases;
 - xii. Disqualification from other specialty surgeries due to obesity (i.e., joint arthroplasty, abdominal wall hernia repair, or organ transplantation);
 - xiii. Chronic kidney disease;
 - xiv. Infertility;
 - xv. Polycystic ovarian syndrome;
- b. Age < 18 years, LSG or laparoscopic RYGB is requested, and one of the following (i. or ii.):
- i. BMI \geq 40 kg/m² or 140% of the 95th percentile (whichever is lower);
 - ii. BMI \geq 35 kg/m² or 120% of the 95th percentile (whichever is lower) with \geq 1 severe comorbidity listed below that has significant short-term effects on health, and that is uncontrolled with lifestyle or pharmacotherapy management:
 - a) Type 2 DM;
 - b) Obstructive sleep apnea;
 - c) Idiopathic intracranial hypertension;
 - d) Nonalcoholic fatty liver disease or nonalcoholic steatohepatitis;
 - e) Blount's disease;
 - f) Slipped capital femoral epiphysis (SCFE);
 - g) Gastroesophageal reflux disease;
 - h) Hypertension;
 - i) Hyperlipidemia;
 - j) Insulin resistance;
- B. Preoperative evaluation and medical clearance requirements within six months of the scheduled surgery include *all* of the following:
1. Cardiac evaluation and *one* of the following categories (a. or b.):
 - a. LOW CARDIAC RISK candidates, with *none* of the risk factors listed in section I.B.1.b. below, need cardiac clearance by a PCP or cardiologist. If additional testing is needed, it should be conducted by a cardiologist;
 - b. HIGH CARDIAC RISK candidates need consultation/evaluation, including electrocardiogram, and cardiac clearance from a cardiologist. High risk candidates include those with *any* of the following:
 - i. History of ischemic heart disease;
 - ii. History of heart failure;
 - iii. History of cerebrovascular disease;
 - iv. Glomerular filtration rate < 60 mL/min⁻¹;
 - v. High-grade arrhythmia;
 - vi. Hemodynamically significant valvular heart disease;
 - vii. Unstable angina;
 - viii. Recent myocardial infarction (within the past 60 days);
 2. Glycemic control evaluation to include A1c and fasting blood glucose;
 3. Pulmonary Evaluation:
 - a. Screening for obstructive sleep apnea with confirmatory polysomnography if screening tests are positive;

- b. Pulmonary function testing and arterial blood gas analysis for candidates with intrinsic lung disease or disordered sleep patterns;
 - c. Specialist consultation for interpretation of any abnormal findings;
 4. Nutritional evaluation, including micronutrient measurements such as, but not limited to, thiamine, calcium, iron studies, vitamin B12, folic acid, and fat soluble vitamins, and treatment of insufficiencies/deficiencies prior to surgery. More extensive testing may be necessary in patients undergoing malabsorptive procedures, based on symptoms and risks;
 5. Nutritional therapy/counseling including all of the following:
 - a. Initial comprehensive diet history to include assessment of current pattern of nutrition and exercise and steps to modify problem eating behaviors;
 - b. Documentation that counseling has been conducted regarding the potential for success of weight loss surgery dependent on post-op diet modification (if patient < 18 years of age, consultation must be with adolescent AND parent/guardian);
 6. Age-appropriate psychiatry/psychology consultation including all of the following:
 - a. An in-person psychological evaluation to assess for major mental health disorders which would contradict surgery and determine ability to comply with post-operative care and guidelines;
 - b. If age < 18 years: evaluation must also include assessment of emotional maturity, decisional capacity, family support and family willingness to participate in lifestyle changes;
 7. A serum TSH level if clinical evidence of hypothyroid is present and treatment if found to be hypothyroid;
 8. A fasting lipid panel obtained and treatment initiated for dyslipidemia;
 9. Gastrointestinal (GI) screening and evaluation for clinically significant GI symptoms with documentation of needed treatment prior to bariatric surgery;
 10. Prophylactic treatment for gouty attacks in patients with a history of gout;
 11. If tobacco user, must stop use > six weeks prior to surgery;
 - C. None of the following contraindications for surgical weight loss procedures:
 1. Medically correctable causes of obesity;
 2. Current or planned pregnancy within 12 to 18 months of the procedure;
 3. Severe cardiac disease with prohibitive anesthetic risks;
 4. Severe coagulopathy;
 5. Current drug or alcohol abuse;
 6. Uncontrolled and untreated eating disorders (e.g., bulimia);
 7. Inability on the part of the patient or parent/guardian to comprehend the risks and benefits of the surgical procedure;
 8. A medical, psychiatric, psychosocial, or cognitive condition that prevents adherence to postoperative dietary and medication regimens or impairs decisional capacity.
- II.** It is the policy of PHW that *repeat bariatric surgery* is considered medically necessary for one of the following:
 - A. To correct complications from a previous bariatric surgery, such as obstruction or strictures (could include conversion surgeries to LSG or RYGB for adults or adolescents; or BPD-DS for adults);

- B. Conversion from LAGB to an LSG, RYGB or BPD-DS; or revision of a primary procedure that has failed due to dilation of the gastric pouch when all of the following criteria are met:
 - 1. All criteria listed above for the initial bariatric procedure are met again;
 - 2. Previous surgery for morbid obesity was at least two years prior to repeat procedure;
 - 3. Weight loss from the initial procedure was less than 50% of the member/enrollee's excess body weight at the time of the initial procedure;
 - 4. If the conversion is requested due to removal of an eroded laparoscopic adjustable band, at least two months have passed between the band removal and the subsequent bariatric procedure;
 - 5. Documented compliance with previously prescribed postoperative nutrition and exercise program;
 - 6. Supporting documentation from the provider should also include a clinical explanation of the circumstances as to why the procedure failed;
- C. Conversion of SG to RYGB for the treatment of gastro-esophageal reflux disease (GERD) when anti-reflux medical therapy has been tried and failed;
- D. Conversion of SG to RYGB or BPD-DS as a bridging procedure for BMI \geq 50 kg/m².

III. It is the policy of PHW that the current medical literature is inadequate to determine the safety, efficacy, and long-term outcomes for the following bariatric surgery procedures:

- A. Distal gastric bypass (very long limb gastric bypass);
- B. Mini gastric bypass—one anastomosis gastric bypass (e.g., mini-gastric bypass, one-anastomosis gastric bypass, single anastomosis gastric bypass, omega loop gastric bypass);
- C. Laparoscopic re-sleeve gastrectomy (LRSG) performed after the resulting gastric pouch is primarily too large or dilates after the original LSG;
- D. Fobi pouch;
- E. Laparoscopic greater curvature plication (Gastric Imbrication);
- F. LAP-BAND when BMI is 30 to 35 with or without comorbid conditions;
- G. Stomach aspiration therapy (e.g., AspireAssist);
- H. Endoscopic Suture Revisions post bariatric surgery;
- I. Single anastomosis duodenoileal bypass (SADI);
- J. Gastric plication/ Endoluminal vertical gastroplasty;
- K. Endoscopic gastrointestinal bypass devices (EGIBD);
- L. Endoscopic sleeve gastroplasty;
- M. Transoral endoscopic surgery;
- N. Vagus Nerve Blocking (e.g., Maestro);
- O. Gastric balloon (e.g., ReShape Duo, Orbera intragastric balloon, Obalon Balloon).

IV. It is the policy of PHW that the following bariatric surgery procedures are considered **not medically necessary**, due to potential complications and a lack of positive outcomes:

- A. Biliopancreatic diversion (BPD) procedure (also known as the Scopinaro procedure);
- B. Jejunioleal bypass (jejuno-colic bypass);
- C. Vertical Banded Gastroplasty (VBG);
- D. Gastric pacing/gastric electrical stimulation;
- E. Gastric wrapping.

- F. For PA Medicaid products, PHW will consider on a case by case basis:
- a. Biliopancreatic diversion (BPD) procedure (also known as the Scopinaro procedure), although BPD with duodenal switch will be considered on a case by case basis; and
 - b. Vertical Banded Gastroplasty (VBG), (this will also be considered on a case by case basis).

Background

Persons with clinically severe obesity are at risk for increased mortality and multiple co-morbidities. These co-morbidities include hypertension, hypertrophic cardiomyopathy, hyperlipidemia, diabetes, cholelithiasis, obstructive sleep apnea, hypoventilation, degenerative arthritis and psychosocial impairments. The majority of severely obese patients losing weight through non-operative methods alone regain all the weight lost over the next five years. Surgical treatment is the only proven method of achieving long term weight control for the morbidly obese. Eating behaviors after surgery improve dramatically due to the restricted size of the stomach, allowing only small amounts of food to be taken in at a time.^{3,4}

The type of surgical procedure performed should be based on body mass index (BMI), comorbidity profile, treatment goals, surgeon's expertise, patient preference and risk stratification.⁵ The most commonly performed bariatric procedure in the United States is laparoscopic sleeve gastrectomy (LSG), followed by laparoscopic Roux-en-Y gastric bypass (RYGB), laparoscopic adjustable gastric banding (LAGB), and Biliopancreatic diversion with duodenal switch (BPD-DS).⁵ The sleeve gastrectomy (SG) continues to trend upwards due to lower rates of complications and nutritional deficiencies while maintaining comparable weight loss and metabolic disease outcomes.⁵ It was the most commonly performed bariatric procedure in the United States and in the world in 2016, and laparoscopic surgery is the preferred methodology.¹

The success of the bariatric surgery relies on the motivation and dedication to the program of the patient. The patient must be able to participate in the treatment and long-term follow up required after surgery. Studies have shown that about 10% of patients may have unsatisfactory weight loss or regain much of the weight they have lost. This may occur due to frequent snacking on high-calorie foods or lack of exercise. Technical problems that may occur include a stretched pouch due to overeating following surgery. Ensuring patients are motivated to lose weight can help prevent some of these issues.

Maximum weight loss usually occurs between 18 and 24 months postoperatively. The average weight loss at five years ranges from 48 to 74% after gastric bypass and 50 to 60% following gastric banding. Several studies have follow-up from five to 15 years with these patients maintaining weight loss of 50 to 60% of excess weight.

The Lap Band is a small bracelet-like band placed around the top of the stomach to produce a small pouch about the size of a thumb. The size of the outlet is controlled by a circular balloon inside the band that can be inflated and deflated with saline solution through an access port placed under the skin. The more inflated the balloon, the narrower the opening and slower passage of food to the rest of the stomach.¹

RYGB creates a small stomach pouch, bypassing most of the stomach, duodenum, and upper intestine.¹ Weight loss occurs through restriction of food intake and by decreasing the absorption of food by re-routing food directly from the pouch into the small intestine. With over 25 years of experience with RYGB in adults, the long-term results are well established for weight loss and improvement in comorbidities, and this surgery now accounts for approximately 20% of bariatric procedures in adolescents.⁶

BPD-DS is a complex operation that includes removing a large portion of the stomach to promote smaller meal sizes, re-routing of food away from much of the small intestine to prevent partial absorption of food, and re-routing of bile and other digestive juices that impair digestion. The operation bypasses most of the duodenum but leaves a small portion for food and the absorption of some vitamins and minerals. BPD-DS produces significant weight loss but has a greater risk of long-term complications due to decreased absorption of food, vitamins, and minerals.¹

American Society for Metabolic and Bariatric Surgery (ASMBS)

Updated guidelines from the ASMBS recommend metabolic and bariatric surgery for patients with BMI ≥ 35 kg/m², regardless of presence, absence, or severity of co-morbidities and for patients with BMI of 30 to 34.9 kg/m² who do not achieve substantial, durable weight loss or comorbidity improvement with reasonable nonsurgical methods, bariatric surgery should be considered. In this population, surgical intervention should be considered after failure of nonsurgical treatments. For patients with type two diabetes, bariatric and metabolic surgery is now recommended for those with BMI ≥ 30 kg/m². LAGB, LSG, and RYGB have been shown to be well-tolerated and effective treatments. Safety and efficacy of these procedures in low-BMI patients appear to be similar to results in patients with severe obesity. Currently, the best evidence for bariatric and metabolic surgery for patients with class I obesity and co-morbid conditions exists for patients in the 18 to 65 age group.^{4,7}

Bariatric Surgery in Adolescents

Weight loss surgery has been performed in small groups of adolescents since the 1970s. Recent data has shown a significant increase in the rate since 2000.² It is likely that we will continue to see a rise in the rate of adolescents undergoing weight loss surgery with the current pediatric obesity epidemic. Children and adolescents who are severely obese are at risk for the same mortality and co-morbidities as adults.^{8,9} These co-morbidities include hypertension, hypertrophic cardiomyopathy, hyperlipidemia, diabetes, cholelithiasis, obstructive sleep apnea, depression and impaired quality of life. In addition, children in the BMI category ≥ 35 kg/m² will almost always remain obese, and 65% will have a BMI ≥ 40 as an adult.⁶

Changes in diet and physical activity must be attempted prior to weight loss surgery in adolescents. A multi-disciplinary, family-based approach should be undertaken to support a staged weight loss plan.¹⁰ However, studies suggest that dietary and behavioral interventions rarely result in significant and sustained weight loss in adolescents. This same multi-disciplinary and family approach must be taken when evaluating and planning for bariatric surgery in an adolescent.^{8,9}

Recently updated guidelines from the ASMBS on pediatric metabolic and bariatric surgery conclude that metabolic and bariatric surgery (MBS) is a proven, effective treatment for severe obesity disease in adolescents and should be considered standard of care. Treatment of severe obesity in adolescents clearly requires a multidisciplinary approach where MBS should not be consigned to the treatment of last resort. Rather, when considered appropriate and within the clinical best practice guidelines, MBS should be readily offered to adolescents with obesity to effectively reverse co-morbidities and achieve overall wellness. Prior weight loss attempts, Tanner stage, and bone age should not be barriers to definitive treatment.^{2,4}

Investigational Procedures

Long-limb or Distal Gastric Bypass for Superobesity: A randomized controlled trial (RCT) was completed by Svanevik et al., but only perioperative outcomes have been reported thus far. Svanevik et al. found that in superobese patients with BMI between 50 and 60 kg/m², distal gastric bypass was associated with longer operating time and more severe complications resulting in reoperation than proximal gastric bypass.¹¹ There is increased risk of adverse nutritional outcomes with longer limb gastric bypass. At this time the long-limb or distal gastric bypass for superobesity is considered investigational, until more long-term studies can be done which reflect better outcomes than existing procedures.

Loop Gastric Bypass (Mini Gastric Bypass, one-anastomosis gastric bypass): The mini gastric bypass has not been universally accepted due to higher rates of alkaline bile reflux and limited long-term research. More long-term research is needed to solidify mini gastric bypass surgery's position as a viable bariatric surgery option.¹

Re-Sleeve Gastrectomy for Failed Laparoscopic Sleeve Gastrectomy: In 2012 Iannelli et al. noted that laparoscopic sleeve gastrectomy (LSG) was rapidly accepted as a valuable bariatric procedure before its effectiveness on weight loss in the long-term is clearly demonstrated.¹² The authors report a feasibility study including 13 patients undergoing a redo LSG for either progressive weight regain after initial weight loss or insufficient weight loss.¹³ AlSabah et al. describe 24 patients who underwent re-sleeve laparoscopic gastrectomy after an initial LSG. Compared to 12 patients that initially had LSG, which was converted to LRYGB, results were similar, with no significant differences in percent of excess weight loss at one year.¹⁴ They conclude that larger and longer follow-up studies are needed to verify results.¹⁴

Fobi Pouch or Silastic® Ring: The Fobi Pouch bariatric operation for obesity is a combination of stomach reduction and gastric bypass. The Silastic ring is placed around the vertically constructed gastric pouch above the anastomosis between the pouch and the intestinal Roux limb. Possible long term nutritional deficiencies involve fat soluble vitamin deficiencies of Calcium, Iron, B12, and Folic Acid. Patients are placed on nutritional supplements for the rest of their lives, and yearly monitoring is needed. The Fobi Pouch gastric bypass takes about double the time that a vertical banded gastroplasty operation takes. There is limited research on the outcomes of the Fobi pouch versus other bariatric surgery procedures.¹⁵

Gastric Imbrication: Fried et al. completed a three year RCT on the safety and efficacy of laparoscopic adjustable gastric banding with and without imbrication sutures.¹⁶ The results of the RCT have demonstrated that single anastomosis gastric bypass (SAGB) combined with a

conservative approach to band adjustments and limited retrogastric dissection is effective and safe with and without imbrication sutures. Not using imbrication sutures results in significant benefits in operative speed with comparable clinical weight loss and intermediate term safety.¹⁶ Sharma et al. conducted a randomized, double blinded trial comparing LSG and laparoscopic gastric imbrication (LGI). They found no differences in weight, age, or BMI preoperatively at six months or three years between the two groups.¹⁷

The AspireAssist System (AspireAssist) was FDA approved in 2016. It is a weight loss device comprised of an endoscopically placed percutaneous gastrostomy tube and an external device to facilitate drainage of about 30% of each meal consumed. It is meant to be used in conjunction with diet and exercise. In 2017 a one-year RCT was performed comparing results of 207 patients treated with AspireAssist.¹ The treatment group (n=137) received AspireAssist and lifestyle counseling, and the control group (n=70) received lifestyle counseling alone. Compared to the control group, those who received the AspireAssist and counseling lost more weight. 58.6% of participants in the AspireAssist group, and 15.3% of participants in the Lifestyle Counseling group lost at least 25% of their excess body weight (P<0.001).¹ Additionally, a prospective observational study was conducted on 25 patients, and by the end of the two-year observation period, only 15 patients were still in the study. They concluded that AspireAssist is an efficient and safe treatment for obesity. There is no research on AspireAssist versus other bariatric surgery procedures.¹⁸

To enhance weight loss, the following endoscopic procedures have been attempted to promote restriction of the pouch or stoma. These revisions have included: sclerotherapy of the site using 6 to 30 mL of sodium morrhuate injected circumferentially; tissue plication systems to reduce the size of the gastrojejunostomy and the gastric pouch; revisional surgery using a tissue plication device known as StomaPhyX to reduce the pouch size; and application of the endoclip to reduce the size of the gastrojejunal anastomosis. There is a lack of long-term outcomes for endoscopic revisions post RYGB.¹

The single anastomosis duodenoileal bypass (SADI), also known as single-anastomosis duodenal switch (SADS) and most descriptively, single-anastomosis duodenoileal bypass with sleeve gastrectomy (SADI-S), combines restrictive, malabsorptive, and probably hormonal mechanisms for weight loss. The sleeve is created first, and the duodenum is divided after the pylorus. SADI creates an anastomosis between the side of the distal ileum and the end of the sleeve-like gastric pouch/duodenum.¹

The ASMBS endorses SADI-S as an appropriate primary metabolic bariatric procedure.¹ Per the ASMBS, the SADI-S procedure is fundamentally a variant of the duodenal switch (DS) operation, in which the transected duodenum is anastomosed to a loop of ileum, as opposed to the classic DS in which a Roux-en-Y configuration is used. However, the ASMBS- notes the publication of long-term safety and efficacy outcomes is still needed and is strongly encouraged, particularly with published details on SG size and common channel length. There remain concerns about intestinal adaptation, nutritional issues, optimal limb lengths, and long-term weight loss/regain after this procedure. As such, ASMBS recommends a cautious approach to the adoption of this procedure, with attention to ASMBS-published guidelines on nutritional and metabolic support of bariatric patients, in particular for DS patients.^{1,19}

The International Federation for the Surgery of Obesity and Metabolic Disorders (IFSO) considers SADI-S safe and effective based on short-term data from studies but recommends that long-term follow up be continued and that randomized controlled trials be performed in the near future.¹⁹ In a 2021 updated statement, IFSO emphasized that SADI-S can result in maintaining significant weight loss for the obese individual, but nutritional deficiencies are a long-term safety concern, and patients need to be aware of this and encouraged to remain in long-term multidisciplinary care.¹⁹

Additionally, the National Institute for Health and Care and Excellence (NICE) encourages further research into SADI-S with a focus on long-term outcomes.⁵ NICE recommendations also state that there are well-recognized complications when treating morbid obesity with SADI-S, including the possibility of serious metabolic complications.⁵ NICE states, “this procedure should only be used with special arrangements for clinical governance, consent and audit or research.”⁵

Endoluminal vertical gastroplasty/gastric plication is an endoscopic approach for suturing the stomach that offers the potential to perform gastric-restrictive procedures endoluminally. The anterior and posterior walls of the stomach are suctioned together, then held in place by either a stapler or T-fastener device to create a tube of stomach similar to the sleeve gastrectomy.¹

Endoscopic gastrointestinal bypass devices (EGIBD) are barrier devices deployed to prevent luminal contents from being absorbed in the proximal small intestine (e.g., ValenTX, EndoBarrier). Data are still lacking about the longevity of these endobarriers and their outcomes once the barrier is removed.¹

Not Medically Necessary Procedures

Biliopancreatic Diversion (BPD) Procedure (Scopinaro procedure): The biliopancreatic diversion (BPD) is a malabsorptive procedure that was introduced as a solution to the high rates of liver failure resulting from bowel exclusion in the jejunioleal bypass. The procedure consists of a partial gastrectomy and gastroileostomy with a long segment of Roux limb and a short common channel, resulting in fat and starch malabsorption. BPD also has a restrictive component. The BPD/DS procedure differs from the BPD in the portion of the stomach that is removed, as well as preservation of the pylorus. This allows more forward flow of the contents of the biliopancreatic limb and avoids the complications of stasis that plagued the jejunioleal bypass (JIB). It is associated with fewer complications than BPD alone. BPD/DS is a complex procedure that is only performed at a few centers in the U.S.¹

Jejunioleal Bypass or Jejunioleal Intestinal Bypass (JIB): The jejunioleal bypass (also called the intestinal bypass) is performed by dividing the jejunum close to the ligament of Treitz and connecting it a short distance proximal to the ileocecal valve, thereby diverting a long segment of small bowel, resulting in malabsorption. This procedure is no longer performed due to the high complication rate and frequent need for revisional surgery. Per the American Society for Metabolic & Bariatric Surgery, the JIB is no longer a recommended bariatric surgical procedure. The lessons learned from the JIB include the crucial importance of long-term follow-up and the dangers of a permanent, severe and global malabsorption.¹

Vertical Banded Gastroplasty (VBG): VBG has fallen out of favor as a restrictive procedure for severe obesity, due largely to the advantages of adjustable gastric banding.¹ VBG requires division of the stomach or intestinal resection, while LAGB does not. In addition, the staples used in VBG may break down and cause weight regain, and VBG requires the use of prosthetic mesh that may increase the incidence of stomach stenosis. Centers for Medicare and Medicaid Services (CMS) states in their National Coverage Determination for bariatric surgery for treatment of co-morbid conditions related to morbid obesity that “VBG procedures are essentially no longer performed.”²⁰

Gastric Balloon: Previous endoscopic technologies used to treat obesity endoscopically, such as the gastric balloon, had limited exposure in the U.S. and were removed from the market because of associated complications, such as balloon deflation with migration and resultant small intestinal obstruction.

Gastric Pacing: A number of procedures have been investigated for weight loss surgery but have not been totally accepted by the surgical community. Gastric pacing has been performed in several trials but has not been shown to have any long-term effect and has been abandoned.

Gastric Wrapping: A gastric wrap is minimally invasive surgery and involves folding the stomach in on itself and then the edges are stitched to turn the stomach into a narrow tube, therefore restricting the amount of food that can be consumed. This surgery is new and not widely offered, and there is a paucity of peer-reviewed scientific literature on this procedure.

Coding Implications

This clinical policy references Current Procedural Terminology (CPT®). CPT® is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2023, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

CPT codes that support medical necessity

CPT®* Codes	Description
43644	Laparoscopy, surgical, gastric restrictive procedure; with gastric bypass and Roux-en-Y gastroenterostomy (roux limb 150 cm or less)
43645	Laparoscopy, surgical, gastric restrictive procedure; with gastric bypass and small intestine reconstruction to limit absorption
43770*	Laparoscopy, surgical, gastric restrictive procedure; placement of adjustable gastric restrictive device (eg, gastric band and subcutaneous port components)
43771	Laparoscopy, surgical, gastric restrictive procedure; revision of adjustable gastric restrictive device component only

CPT®* Codes	Description
43772	Laparoscopy, surgical, gastric restrictive procedure; removal of adjustable gastric restrictive device component only
43773	Laparoscopy, surgical, gastric restrictive procedure; removal and replacement of adjustable gastric restrictive device component only
43774	Laparoscopy, surgical, gastric restrictive procedure; removal of adjustable gastric restrictive device and subcutaneous port components
43775	Laparoscopy, surgical, gastric restrictive procedure; longitudinal gastrectomy (ie, sleeve gastrectomy)
43843	Gastric restrictive procedure, without gastric bypass, for morbid obesity; other than vertical-banded gastroplasty
43845	Gastric restrictive procedure with partial gastrectomy, pylorus-preserving duodenoileostomy and ileoileostomy (50 to 100 cm common channel) to limit absorption (biliopancreatic diversion with duodenal switch)
43846	Gastric restrictive procedure, with gastric bypass for morbid obesity; with short limb (150 cm or less) Roux-en-Y gastroenterostomy
43848*	Revision, open, of gastric restrictive procedure for morbid obesity, other than adjustable gastric restrictive device (separate procedure)
43860	Revision of gastrojejunal anastomosis (gastrojejunostomy) with reconstruction, with or without partial gastrectomy or intestine resection; without vagotomy
43865	Revision of gastrojejunal anastomosis (gastrojejunostomy) with reconstruction, with or without partial gastrectomy or intestine resection; with vagotomy
43886	Gastric restrictive procedure, open; revision of subcutaneous port component only
43887	Gastric restrictive procedure, open; removal of subcutaneous port component only
43888	Gastric restrictive procedure, open; removal and replacement of subcutaneous port component only

*Some codes may be used for both medically necessary and not medically necessary indications.

CPT codes that do not support medical necessity

CPT®*	Description
43290	Esophagogastroduodenoscopy, flexible, transoral; with deployment of intragastric bariatric balloon
43291	Esophagogastroduodenoscopy, flexible, transoral; with removal of intragastric bariatric balloon(s)
43632	Gastrectomy, partial, distal; with gastrojejunostomy
43647	Laparoscopy, surgical; implantation or replacement of gastric neurostimulator electrodes, antrum
43648	Laparoscopy, surgical; revision or removal of gastric neurostimulator electrodes, antrum
43842	Gastric restrictive procedure, without gastric bypass, for morbid obesity; vertical-banded gastroplasty
43847	Gastric restrictive procedure, with gastric bypass for morbid obesity; with small intestine reconstruction to limit absorption

43881	Implantation or replacement of gastric neurostimulator electrodes, antrum, open
43882	Revision or removal of gastric neurostimulator electrodes, antrum, open
64590	Insertion or replacement of peripheral or gastric neurostimulator pulse generator or receiver, direct or inductive coupling
64595	Revision or removal of peripheral or gastric neurostimulator pulse generator or receiver

HCPCS codes that support medical necessity

HCPCS Codes	Description
S2083	Adjustment of gastric band diameter via subcutaneous port by injection or aspiration of saline

Reviews, Revisions, and Approvals	Date of Last Revision	Approval Date
Policy Developed	11/17	01/18
Modified Sections I.B. and I.C. requiring a 6 month trial of an exercise/weight loss program, or a medical condition that would supercede the need for such a program, and instead required “previous attempts at weight loss” per the American Society for Metabolic and Bariatric Surgery updated position statement on insurance mandated preoperative weight loss requirements (2016). Removed requirement in II.F. for 6 months of nutritional counseling, while still requiring monthly nutritional counseling until date of surgery. Removed requirement for documented compliance with exercise program in section II.G. Modified II.A.2 removing requirement for specific cardiac testing (stress test, echocardiogram) for high cardiac risk candidates and revised to state they require consultation/evaluation and cardiac clearance from a cardiologist. Replaced cardiac risk qualifiers with that from the reconstructed RCRI, in addition to significant arrhythmias and valvular heart disease. Reworded hypothyroidism screening criteria to require testing if signs/symptoms of hypothyroidism other than obesity. For H. Pylori testing, removed requirement for screening in high prevalence areas, and required treatment if positive. Modified Section VI.A adding that for Biliopancreatic diversion (BPD) procedure (also known as the Scopinaro procedure), BPD with duodenal switch will be considered on a case by case basis. In Section VI.C added that Vertical Banded Gastroplasty (VBG), will also be considered on a case-by-case basis.	09/18	
Revised and reorganized section I.A.1 by BMI and type of procedures considered medically necessary. I.A.1.c, added medically necessary BMI category of > 30- < 35 when criteria is met. Revised I.A.2, a and b, clarifying weight parameters to reflect current terminology. I.A.2.a, removed requirement for co-	12/2020	2/9/2021

Reviews, Revisions, and Approvals	Date of Last Revision	Approval Date
<p>morbidities. I.A.2.c, added comorbidities to this section. Removed II.D, requirement that Tanner stage, or bone age should be completed. III.V, added single anastomosis duodenoileal bypass (SADI); gastric plication/ endoluminal vertical gastroplasty; and endoscopic gastrointestinal bypass devices (barrier devices) as investigational. Updated background. Coding updates</p> <p>In section IV, removed contraindications that are addressed elsewhere in the policy for clarity.</p> <p>Restructured criteria in section I. Medical History. Moved codes 43842 and 43847 to table of codes that do not support medical necessity. Added the following codes to the table that does not support medical necessity: 43647, 43881, 64590.</p> <p>In glycemic control section, changed HbA1C requirement to <7% instead of 6.5-7%. Noted that glycemic control requirement doesn't apply to those who qualify for surgery based on BMI between 30 and 35 with type 2 DM.</p> <p>Changes to section III. Repeat surgeries: Clarified in III.A.1. that repair of complications could include revisions to LSG or RYGB for adults or adolescents, or to BPD-DS for adults. Added to III.A.2 that LSG was an acceptable revision procedure. Edited III.A.2.b. to say that previous surgery was 2 years prior, instead of 3. Added criteria to III.A.2.d. that if the conversion is requested due to removal of an eroded laparoscopic adjustable band, at least two months have passed between the band removal and the subsequent bariatric procedure. Added indication in III.A.3 for conversion of sleeve gastrectomy to Roux-en-Y gastric bypass for the treatment of gastro-esophageal reflux disease (GERD) when anti-reflux medical therapy has been tried and failed. Updated description and background information regarding gastric banding in adolescents.</p>		
<p>Added Coronary artery disease as a comorbidity under A.1.b.ii. Edits made to ICD-10 codes; M54-M54.9 now M54.00-M54.9; T81.1X+-T81.9X now T81.10X+ - T81.9XX+; and T85.59 – T85.59 now T85.590+ - T85.598+. References reviewed and updated. Specified that H. Pylori screening should be conducted using a urea breath test or stool antigen test. Added the following ICD-10 code ranges: M17.0-M17.9, M19.171-M19.179 and M19.271-M19.279. 10/1/20 ICD 10 updates: Replaced category K21.0-K21.9 with K21.00- K21.9. Removed “member” from II.C.4. and II.G. Reworded II.G with no impact on criteria. Replaced “member” with “member/ enrollee” in all other instances. Section I: Added BMI criteria for Asian ethnicity to IA.1.a, IA.1.b and IA.1.c. Added high risk of T2D to list of</p>	12/13/2021	

Reviews, Revisions, and Approvals	Date of Last Revision	Approval Date
<p>severe obesity related complications; added “inadequate glycemic control...” to I.A.1.c.i. Section II: Removed criteria for ECG during cardiac clearance except for high risk; in II.B, removed criteria options of Fasting blood glucose level of ≤ 110 mg/dL and 2-hour postprandial blood glucose concentration of ≤ 140 mg/dL; added note about medical director review if A1C ≥ 8; removed requirement of chest x-ray and specific criteria for PSG, noting that PSG is warranted if OSA screening is positive in II.C. Pulmonary Evaluation; added examples of nutritional tests to be conducted, and that malabsorptive procedures may require further testing to section II. D; removed requirement of 1 year abstinence of drug & alcohol use and urine drug screen if history of abuse in II.F; added “current drug and alcohol abuse” to list of contraindications; added clinically significant GI symptoms should be evaluated & treated prior to surgery in II.I. In III.A.2.e, removed option for non-compliance with post-operative regimen if completing a multidisciplinary bariatric program. In III.A.2.f., removed option for non-compliance. Reworded V, replacing “investigational” with “current medical literature is inadequate to determine the safety, efficacy and long-term outcomes” and added one-anastomosis gastric bypass; endoscopic sleeve gastropasty; transoral endoscopic surgery; vagus nerve blocking (e.g., Maestro) and gastric balloon (e.g., ReShape Duo, Orbera intragastric balloon, Obalon Balloon) to this list. Updated background. Added the following CPT codes as not supporting medical necessity: 43648, 43882, 64595, 0312T, 0313T, 0314T, 0315T, 0316T and 0317T. References reviewed, updated and reformatted. Changed “review date” in the header to “date of last revision” and “date” in the revision log header to “revision date.”</p>		
<p>Annual review. Description updated with no impact on criteria. Criteria I.A. procedures listed with abbreviations with no impact on criteria. Background updated with no impact on criteria. Corrected ICD10 code I10.0 to I10. Section III: updated abbreviations in III.3 with no clinical significance; added indication for SG to RYGB or BPD-DS DS as a bridging procedure for BMI ≥ 50 kg/m² in III.4. References reviewed and updated.</p>	2/20/2023	
<p>Annual review. Updated policy format. Updated policy statement in I, I.A.1, and I.A.1.a. In I.A.1.a.i updated policy statement and BMI threshold to ≥ 35 or ≥ 32.5 kg/m² for South Asian, Southeast Asian, and East Asian adults. In I.A.1.a.ii BMI threshold was updated to "BMI ≥ 30 and < 35 kg/m², or < 27.5 kg/m² and < 32.5 kg/m² for South Asian, Southeast Asian, and East Asian adults and policy statements in I.A.1.a.ii, I.A.1.a.ii.a),</p>	04/2024	

Reviews, Revisions, and Approvals	Date of Last Revision	Approval Date
<p>and c). Moved Type 2 diabetes mellitus (DM) to I.A.1.a.ii.b) as an absolute co-morbidity. Added "pseudotumor cerebri" and "disqualification from other surgeries..." to I.A.1.a.ii.c). Updated policy statement in I.A.1.b.ii. Updated I.B.2 to "Glycemic control evaluation to include A1c and fasting blood glucose". Removed criteria I.B.5.c. requiring prescribed exercise program as part of nutritional counseling. Moved IV. Contraindications to I.C and added "severe cardiac disease with prohibitive anesthetic risks," "uncontrolled and untreated eating disorders (eg, bulimia)," "inability on the part of the patient or parent/guardian to comprehend the risks and benefits of the surgical procedure," and "a medical, psychiatric, psychosocial, or cognitive condition that prevents adherence to postoperative dietary and medication regimens or impairs decisional capacity." Background updated with no clinical impact. Removed deleted CPT codes 0312T-0317T and added CPT codes 43290, 43291, and 43632 to not medically necessary table. Removed ICD-10 codes and table. Annual review. Removed high risk for type 2 DM in Criteria I.A.1.a.ii.c)i). Changed Criteria I.A.1.a.ii.c)ii) from poorly controlled hypertension to hypertension. Removed severe urinary incontinence from Criteria I.A.1.a.ii.c)xii). Updated verbiage in Criteria I.A.1.a.ii.c)xiii) to bone and joint diseases. Removed idiopathic intracranial hypertension from Criteria I.A.1.a.ii.c)xiv) since duplicative since pseudotumor cerebri is in criteria. Added chronic kidney disease in Criteria I.A.1.a.ii.c)xiii). Added infertility in Criteria I.A.1.a.ii.c)xiv). Added polycystic ovarian syndrome in Criteria I.A.1.a.ii.c)xv). Clarified verbiage in Criteria I.A.1.b. Updated Criteria I.A.1.b.ii.d) to state nonalcoholic fatty liver disease or nonalcoholic steatohepatitis. Criteria Clarified verbiage in Criteria I.B.1. and in Criteria I.B.1.a. Updated Criteria I.B.1.b.iv. from glomerular filtration rate (GFR) < 30 mL/min-1 to GFR < 60 mL/min-1. Added Criteria I.B.1.b.vii. to include unstable angina. Added Criteria I.B.1.b.viii. to include recent myocardial infarction (within the past 60 days). Updated Criteria I.B.4. to include thiamine, calcium, and fat-soluble vitamins. In I.B.5., removed requirement for monthly nutritional counseling. Updated verbiage to Criteria I.B.9. to state gastrointestinal (GI) screening and evaluation for clinically significant GI symptoms with documentation of needed treatment prior to bariatric surgery. Removed Criteria I.B.10. for Helicobacter pylori screening. Minor rewording in Criteria I.B.11. Updated Criteria III.B. to include other names of procedure for clarification. Minor rewording in Criteria III.K. Removed one-anastomosis gastric bypass in Criteria III.L. since</p>		

Reviews, Revisions, and Approvals	Date of Last Revision	Approval Date
duplicative. Updated Background with no impact on criteria. References reviewed and updated. Reviewed by internal and external specialists.		

References

1. Lim RB. Bariatric procedures for the management of severe obesity: Descriptions. UpToDate. www.uptodate.com. Published April 13, 2023. Accessed February 26, 2024.
2. Pratt JSA, Browne A, Browne NT, et al. ASMBS pediatric metabolic and bariatric surgery guidelines, 2018. *Surg Obes Relat Dis*. 2018;14(7):882 to 901. doi:10.1016/j.soard.2018.03.019
3. Lim RB. Bariatric surgery for management of obesity: Indications and preoperative preparation. UpToDate. www.uptodate.com. Published January 04, 2023. Accessed February 20, 2024.
4. Eisenberg D, Shikora SA, Aarts E, et al. 2022 American Society of Metabolic and Bariatric Surgery (ASMBS) and International Federation for the Surgery of Obesity and Metabolic Disorders (IFSO) Indications for Metabolic and Bariatric Surgery [published correction appears in *Obes Surg*. 2022 Nov 29;:]. *Obes Surg*. 2023;33(1):3 to 14. doi:10.1007/s11695-022-06332-1
5. National Institute for Health and Care Excellence (NICE). Single-anastomosis duodeno-ileal bypass with sleeve gastrectomy for treating morbid obesity. Interventional procedures guidance [IPG569]. <https://www.nice.org.uk/guidance/IPG569>. Published November 23, 2016. Accessed February 28, 2024.
6. Inge TH. Surgical management of severe obesity in adolescents. UpToDate. www.uptodate.com. Published August 08, 2023. Accessed February 29, 2024.
7. Aminian A, Chang J, Brethauer SA, Kim JJ; American Society for Metabolic and Bariatric Surgery Clinical Issues Committee. ASMBS updated position statement on bariatric surgery in class I obesity (BMI 30 to 35 kg/m²). *Surg Obes Relat Dis*. 2018;14(8):1071 to 1087. doi:10.1016/j.soard.2018.05.025
8. Styne DM, Arslanian SA, Connor EL, et al. Pediatric Obesity-Assessment, Treatment, and Prevention: An Endocrine Society Clinical Practice Guideline. *J Clin Endocrinol Metab*. 2017;102(3):709 to 757. doi:10.1210/jc.2016-2573
9. Health Technology Assessment. Comparative effectiveness review of bariatric surgeries for treatment of obesity in adolescents. Hayes. www.hayesinc.com. Published January 21, 2019 (annual review January 20, 2022). Accessed February 26, 2024.
10. Armstrong SC, Bolling CF, Michalsky MP, Reichard KW; SECTION ON OBESITY, SECTION ON SURGERY. Pediatric Metabolic and Bariatric Surgery: Evidence, Barriers, and Best Practices. *Pediatrics*. 2019;144(6):e20193223. doi:10.1542/peds.2019-3223
11. Svanevik M, Ristad H, Hofsvø D, et al. Perioperative Outcomes of Proximal and Distal Gastric Bypass in Patients with BMI Ranged 50 to 60 kg/m(2)--A Double-Blind, Randomized Controlled Trial. *Obes Surg*. 2015;25(10):1788 to 1795. doi:10.1007/s11695-015-1621-y

12. Jensen MD, Ryan DH, Apovian CM, et al. 2013 AHA/ACC/TOS guideline for the management of overweight and obesity in adults: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and The Obesity Society [published correction appears in *J Am Coll Cardiol*. 2014 Jul 1;63(25 Pt B):3029 to 3030]. *J Am Coll Cardiol*. 2014;63(25 Pt B):2985 to 3023. doi:10.1016/j.jacc.2013.11.004
13. Iannelli A, Schneck AS, Noel P, Ben Amor I, Krawczykowski D, Gugenheim J. Re-sleeve gastrectomy for failed laparoscopic sleeve gastrectomy: a feasibility study. *Obes Surg*. 2011; 21(7):832 to 835. doi:10.1007/s11695-010-0290-0
14. AlSabah S, Alsharqawi N, Almulla A, et al. Approach to Poor Weight Loss After Laparoscopic Sleeve Gastrectomy: Re-sleeve Vs. Gastric Bypass. *Obes Surg*. 2016;26(10):2302 to 2307. doi:10.1007/s11695-016-2119-y
15. Fobi MA, Lee H. The surgical technique of the Fobi-Pouch operation for obesity (the transected silastic vertical gastric bypass). *Obes Surg*. 1998;8(3):283-288. doi:10.1381/096089298765554485
16. Fried M, Dolezalova K, Sramkova P. Adjustable gastric banding outcomes with and without gastrogastic imbrication sutures: a randomized controlled trial. *Surg Obes Relat Dis*. 2011;7(1):23 to 31. doi:10.1016/j.soard.2010.09.018
17. Sharma S, Narwaria M, Cottam DR, Cottam S. Randomized double-blinded trial of laparoscopic gastric imbrication v laparoscopic sleeve gastrectomy at a single Indian institution. *Obes Surg*. 2015;25(5):800 to 804. doi:10.1007/s11695-014-1497-2
18. Norén E, Forssell H. Aspiration therapy for obesity; a safe and effective treatment. *BMC Obes*. 2016;3:56. Published 2016 Dec 28. doi:10.1186/s40608-016-0134-0
19. Pennestrì F, Sessa L, Prioli F, et al. Single anastomosis duodenal-ileal bypass with sleeve gastrectomy (SADI-S): experience from a high-bariatric volume center. *Langenbecks Arch Surg*. 2022;407(5):1851 to 1862. doi:10.1007/s00423-022-02501-z
20. National coverage determination: bariatric surgery for treatment of co-morbid conditions related to morbid obesity (100.1). Centers for Medicare and Medicaid Services Web site. <https://www.cms.gov/medicare-coverage-database/new-search/search.aspx>. Published September 24, 2013. Accessed February 22, 2024.
21. Buchwald H; Consensus Conference Panel. Bariatric surgery for morbid obesity: health implications for patients, health professionals, and third-party payers. *J Am Coll Surg*. 2005;200(4):593 to 604. doi:10.1016/j.jamcollsurg.2004.10.039
22. Spear BA, Barlow SE, Ervin C, et al. Recommendations for treatment of child and adolescent overweight and obesity. *Pediatrics*. 2007;120 Suppl 4:S254 to S288. doi:10.1542/peds.2007-2329F
23. Cohn SL, Fleisher LA. Evaluation of cardiac risk prior to noncardiac surgery. UpToDate. www.uptodate.com. Published July 26, 2023. Accessed February 20, 2024.
24. Colquitt JL, Pickett K, Loveman E, Frampton GK. Surgery for weight loss in adults. *Cochrane Database Syst Rev*. 2014;2014(8):CD003641. Published 2014 Aug 8. doi:10.1002/14651858.CD003641.pub4
25. Thompson CC, Abu Dayyeh BK, Kushner R, et al. Percutaneous Gastrostomy Device for the Treatment of Class II and Class III Obesity: Results of a Randomized Controlled Trial. *Am J Gastroenterol*. 2017;112(3):447 to 457. doi:10.1038/ajg.2016.500
26. Davis C, Tait G, Carroll J, Wijeyesundera DN, Beattie WS. The Revised Cardiac Risk Index in the new millennium: a single-centre prospective cohort re-evaluation of the original

- variables in 9,519 consecutive elective surgical patients. *Can J Anaesth.* 2013; 60(9):855 to 863. doi:10.1007/s12630-013-9988-5
27. Health Technology Assessment. Intra-gastric balloons for treatment of obesity. Hayes. www.hayesinc.com. Published March 29, 2018 (annual review March 16, 2022). Accessed February 26, 2024.
 28. Kim JJ, Rogers AM, Ballem N, Schirmer B; American Society for Metabolic and Bariatric Surgery Clinical Issues Committee. ASMBS updated position statement on insurance mandated preoperative weight loss requirements. *Surg Obes Relat Dis.* 2016;12(5):955 to 959. doi:10.1016/j.soard.2016.04.019
 29. Mechanick JI, Apovian C, Brethauer S, et al. Clinical Practice Guidelines for the Perioperative Nutrition, Metabolic, and Nonsurgical Support of Patients Undergoing Bariatric Procedures - 2019 Update: Cosponsored by American Association of Clinical Endocrinologists/American College of Endocrinology, The Obesity Society, American Society for Metabolic and Bariatric Surgery, Obesity Medicine Association, and American Society of Anesthesiologists. *Obesity (Silver Spring).* 2020;28(4):O1 to O58. doi:10.1002/oby.22719
 30. Health Technology Assessment. Comparative effectiveness review of mini gastric bypass—one anastomosis gastric bypass for the treatment of obesity: a review of reviews. Hayes. www.hayesinc.com. Published May 30, 2019 (annual review January 20, 2023). Accessed February 29, 2024.
 31. Kallies K, Rogers AM; American Society for Metabolic and Bariatric Surgery Clinical Issues Committee. American Society for Metabolic and Bariatric Surgery updated statement on single-anastomosis duodenal switch. *Surg Obes Relat Dis.* 2020;16(7):825 to 830. doi:10.1016/j.soard.2020.03.020
 32. Michalsky M, Reichard K, Inge T, Pratt J, Lenders C; American Society for Metabolic and Bariatric Surgery. ASMBS pediatric committee best practice guidelines. *Surg Obes Relat Dis.* 2012;8(1):1 to 7. doi:10.1016/j.soard.2011.09.009
 33. National Institute for Health and Care Excellence (NICE). Single-anastomosis duodeno-ileal bypass with sleeve gastrectomy for treating morbid obesity. Interventional procedures guidance [IPG569]. <https://www.nice.org.uk/guidance/IPG569>. Published November 23, 2016. Accessed February 28, 2024.
 34. Rosenthal RJ, Szomstein S, Menzo EL. Laparoscopic sleeve gastrectomy. UpToDate. www.uptodate.com. Published May 16, 2022. Accessed February 26, 2024.
 35. National Clinical Guideline Centre (UK). *Obesity: Identification, Assessment and Management of Overweight and Obesity in Children, Young People and Adults*. London: National Institute for Health and Care Excellence (UK); November 2014. National Institute for Health and Care Excellence (NICE); November 2014.
 36. Carter J, Chang J, Birriel TJ, et al. ASMBS position statement on preoperative patient optimization before metabolic and bariatric surgery. *Surg Obes Relat Dis.* 2021;17(12):1956-1976. doi:10.1016/j.soard.2021.08.024
 37. Kushner RF, Herron DM, Herrington H. Bariatric surgery: Postoperative nutritional management. UpToDate. www.uptodate.com. Published August 04, 2022. Accessed February 29, 2024.
 38. Parikh M, Chung M, Sheth S, et al. Randomized pilot trial of bariatric surgery versus intensive medical weight management on diabetes remission in type 2 diabetic patients who

do NOT meet NIH criteria for surgery and the role of soluble RAGE as a novel biomarker of success. *Ann Surg.* 2014;260(4):617 to 624. doi:10.1097/SLA.0000000000000919

39. Halvorsen S, Mehilli J, Cassese S, et al. 2022 ESC Guidelines on cardiovascular assessment and management of patients undergoing non-cardiac surgery: Developed by the task force for cardiovascular assessment and management of patients undergoing non-cardiac surgery of the European Society of Cardiology (ESC) Endorsed by the European Society of Anaesthesiology and Intensive Care (ESAIC). *European Heart Journal.* 2022;43(39):3826 to 3924.