

Clinical Policy: Neuromuscular and Peroneal Nerve Electrical Stimulation (NMES)

Reference Number: PA.CP.MP.48

Plan Effective Date: 01/2023

Date of Last Revision: 06/2024

[Coding Implications](#)

[Revision Log](#)

Description

This policy describes the medical necessity requirements for the use of neuromuscular electrical stimulation (NMES) and functional electrical stimulation (FES).

Policy/Criteria

- I. It is the policy of PA Health & Wellness (PHW)® that neuromuscular electrical stimulation (NMES) is **medically necessary** when used as one component of a comprehensive rehab program for the treatment of disuse atrophy when the nerve supply to the atrophied muscle is intact and has any of the following atrophy indications:
 - A. Contractures due to burn scarring;
 - B. Previous casting or splinting of a limb;
 - C. Major knee surgery with failure to respond to physical therapy;
 - D. Recent hip replacement and NMES will be used until physical therapy begins.
- II. It is the policy of PHW that functional electrical stimulation (FES) is **medically necessary** for spinal cord injury (SCI) when all of the following criteria are met:
 - A. The member/enrollee has brisk muscle contraction to stimulation and sensory perception of electrical stimulation sufficient for muscle contraction;
 - B. At least six months have passed since recovery from spinal cord injury and restorative surgery;
 - C. Member/enrollee is highly motivated, committed, and has the cognitive ability to use FES devices for walking;
 - D. Successful completion of a training program consisting of at least 32 physical therapy sessions with the device over a three-month period;
 - E. Member/enrollee demonstrates a willingness to use the device long-term;
 - F. None of the following contraindications are present:
 1. Cardiac pacemaker;
 2. Severe scoliosis or severe osteoporosis;
 3. Skin disease or cancer at area of stimulation;
 4. Irreversible contracture;
 5. Autonomic dysreflexia;
 - G. If lower extremity FES is requested, all of the following:
 1. Intact lower motor units (L1 and below, including both muscle and peripheral nerve);
 2. Muscle and joint stability adequate for weight bearing at upper and lower extremities and can demonstrate balance and control to maintain an upright support posture independently;
 3. Transfers independently and demonstrates independent standing tolerance for at least three minutes;
 4. Demonstrates hand and finger function to manipulate controls;
 5. No hip and knee degenerative disease and no history of long bone fracture secondary to osteoporosis.

III. It is the policy of PHW that peroneal nerve stimulators (e.g., NESS L300, NESS L300 Plus, L300 Go System, WalkAide, Odstock [ODFS[®]] Dropped Foot Stimulator) are **medically necessary** for incomplete spinal cord injury.

IV. It is the policy of PHW that peroneal nerve stimulators (e.g., NESS L300, NESS L300 Plus, L300 Go System, WalkAide, Odstock [ODFS[®]] Dropped Foot Stimulator) have not been proven safe and effective for any indication other than incomplete spinal cord injury, including, but not limited to: foot drop in cerebral palsy, multiple sclerosis, traumatic brain injury, or stroke.

V. It is the policy of PHW that neuromuscular electrical stimulation for any other indication (e.g., idiopathic scoliosis, heart failure) is not proven safe and effective.

Background

Neuromuscular electrical stimulation (NMES) involves the use of a device which transmits an electrical impulse to the skin over selected muscle groups by way of electrodes.^{1,2} There are two broad categories of NMES. The first type of device stimulates the muscle when the patient is in a resting state to treat muscle atrophy.¹ The second type, known as functional electrical stimulation (FES), is used to enhance functional activity of neurologically impaired patients.¹

NMES can be performed at low, medium, or high intensity to elicit mild, moderate, or strong muscle contractions. NMES can be performed on upper or lower limbs. When used at very low intensity to stimulate barely perceptible contractions, this technique is referred to as threshold NMES or threshold electrical stimulation (TES).^{1,4} Regardless of the intensity of NMES, patients are encouraged to exercise the affected muscles voluntarily to maintain and improve strength and function. For chronic disorders, this exercise may be in the form of regular participation in sports activities. For acute conditions, such as rehabilitation shortly after surgery or a stroke, patients must often undergo intensive physical and occupational therapy.^{1,4}

FES is the application of electrical stimulation that can be used to activate muscles of the upper or lower limbs to produce functional movement patterns, such as standing and walking in patients with paraplegia.^{1,4} Although FES is used to treat the effects of upper motor neuron lesions, it is not normally suitable for patients with lower motor neuron lesions.¹⁰ FES can also be used therapeutically for cycling of the upper and/or lower limbs, with the goal of strengthening to produce functional movement patterns.⁹

FES has been shown to strengthen muscles, improve circulation, heal tissue, slow muscle atrophy, and reduce pain and spasticity.⁴

There is evidence from preliminary studies that FES can improve gait in some patients; however, additional larger randomized trials are needed.^{3,2,6}

The only settings where skilled therapists can provide both types of NMES services are inpatient hospitals, outpatient hospitals, comprehensive outpatient rehabilitation facilities, and outpatient

rehabilitation facilities. The physical therapy needed to perform these services requires that the patient be in a one-on-one training program.¹

Coding Implications

This clinical policy references Current Procedural Terminology (CPT®). CPT® is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2023, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS®* Codes	Description
E0745	Neuromuscular stimulator, electronic shock unit
E0764	Functional neuromuscular stimulation, transcutaneous stimulation of sequential muscle groups of ambulation with computer control, used for walking by spinal cord injured, entire system, after completion of training program
E0770	Functional electrical stimulator, transcutaneous stimulation of nerve and/or muscle groups, any type, complete system, not otherwise specified

HCPCS codes that do not support coverage criteria

HCPCS Codes	Description
E0744	Neuromuscular stimulator for scoliosis

Reviews, Revisions, and Approvals	Revision Date	Approval Date
Original approval date 09/11. References reviewed and updated. Template update and approved 12/11. References reviewed and updated. Approved with no changes 9/12-9/14.	09/11	09/11
References reviewed and updated. Approved by MPC. Coding update only.	09/15	09/15
References reviewed and updated. Approved by MPC. No changes.	09/16	09/16
References reviewed and updated. Approved by MPC. No changes.	07/17	07/17
References reviewed and updated. Approved by MPC. No changes.	07/18	07/18
References reviewed and updated. Approved by MPC. No changes.	07/19	07/19
References reviewed and updated. Approved by MPC. No changes.	07/20	07/20
Transitioned to CNC template. Replaced “members” with “members/enrollees” in all instances.	12/20	
Annual review completed. References reviewed and updated. Changed “review date” in the header to “date of last revision” and “date” in the revision log header to “revision date.” Integrated NMES, FES, and peroneal stimulator criteria from PA.CP.MP.107 DME and Legacy WellCare Neuromuscular Electrical Stimulation (NMES)	07/21	07/21

Reviews, Revisions, and Approvals	Revision Date	Approval Date
PA.CP.MP.48 policy. Renamed to “Neuromuscular and Peroneal Nerve Electrical Stimulation.” Added section III and IV criteria. Added code E0744 to “HCPCS codes that do not support coverage criteria.” Specialist reviewed.		
Annual review. Criteria IV. verbiage updated for clarity. Background updated with no impact on criteria. References reviewed and updated. Specialist reviewed.	07/22	07/22
New Plan specific policy created	12/19/2022	
Annual review completed. Combined criteria applicable to LE units into section II.G. Additional contraindications added to Section F. Minor rewording with no clinical significance. Background updated with no impact to criteria. ICD-10-CM Diagnosis Code table removed. References reviewed and updated. Internal specialist reviewed.	07/2023	
Annual review. Removed contraindications under II.F. including uncontrolled cardiac arrhythmias, unstable angina, joint replacement in a location targeted by FES and seizure disorder. Background updated with no impact on criteria. References reviewed and updated. Reviewed by external specialist.	06/2024	8/21/24

References

1. National coverage determination. Neuromuscular electrical stimulation (NMES) (160.12). Centers for Medicare and Medicaid Services website. <http://www.cms.hhs.gov/mcd/search.asp>. Published October 01, 2006. Accessed May 7, 2024.
2. Health Technology Assessment. Functional electrical stimulation for foot drop in acute or subacute phases of stroke recovery. Hayes. www.hayesinc.com. Published June 01, 2022 (annual review June 30, 2023). Accessed May 7, 2024.
3. Health Technology Assessment. Functional electrical stimulation (FES) for treatment of foot drop in multiple sclerosis patients. Hayes. www.hayesinc.com. Published November 17, 2021 (annual review November 14, 2023). Accessed May 7, 2024.
4. Doucet BM, Lam A, Griffin L. Neuromuscular electrical stimulation for skeletal muscle function. *Yale J Biol Med*. 2012;85(2):201 to 215.
5. Celli BR. Pulmonary rehabilitation. UpToDate. www.uptodate.com. Updated May 3, 2024. Accessed May 7, 2024.
6. Olek MJ, Narayan RN, Frohman EM, Frohman TC. Symptom management of multiple sclerosis in adults. UpToDate. www.uptodate.com. Updated October 5, 2023. Accessed May 7, 2024.
7. de Freitas GR, Szpoganicz C, Ilha J. Does Neuromuscular Electrical Stimulation Therapy Increase Voluntary Muscle Strength After Spinal Cord Injury? A Systematic Review. *Top Spinal Cord Inj Rehabil*. 2018;24(1):6 to 17. doi:10.1310/sci16-00048
8. Jones S, Man WD, Gao W, Higginson IJ, Wilcock A, Maddocks M. Neuromuscular electrical stimulation for muscle weakness in adults with advanced disease. *Cochrane*

Database Syst Rev. 2016;10(10):CD009419. Published 2016 Oct 17.
doi:10.1002/14651858.CD009419.pub3

9. Sansare A, Harrington AT, Wright H, Alesi J, Behboodi A, Verma K, Lee SCK. Aerobic Responses to FES-Assisted and Volitional Cycling in Children with Cerebral Palsy. *Sensors (Basel)*. 2021 Nov 15;21(22):7590. doi: 10.3390/s21227590. PMID: 34833666; PMCID: PMC8622737.
10. National Institute for Health and Care Excellence (NICE). Functional electrical stimulation for drop foot of central neurological origin [IPG278].
<https://www.nice.org.uk/guidance/ipg278/chapter/2-The-procedure>. Published January 28, 2009. Accessed May 7, 2024.