

**Prior Authorization Review Panel
MCO Policy Submission**

A separate copy of this form must accompany each policy submitted for review.
Policies submitted without this form will not be considered for review.

Plan: PA Health & Wellness (PHW)	Submission Date: 10/01/2024
Policy Number: PA.CP.MP.53	Effective Date: 01/01/2018 Revision Date: 09/2024
Policy Name: Ferriscan R2-MRI	
Type of Submission – <u>Check all that apply</u>: ☐ New Policy ☐ Revised Policy* ☐ Annual Review – No Revisions ☐ Statewide PDL ☒ Retiring Policy	
*All revisions to the policy <u>must</u> be highlighted using track changes throughout the document. Please provide any clarifying information for the policy below: Policy being retired with no replacement indicated.	
Name of Authorized Individual (Please type or print): Dr. Craig A. Butler, MD.	Signature of Authorized Individual: <i>Craig A Butler MD</i>

Clinical Policy: Ferriscan R2-MRI

Reference Number: PA.CP.MP.53

Effective Date: 10/18

Date of Last Revision: 09/2024

[Coding Implications](#)
[Revision Log](#)

Description

FerriScan[®] R2-MRI is a magnetic resonance imaging (MRI)-based solution for measuring liver iron concentration (LIC) in patients with iron overload.

Policy/Criteria

- I. It is the policy of PA Health & Wellness[®] that the FerriScan[®] R2-MRI is **medically necessary** for the measurement of liver iron concentration in suspected cases of iron overload due to the following conditions:
 - A. Hereditary hemochromatosis;
 - B. Iron-loading anemias with or without multiple transfusions:
 1. Thalassemia major or thalassemia intermedia;
 2. Sideroblastic anemia;
 3. Chronic hemolytic anemias (e.g., sickle cell disease);
 4. Inherited or acquired aplastic anemia;
 5. Myelodysplastic syndromes;
 6. Hematopoietic stem cell transplantation;
 - C. Dietary iron overload;
 - D. Iron overload in liver diseases:
 1. Hepatitis C or B;
 2. Alcohol-induced liver disease;
 3. Porphyria cutanea tarda;
 4. Fatty liver disease;
 5. Gestational alloimmune liver disease;
 6. Suspicion of rare genetic variants affecting iron absorption or distribution⁴;
 - E. Neonatal iron overload;
 - F. Aceruloplasminemia;
 - G. Repeated hemin infusions for acute porphyrias;
 - H. Hemodialysis for end stage renal failure.

Background

Iron overload is a potentially life-threatening problem that is commonly overlooked due to nonspecific symptoms that tend to develop slowly over time. Excess iron does not only affect the liver, but can also accumulate in, and damage other organs like the heart, skin and endocrine organs, as well as joints. Clinical issues resulting from excess iron include tissue damage, inflammation, and fibrosis. Left untreated, iron overload can result in organ toxicity, end-organ damage and dysfunction due to oxidative stress resulting in excess oxygen radicals and injury from tissue peroxidation. Once identified, iron overload is treated with therapeutic phlebotomy and chelation therapy as well as exchange transfusion in sickle cell disease.^{4,5}

Disorders associated with hepatic iron deposition include^{4,5}:

- Hereditary hemochromatosis;

CLINICAL POLICY

Ferriscan R2-MRI

- Syndromes of ineffective erythropoiesis such as beta thalassemia, sideroblastic anemia and other inherited anemias;
- Chronic liver disease;
- Gestational alloimmune liver disease
- Alcoholic liver disease;
- Hepatitis;
- Nonalcoholic fatty liver disease;
- Cirrhosis;
- Wilson disease;
- Porphyria cutanea tarda;
- Hematopoietic stem cell transplantation,
- Myelodysplastic syndrome,
- Dialysis;
- Blood transfusions for sickle cell disease.

FerriScan[®] is a non-invasive technology based on magnetic resonance imaging (MRI). It has a high sensitivity and specificity for the measurement of liver iron concentration (LIC) over the entire range encountered in clinical practice. It can be set up on most 1.5 Tesla MRI scanners, which are the most common type of clinical scanner, and it was announced by Resonance Health in 2022 that FerriScan is now available on 3 Tesla MRI machines.¹⁶ FerriScan works by making a map of the LIC and calculating the mean LIC. The results are unaffected by the presence of fibrosis or cirrhosis. Image data is acquired on an MRI scanner and is electronically transmitted to a data analysis center. All data is analyzed to ensure correct acquisition, and the LIC results are transmitted back to the originating MRI center.

Measurements have been shown to have a high degree of sensitivity and specificity for LIC measured by biopsy. Ferriscan has become increasingly accurate in the determination of hepatic and cardiac iron deposition and is replacing direct tissue biopsy in the assessment of iron overload.⁴ FerriScan images give information on liver iron distribution. The mean LIC value given in the FerriScan report is then used to guide chelation therapy.

The operational principle of the R2-MRI Analysis System is based on fitting signal decay curves to the image signal intensities (e.g., of the liver) at the different echo times for the magnetic resonance data set on a voxel-by-voxel (3-D pixel) basis to determine transverse relaxation rate (R2) images. These may be further transformed by a defined calibration to provide a quantitative measure of liver iron concentrations.

Although magnetic resonance evaluation for hepatic iron concentration is improved compared with older programs, this type of imaging will not detect cellular liver damage due to iron overload.

The American College of Radiology's 2020 Practice Parameter for the performance of MRI of the liver states that indications for MRI of the liver include, but are not limited to, evaluation and noninvasive quantification of iron, fat, and fibrosis in chronic liver disease, such as hemochromatosis, hemosiderosis, nonalcoholic steatohepatitis (NASH), and hepatitis in adults and pediatric patients. Additionally, multiple studies have confirmed the clinical utility of R2-

CLINICAL POLICY

Ferriscan R2-MRI



MRI in the measurement of LIC for iron-overloading conditions such as thalassemia and sickle cell anemia.⁹ A study of R2-MRI results vs. simulated liver biopsy results found R2-MRI to be superior to liver biopsy for serial LIC observations.¹⁰ Furthermore, a review of the current state of liver iron quantification by MRI states that R2-MRI provides validated measurement of LIC and has advantages over liver biopsy, in that it is non-invasive.¹¹

The R2-MRI Analysis System (Inner Vision Biometrics PTY LTD) received 510(k) clearance (K043271) from the United States Food and Drug Administration (FDA) on January 21, 2005. In January 2013, the FDA authorized the FerriScan R2-MRI to be marketed as an imaging companion diagnostic device for the safe and effective use of Exjade in patients with non-transfusion-dependent thalassemia.

Coding Implications

This clinical policy references Current Procedural Terminology (CPT®). CPT® is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2022, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

CPT® Codes	Description
76498	Unlisted MRI procedure

Reviews, Revisions, and Approvals	Date	Approval Date
References reviewed and updated. Codes reviewed and updated.	10/18	10/18
Changed “thalassemia major and thalassemia intermedia” to “thalassemia major or thalassemia intermedia.” Changed “hepatitis C and B” to “hepatitis C or B”	01/19	02/19
Replaced codes D61.89 and D61.9 with expanded range of D61.01-D61.9. Replaced “member” with “member/enrollee/enrollee” in all instances. References reviewed and updated. Reviewed by specialist.	12/2020	1/18/2021
References reviewed and updated. Reviewed by specialist.	10/2021	
Annual review. Added “Hemodialysis for end stage renal failure” as an indication. Changed “review date” in the header to “date of last revision” and “date” in the revision log header to “revision date.” Updated background with no clinical significance or impact on criteria. ICD-10 codes removed from policy. References reviewed and updated. Reviewed by specialist.	02/22/2023	
Annual Review. Added criterion I.B.6. Hematopoietic stem cell transplantation as an iron-loading anemia indication. Added criterion I.G.6., “Suspicion of rare genetic variants affecting iron absorption or distribution.” Updated background with no clinical significance or	10/2023	01/2024

Reviews, Revisions, and Approvals	Date	Approval Date
impact to criteria. Reviewed and updated references. Reviewed and updated CPT code description. Reviewed by external specialist		
Policy being retired with no replacement indicated.	09/2024	9/30/2024

References

1. The American College of Radiology. ACR–SAR–SPR practice parameter for the performance of magnetic resonance imaging (MRI) of the liver. <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/MR-Liver.pdf>. Published 2020. Accessed August 30, 2023.
2. FerriScan® R2-MRI Fact Sheet. Resonance Health. <https://ferriscan.com/wp-content/uploads/2019/10/FerriScan-Fact-Sheet-May-2019.pdf>. Accessed September 26, 2023.
3. Fischer R, Harmatz PR. Non-invasive assessment of tissue iron overload. *Hematology Am Soc Hematol Educ Program*. 2009;215 to 221. doi:10.1182/asheducation-2009.1.215
4. Bacon BR, Kwiatkowski JL. Approach to the Patient with Suspected Iron Overload. UpToDate. www.uptodate.com. Updated June 09, 2022. Accessed August 30, 2023.
5. Fiel, MI. Methods to determine hepatic iron content. UpToDate. www.uptodate.com. Updated January 24, 2022. Accessed August 30, 2023.
6. Taher A, Musallam KM, El Rassi F, et al. Levels of non-transferrin-bound iron as an index of iron overload in patients with thalassaemia intermedia. *Br J Haematol*. 2009;146(5):569 to 572. doi:10.1111/j.1365-2141.2009.07810.x
7. Kwiatkowski JL, Kim HY, Thompson AA, et al. Chelation use and iron burden in North American and British thalassemia patients: a report from the Thalassemia Longitudinal Cohort. *Blood*. 2012;119(12):2746 to 2753. doi:10.1182/blood-2011-04-344507
8. Wood JC, Pressel S, Rogers ZR, et al. Liver iron concentration measurements by MRI in chronically transfused children with sickle cell anemia: baseline results from the TWiTCH trial. *Am J Hematol*. 2015;90(9):806 to 810. doi:10.1002/ajh.24089
9. Wood JC, Zhang P, Rienhoff H, Abi-Saab W, Neufeld EJ. Liver MRI is more precise than liver biopsy for assessing total body iron balance: a comparison of MRI relaxometry with simulated liver biopsy results. *Magn Reson Imaging*. 2015;33(6):761 to 767. doi:10.1016/j.mri.2015.02.016
10. Hernando D, Levin YS, Sirlin CB, Reeder SB. Quantification of liver iron with MRI: state of the art and remaining challenges. *J Magn Reson Imaging*. 2014;40(5):1003 to 1021. doi:10.1002/jmri.24584
11. Tamary H, Dgany O. Congenital Dyserythropoietic Anemia Type 1. <https://www.ncbi.nlm.nih.gov/books/NBK5313/>. Published April 21, 2009. Accessed August 30, 2023.
12. d'Assignies G, Paisant A, Bardou-Jacquet E, et al. Non-invasive measurement of liver iron concentration using 3-Tesla magnetic resonance imaging: validation against biopsy. *Eur Radiol*. 2018;28(5):2022 to 2030. doi:10.1007/s00330-017-5106-3
13. Jhaveri KS, Kannengiesser SAR, Ward R, Kuo K, Sussman MS. Prospective Evaluation of an R2* Method for Assessing Liver Iron Concentration (LIC) Against FerriScan: Derivation of the Calibration Curve and Characterization of the Nature and Source of

- Uncertainty in the Relationship. *J Magn Reson Imaging*. 2019;49(5):1467 to 1474. doi:10.1002/jmri.26313
14. Kowdley KV, Brown KE, Ahn J, Sundaram V. ACG Clinical Guideline: Hereditary Hemochromatosis [published correction appears in *Am J Gastroenterol*. 2019 Dec;114(12):1927]. *Am J Gastroenterol*. 2019;114(8):1202 to 1218. doi:10.14309/ajg.0000000000000315
 15. Ghoti H, Rachmilewitz EA, Simon-Lopez R, et al. Evidence for tissue iron overload in long-term hemodialysis patients and the impact of withdrawing parenteral iron. *Eur J Haematol*. 2012;89(1):87 to 93. doi:10.1111/j.1600-0609.2012.01783.x
 16. Resonance Health. FerriScan[®] Now Available on 3 Tesla (3T) MRI Scanning Machines. <https://company-announcements.afr.com/asx/rht/1bf849e5-0b9e-11ed-8ad3-be99ac8a3e10.pdf>. July 25, 2022. Accessed August 30, 2023.