

Prior Authorization Review Panel

CHC-MCO Policy Submission

A separate copy of this form must accompany each policy submitted for review. Policies submitted without this form will not be considered for review.

Plan: PA Health & Wellness	Submission Date: 11/01/2022	
Policy Number: PA.CP.PHAR.332	Effective Date: 01/2020 Revision Date: 10/2022	
Policy Name: Pasireotide (Signifor, Signifor LAR)		
Type of Submission – <u>Check all that apply</u> :		
 □ New Policy ✓ Revised Policy* □ Annual Review - No Revisions □ Statewide PDL - Select this box when submitting policies for when submitting policies for drug classes included on the S 	-	
*All revisions to the policy <u>must</u> be highlighted using track chan	ges throughout the document.	
Please provide any changes or clarifying information for the poli	cy below:	
4Q 2022 annual review: for acromegaly, added confirmatory diagnostic requirements (IGF-I or GH) per PS/ES practice guidelines; references reviewed and updated.		
Name of Authorized Individual (Please type or print):	Signature of Authorized Individual:	
Venkateswara R. Davuluri, MD	Raulun	



Clinical Policy: Pasireotide (Signifor, Signifor LAR)

Reference Number: PA.CP.PHAR.332 Effective Date: 01/2018 Last Review Date: 10/2022

Coding Implications Revision Log

Description

Pasireotide (Signifor[®], Signifor[®] LAR) is a somatostatin analog.

FDA Approved Indication(s)

Signifor and Signifor LAR are indicated for the treatment of patients with Cushing's disease for whom pituitary surgery is not an option or has not been curative. Signifor is specifically indicated in adults.

Signifor LAR is also indicated for the treatment of patients with:

• Acromegaly who have had an inadequate response to surgery and/or for whom surgery is not an option

Policy/Criteria

It is the policy of PA Health and Wellness[®] that Signifor and Signifor LAR is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

- A. Acromegaly (must meet all):
 - 1. Diagnosis of acromegaly as evidenced by one of the following (a or b):
 - a. Pre-treatment insulin-like growth factor-I (IGF-I) level above the upper limit of normal based on age and gender for the reporting laboratory;
 - b. Serum growth hormone (GH) level $\geq 1 \ \mu g/mL$ after a 2-hour oral glucose tolerance test;
 - 2. Request is for Signifor LAR;
 - 3. Prescribed by or in consultation with an endocrinologist;
 - 4. Age \geq 18 years;
 - 5. Inadequate response to surgical resection or pituitary irradiation (*see Appendix D*), or member is not a candidate for such treatment;
 - 6. Dose does not exceed (a and b):
 - a. 60 mg every 4 weeks;
 - b. 1 vial every 4 weeks.

Approval duration: 6 months

B. Cushing's Disease (must meet all):

- 1. Diagnosis of Cushing's disease;
- 2. Prescribed by or in consultation with an endocrinologist;
- 3. Age \geq 18 years;
- 4. Member meets one of the following (a or b):
 - a. Pituitary surgery was not curative;
 - b. Ineligible for pituitary surgery;
- 5. Dose does not exceed one of the following (a or b):



- a. Signifor (i and ii):
 - i. 1.8 mg per day;
 - ii. 2 ampules per day;
- b. Signifor LAR (i and ii):
 - i. 40 mg every 4 weeks
 - ii. 1 vial every 4 weeks.

Approval duration: 6 months

C. Other diagnoses/indications: Refer to PA.CP.PMN.53

II. Continued Approval

- A. Acromegaly (must meet all):
 - 1. Currently receiving medication via PA Health & Wellness benefit or member has previously met all initial approval criteria; or the Continuity of Care policy (PA.LTSS.PHAR.01) applies;
 - 2. Request is for Signifor LAR;
 - 3. Member is responding positively to therapy (*see Appendix D*);
 - 4. If request is for a dose increase, new dose does not exceed (a and b):
 - a. 60 mg every 4 weeks;
 - b. 1 vial every 4 weeks.

Approval duration: 12 months

B. Cushing's Disease (must meet all):

- 1. Currently receiving medication via PA Health & Wellness benefit or member has previously met all initial approval criteria; or the Continuity of Care policy (PA.LTSS.PHAR.01) applies;
- 2. Member is responding positively to therapy (see Appendix D);
- 3. If request is for a dose increase, new dose does not exceed one of the following (a or b):
 - a. Signifor (i and ii):
 - i. 1.8 mg per day;
 - ii. 2 ampules per day;
 - b. Signifor LAR (i and ii):
 - i. 40 mg every 4 weeks;
 - ii. 1 vial every 4 weeks.

Approval duration: 12 months

C. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via PA Health & Wellness benefit or member has previously met all initial approval criteria or the Continuity of Care Policy (PA.LTSS.PHAR.01) applies.

Approval duration: Duration of request or 6 months (whichever is less); or

- 2. Refer to the off-label use policy if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): PA.CP.PMN.53
 - a. The requested dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use.



III. Diagnoses/Indications for which coverage is NOT authorized:

A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policy – PA.CP.PMN.53 or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key FDA: Food and Drug Administration GH: growth hormone IGF-I: insulin-like growth factor

Appendix B: Therapeutic Alternatives Not applicable

Appendix C: Contraindications/Boxed Warnings None reported

Appendix D: General Information

- Treatment response for Cushing's disease may be defined as reduction in 24-hour urinary free cortisol (UFC) levels and/or improvement in signs or symptoms of the disease. Maximum urinary free cortisol reduction is typically seen by two months of treatment.
- Examples of treatment response to acromegaly therapy (including somatostatin analogs, surgical resection or pituitary irradiation) include improvement from baseline in or normalization of growth hormone (GH) and/or age- and sex-adjusted insulin-like growth factor (IGF-1) serum concentrations, or tumor mass control.

Dosage and Administration			
Drug Name	Indication	Dosing Regimen	Maximum Dose
Pasireotide (Signifor)	Cushing's disease	Initial: 0.6 mg or 0.9 mg SC BID	1.8 mg/day
		Recommended dosing range: 0.3 mg to 0.9 mg SC BID	
Pasireotide (Signifor LAR) [*]	Cushing's disease	10 mg to 40 mg IM every 4 weeks	40 mg/4 weeks
	Acromegaly	40 mg to 60 mg IM every 4 weeks	60 mg/4 weeks

V. Dosage and Administration

*Signifor LAR must be administered by a healthcare professional

IV. Product Availability

Drug Name	Availability
Pasireotide (Signifor)	Single-dose ampules for injection: 0.3 mg/mL, 0.6 mg/mL,
	0.9 mg/mL

Drug Name	Availability
Pasireotide (Signifor LAR)	Vial for reconstitution and injectable suspension: 10 mg, 20
	mg, 30 mg, 40 mg, 60 mg

VI. References

- 1. Signifor Prescribing Information. East Hanover, NJ: Novartis Pharmaceuticals Corporation; March 2020. Available at: <u>https://www.signifor.com/pdf/signifor-pi.pdf</u>. Accessed July 20, 2022.
- 2. Signifor LAR Prescribing Information. East Hanover, NJ: Novartis Pharmaceuticals Corporation; June 2020. Available at https://www.signiforlar.com/pdf/signifor-lar-pi.pdf. Accessed July 20, 2022.
- 3. Melmed S, Bronstein MD, Chanson P. A Consensus Statement on acromegaly therapeutic outcomes. Nat Rev Endocrinol. 2018 Sep;14(9):552-561.
- 4. Katznelson L, Laws ER, Melmed S, et al. Acromegaly: An Endocrine Society clinical practice guideline. J Clin Endocrinol Metab. 2014; 99(11): 3933-3951.
- 5. Fleseriu M, Biller BMK, Freda PU, et al. A Pituitary Society update to acromegaly management guidelines. Pituitary. 2021; 24: 1-13.
- 6. Guistina A, Barkhoudarian G, Beckers A, et al. Multidisciplinary management of acromegaly: A consensus. Rev Endocr Metab Disord. 2020; 21(4): 667-678.
- Nieman LK, Biller BMK, Findling JW, et al. Treatment of Cushing's syndrome: An Endocrine Society clinical practice guideline. J Clin Endocrinol Metab. 2015; 100(8): 2807-2831.
- 8. Fleseriu M, Auchus R, Bancos I, et al. Consensus on diagnosis and management of Cushing's disease: a guideline update. Lancet Diabetes Endocrinol. 2021; 9(12): 847-875.

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-todate sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J2502	Injection, pasireotide long acting, 1 mg

Reviews, Revisions, and Approvals	Date	Approval Date
4Q 2018 annual review: Signifor added to policy; criteria added for new FDA indication for Signifor LAR: Cushing's disease; new strengths of Signifor LAR added; requirement for inadequate response to surgery or pituitary irradiation added for acromegaly; references reviewed and updated.	08/2018	
4Q 2019 annual review: No changes per Statewide PDL implementation 01-01-2020	10/2019	

Reviews, Revisions, and Approvals	Date	Approval Date
4Q 2020 annual review: increased acromegaly initial approval duration from 3 months to 6 months to align with approach for other acromegaly policies; references reviewed and updated.	08/2020	
4Q 2021 annual review: no significant changes; updated J code; references reviewed and updated	10/2021	
4Q 2022 annual review: for acromegaly, added confirmatory diagnostic requirements (IGF-I or GH) per PS/ES practice guidelines; references reviewed and updated.	10/2022	