

## Clinical Policy: Pegvaliase-pqpz (Palynziq)

Reference Number: PA.CP.PHAR.140

Effective Date: 10.17.18

Last Review Date: 10/2023

[Revision Log](#)

### Description

Pegvaliase-pqpz (Palynziq™) is a PEGylated phenylalanine ammonia lyase (PAL) enzyme that converts phenylalanine to ammonia and trans-cinnamic acid. It substitutes for the deficient phenylalanine hydroxylase (PAH) enzyme activity in patients with phenylketonuria (PKU) and reduces blood phenylalanine concentrations.

### FDA Approved Indication(s)

Palynziq is indicated to reduce blood phenylalanine concentrations in adult patients with PKU who have uncontrolled blood phenylalanine concentrations > 600 µmol/L on existing management.

### Policy/Criteria

*Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.*

It is the policy of PA Health & Wellness® that Palynziq is **medically necessary** when the following criteria are met:

### I. Initial Approval Criteria

#### A. Phenylketonuria (must meet all):

1. Diagnosis of PKU;
2. Prescribed by or in consultation with an endocrinologist, metabolic disease specialist, or genetic disease specialist;
3. Age ≥ 18 years;
4. Recent (within 90 days) phenylalanine (Phe) blood level is > 600 µmols/L;
5. Member is currently on a phenylalanine-restricted diet and will continue this diet during treatment with Palynziq;
6. Failure of Kuvan® at up to maximally indicated doses, unless contraindicated or clinically significant adverse effects are experienced;
7. Palynziq is not prescribed concurrently with Kuvan;
8. Dose does not exceed 20 mg per day.

**Approval duration: 12 months**

#### B. Other diagnoses/indications

1. Refer to the off-label use policy if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): PA.CP.PMN.53.

### II. Continued Therapy

#### A. Phenylketonuria (must meet all):

1. Currently receiving medication via PA Health & Wellness benefit or member has previously met all initial approval criteria or the Continuity of Care Policy (PA.LTSS.PHAR.01) applies;
2. Member is currently on a phenylalanine-restricted diet and will continue this diet during treatment with Palynziq;
3. Member meets one of the following (a, b, or c):
  - a. Member has achieved blood Phe control (i.e., blood Phe level is  $\leq 600$   $\mu\text{mol/L}$ );
  - b. Request is for 40 mg per day and member has previously used 20 mg per day continuously for at least 6 months without achieving blood Phe control;
  - c. Request is for 60 mg per day and member meets both of the following (i and ii):
    - i. Member has previously used 40 mg per day continuously for at least 16 weeks without achieving blood Phe control;
    - ii. Member has not used 60 mg per day continuously for more than 16 weeks without achieving blood Phe control;
4. If request is for a dose increase, new dose does not exceed 60 mg per day.

**Approval duration: 12 months**

**B. Other diagnoses/indications (must meet 1 or 2):**

1. Currently receiving medication via PA Health & Wellness benefit or member has previously met all initial approval criteria or the Continuity of Care Policy (PA.LTSS.PHAR.01) applies.  
**Approval duration: Duration of request or 6 months (whichever is less);** or
2. Refer to the off-label use policy if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): PA.CP.PMN.53.

**III. Diagnoses/Indications for which coverage is NOT authorized:**

- A.** Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policy – PA.CP.PMN.53 or evidence of coverage documents.

**IV. Appendices/General Information**

*Appendix A: Abbreviation/Acronym Key*

FDA: Food and Drug Administration

PAH: phenylalanine hydroxylase

PAL: phenylalanine ammonia lyase

Phe: phenylalanine

PKU: phenylketonuria

*Appendix B: Therapeutic Alternatives*

*This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.*

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
Kuvan (sapropterin)	Age 1 month to $\leq 6$ years (starting dose): 10 mg/kg PO QD Age $\geq 7$ years (starting dose): 10 to 20 mg/kg PO QD	20 mg/kg/day

Therapeutic alternatives are listed as Brand name<sup>®</sup> (generic) when the drug is available by brand name only and generic (Brand name<sup>®</sup>) when the drug is available by both brand and generic.

*Appendix C: Contraindications/Boxed Warnings*

- Contraindication(s): none reported
- Boxed warning(s): risk of anaphylaxis

*Appendix D: General Information*

- Palynziq has a black box warning for the potential to cause anaphylaxis and enrollment in a REMS program is required, along with supervision of the initial dose by a healthcare professional and the need to carry auto-injectable epinephrine at all times while using Palynziq. Use of premedication with H<sub>1</sub> blockers, H<sub>2</sub> blockers, and/or antipyretics can also be considered.
- Per the Palynziq PI, discontinuation of Palynziq is recommended if a patient has not achieved an adequate response (blood Phe concentration  $\leq 600 \mu\text{mol/L}$ ) after 16 weeks of continuous treatment with the maximum dosage of 60 mg QD.

**V. Dosage and Administration**

Indication	Dosing Regimen	Maximum Dose
PKU	<p>Initiate dosing with 2.5 mg SC once weekly for 4 weeks. Administer the initial dose under the supervision of a healthcare provider.</p> <p>Titrate the Palynziq dosage in a step-wise manner, based on tolerability, over <math>\geq 5</math> weeks, to achieve a dosage of 20 mg SC QD.</p> <p>Maintain the Palynziq dosage at 20 mg SC QD for <math>\geq 24</math> weeks. Consider increasing the Palynziq dosage to 40 mg SC QD in patients who have been maintained continuously on 20 mg QD for <math>\geq 24</math> weeks and who have not achieved a blood Phe concentration <math>\leq 600 \mu\text{mol/L}</math>.</p> <p>Consider increasing the dosage to a maximum of 60 mg SC QD in patients who have been on 40 mg QD continuously for <math>\geq 16</math> weeks and who have not achieved a blood Phe concentration <math>\leq 600 \mu\text{mol/L}</math>.</p> <p>Discontinue Palynziq in patients who have not achieved a response (blood Phe concentration <math>\leq 600 \mu\text{mol/L}</math>) after 16 weeks of continuous treatment with the maximum dosage of 60 mg QD.</p>	60 mg/day

**VI. Product Availability**

Injection, single-dose prefilled syringe: 2.5 mg/0.5 mL, 10 mg/0.5 mL, 20 mg/mL

**VII. References**

1. Palynziq Prescribing Information. Novato, CA: BioMarin Pharmaceutical Inc.; November 2020. Available at: <http://www.palynziq.com>. Accessed June 29, 2023.
2. Vockley J, Andersson HC, et al. Phenylalanine hydroxylase deficiency: diagnosis and management guideline. Genet Med. Feb 2014;16(2):188-200.
3. Thomas J, Levy H, et al. Pegvaliase for the treatment of phenylketonuria: results of a long-term phase 3 clinical trial program (PRISM). Molecular Genetics and Metabolism. 2018;124:27-38.
4. Harding CO, Amato RS, et al. Pegvaliase for the treatment of phenylketonuria: a pivotal, double-blind randomized discontinuation phase 3 clinical trial. Molecular Genetics and Metabolism. 2018;124:20-26.

**Coding Implications**

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J3590	Unclassified biologics

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created	10/2018	
4Q 2019 annual review: No changes per Statewide PDL implementation 01-01-2020	10/2019	
4Q 2020 annual review: added age limit; added requirement for current and continued use of Phe-restricted diet; added requirement for a prior trial of Kuvan; referenced reviewed and updated.	10/2020	
4Q 2021 annual review: RT4: revised continuation criteria to reflect updated dosing recommendations in the package labeling; references reviewed and updated.	10/2021	
4Q 2022 annual review: no significant changes; references reviewed and updated.	10/2022	
4Q 2023 annual review: no significant changes; references reviewed and updated.	10/2023	