

Clinical Policy: IV Moderate Sedation, IV Deep Sedation, and General Anesthesia for Dental Procedures

Reference Number: PA.CP.MP.61

Effective Date: 01/2018

Date of Last Revision: 10/2023

[Coding Implications](#)

[Revision Log](#)

Description

The administration of intravenous moderate sedation and intravenous deep sedation/general anesthesia in the dental office, inpatient hospital, or ambulatory surgical center may be necessary to safely provide dental care. These procedures generally are safe when administered by trained, certified providers in the appropriate setting but are not without risk. According to the American Academy of Pediatrics and the American Academy of Pediatric Dentistry (AAPD), the sedation of children is different from the sedation of adults, and the in-office use of deep sedation or general anesthesia may be appropriate on select pediatric dental patients administered in appropriately equipped and staffed facilities. Pediatric patients are subject to higher risk of adverse outcomes with sedation. This policy governs only the services cited. For guidance in prior authorization processes refer to PA Health & Wellness Provider Manual and Pennsylvania PROMISe™ 837 Dental/ADA form.

Policy/Criteria

- I. It is the policy of PA Health and Wellness® (PHW) that intravenous moderate sedation, intravenous deep sedation, and general anesthesia are **medically necessary** when all of the following general criteria and place of service-specific criteria are met in either II or III:
 - A. One or more of the following indications:
 1. Documented extreme anxiety or fear and documentation of failed local anesthesia, inhalation of nitrous oxide, or oral conscious sedation attempts;
 2. Dental and medical records documentation substantiates a physical, cognitive, or developmental disability (including, but not limited to, congenital cardiac defect, cerebral palsy, epilepsy) that would render the patient non-compliant;
 3. Underlying medical conditions are supported by patient records;
 4. There is documentation of allergy or sensitivity to local anesthesia;
 5. Lengthy and numerous (more than four) restorative procedures for pediatric member/enrollee;
 6. Documentation indicates a pediatric member/enrollee has resisted all other conventional behavior management procedures;
 7. Documentation indicates a pediatric member/enrollee has history of uncooperative behaviors that may result in the compromised safety of the member/enrollee, provider, or staff;
 8. Documented history of relevant social impact conditions;
 9. Complex and extensive surgical procedures such as two or more impacted teeth in two or more quadrants, or two or more extractions of permanent teeth per quadrant in at least two quadrants.
 - B. None of the following:
 1. Member/enrollee has predisposing medical and/or physical conditions that would make general anesthesia unsafe;

IV Moderate Sedation, IV Deep Sedation and General Anesthesia for Dental Procedures

2. Member/enrollee is cooperative and has minimal dental needs;
 3. General anesthesia or moderate or deep sedation is requested for member/enrollee or dentist convenience;
 4. Member/enrollee has history of tolerating similar dental procedures without the need for sedation;
 5. Member/enrollee has a choice of an alternative option for treatment;
 6. Member/enrollee has an allergy to oral sedatives or reversal agents;
 7. Parent or guardian objects to the sedation or general anesthesia;
 8. There are no covered and approved dental services to be rendered.
- C. Required documentation, all the following:
1. Current (less than six months) diagnostic quality pre-operative radiographs (when achievable) or intraoral photographic images if radiographs are not achievable or if requested treatment is not visualized in the radiographs;
 2. Patient chart and treatment notes document conditions supporting medical necessity and all relevant dental and medical diagnoses and/or ICD-10 diagnosis codes;
 3. Patient chart and treatment notes document applicable non-clinical and/or relevant social conditions;
 4. Complete treatment plan;
 5. ASA physical status classification (ASA Classes III and IV for member/enrollees under the age of eight will not be approved for a dental office setting);
 6. Specific history of failed anesthesia, sedation, or analgesia/anxiolysis attempts;
 7. Referring dentist evaluation records;
 8. State-mandated outpatient (OP) form, if applicable.
- II. It is the policy of PHW[®] that in-office IV sedation or general anesthesia is **medically necessary** and the appropriate venue when the general criteria in section I are met, the member/enrollee has no significant medical comorbidities and any of the following:
- A. Member/enrollee exhibits moderate or situational anxiety;
 - B. Pediatric member/enrollee with limited treatment need (four to six teeth requiring restoration, pulpotomy, or extraction);
 - C. Pediatric member/enrollee has social impact conditions necessitating completion of all treatment needs during a single visit;
 - D. Pediatric member/enrollee under the age of eight with medically necessary dental services and with an ASA I or II classification;
 - E. Pediatric member/enrollee under the age of eight with medically necessary dental services and with an uncompromised airway;
 - F. Member/enrollee requires extraction of two or more impacted teeth in two or more quadrants;
 - G. Member/enrollee requires extraction of two or more permanent teeth per quadrant involving two or more quadrants.
- III. It is the policy of PHW[®] that ambulatory surgical center (ASC) or hospital out-patient IV sedation or general anesthesia is **medically necessary** and the appropriate venue when the general criteria in section I. are met and any of the following:

IV Moderate Sedation, IV Deep Sedation and General Anesthesia for Dental Procedures

- A. Member/enrollee has compromising medical comorbidities;
- B. Local factors (e.g., access to care) or state regulations permit treating cases in an ASC that meet in-office approval conditions;
- C. Pediatric member/enrollee with extensive treatment needs (seven or more teeth requiring restoration, pulpotomy, or extraction of primary teeth);
- D. Pediatric member/enrollee with a combination of at least four teeth requiring restoration, pulpotomy, or extraction of primary teeth and the presence of at least one major medical comorbidity (e.g., documented anxiety disorder, uncontrolled asthma, pre-combative or combative behavior);
- E. Pediatric member/enrollee with social impact conditions necessitating completion of all treatment needs during a single visit and in-office IV sedation or general anesthesia is not accessible;
- F. Pediatric member/enrollee with social impact conditions necessitating completion of all treatment needs during a single visit in combination with a major medical comorbidity;
- G. Pediatric member/enrollee under the age of eight years with medically necessary dental services meeting clinical criteria and with an ASA III or IV classification;
- H. Pediatric member/enrollee under the age of eight years with medically necessary dental services and a compromised or obstructed airway;
- I. Member/enrollee requires extraction of two or more impacted teeth in two or more quadrants in combination with a major medical comorbidity;
- J. Member/enrollee requires extraction of two or more permanent teeth per quadrant involving two or more quadrants in combination with a major medical comorbidity.

Background*Guidelines*

According to the American Dental Association (ADA), dentists must comply with their state laws, rules and/or regulations when providing sedation and anesthesia and follow the educational and training requirements for the level of sedation intended. The ADA maintains clinical guidelines and educational/training requirements for all levels of sedation and includes specific information for the following:

- Patient history and evaluation
- Pre-operative preparation
- Personnel and equipment requirements
- Monitoring and documentation (including consciousness, oxygenation, ventilation, and circulation)
- Recovery and discharge
- Emergency management

Providers are encouraged to embrace and utilize the following American Academy of Pediatrics (AAP), American Academy of Pediatric Dentistry (AAPD), American Society of Anesthesiologists (ASA), Society for Pediatric Anesthesia, American Society of Dentist Anesthesiologists, and Society for Pediatric Sedation guidelines and/or recommendations⁵.

- High-risk patients and pediatric patients under the age of eight should be evaluated by a primary care physician or physician anesthesiologist prior to scheduling a procedure.

IV Moderate Sedation, IV Deep Sedation and General Anesthesia for Dental Procedures

- Prolonged and extensive procedures with longer periods of sedation and anesthesia care are of concern in the office-based setting and qualified anesthesia providers, in consultation with such patients, should consider more suitable facilities for the procedure.
- *The sedation must be administered by a qualified anesthesia provider (a medical anesthesiologist, certified registered nurse anesthetist, dentist anesthesiologist or second oral surgeon). At a minimum, the sedation provider must be "able to provide advanced pediatric life support (PALS) and capable of rescuing a child with apnea, laryngospasm, and airway obstruction. Required skills include the ability to open the airway, suction secretions, provide CPAP, insert supraglottic devices (oral airway, nasal trumpet, laryngeal mask airway) and perform successful bag-valve-mask ventilation, tracheal intubation, and cardiopulmonary resuscitation.*
- A second observer, who is also skilled, and PALS certified, must also be present to assist with the anesthetic emergency while additional support is summoned; this is particularly critical in a dental office-based setting as the only backup is to call 911.
- PALS-certified observers should be present to monitor the patient throughout procedures performed with sedation and have no other responsibilities during deep sedation and/or general anesthesia.
- Oral surgeons and other dental practitioners who provide intravenous deep sedation or general anesthesia should discontinue the use of the single provider/operator model for the care of pediatric patients under the age of eight when sedated in dental offices.

Drugs, Medicaments, and Gases

Drugs, medicaments, and gases are used for the following levels of anxiolysis/analgesia, oral conscious sedation, IV moderate sedation, IV deep sedation/general anesthesia and include, but are not limited to the following:

Anxiolysis/Analgesia

- Nitrous oxide and oxygen

Oral Conscious Sedation

- Midazolam – Schedule IV
- Hydroxyzine
- Diphenhydramine

IV Moderate Sedation

- Benzodiazapine
- Opioids – Schedule II

IV Deep Sedation/General Anesthesia

- Propofol
- Benzodiazapine
- Ketamine – Schedule III
- Dexmedetomidine
- Sevoflurane
- Desflurane
- Isoflurane
- Methohexital – Schedule IV
- Succinylcholine
- Opioids – Schedule II

Sedation/General Anesthesia Definitions

IV Moderate Sedation: A drug-induced depression of consciousness during which patients respond purposefully to verbal commands, either alone or accompanied by light tactile stimulation. No interventions are required to maintain a patent airway, and spontaneous ventilation is adequate. Cardiovascular function is usually maintained.

IV Moderate Sedation, IV Deep Sedation and General Anesthesia for Dental Procedures

IV Deep Sedation: A drug-induced depression of consciousness during which patients cannot be easily aroused but respond purposefully after repeated verbal or painful stimulation (e.g., purposefully pushing away the noxious stimuli). Reflex withdrawal from a painful stimulus is not considered a purposeful response and is more consistent with a state of general anesthesia. The ability to maintain ventilator function may be impaired. Patients may require assistance in maintaining a patent airway, and spontaneous ventilation may be inadequate. Cardiovascular function is usually maintained.

General Anesthesia: A drug-induced loss of consciousness during which patients are not arousable, even by painful stimulation. The ability to independently maintain respiratory function is often impaired. Patients often require assistance in maintaining a patent airway, and positive-pressure ventilation may be required because of depressed spontaneous ventilation or drug-induced depression of neuromuscular function. Cardiovascular function may be impaired.

Coding Implications

This clinical policy references Current Procedural Terminology (CPT®). CPT® is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2022, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

This clinical policy references Current Dental Terminology (CDT®). CDT® is a registered trademark of the American Dental Association. All CDT codes and descriptions are copyrighted 2022, American Dental Association. All rights reserved. CDT codes and CDT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

Coding Limitations/Exclusions

1. One D9222 or D9239 per date of service.
2. More than four units of D9223 or D9243 per date of service requires a copy of the sedation log, subject to state-specific regulations.

CPT® Codes	Description
00170	Anesthesia for intraoral procedures, including biopsy; not otherwise specified
00190	Anesthesia for procedures on facial bones or skull; not otherwise specified

--	--

IV Moderate Sedation, IV Deep Sedation and General Anesthesia for Dental Procedures

HCPCS Codes	Description		
D9222	Deep sedation/general anesthesia – first 15 minutes		
D9223	Deep sedation/general anesthesia – each subsequent 15-minute increment		
D9239	Intravenous moderate (conscious) sedation/analgesia – first 15 minutes		
D9243	Intravenous moderate (conscious) sedation/analgesia – each subsequent 15 minute increment		
D9248	Non-intravenous conscious sedation		
Reviews, Revisions, and Approvals		Review Date	Approval Date
References reviewed and updated. Code updates		05/18	07/18
References reviewed and updated. Specialist reviewed.		10/19	2/18/2020
Annual Review performed, References reviewed and updated. Specialist reviewed.		2/26/2021	
Update reference PA Health & Wellness Provider Manual and reference to April 15, 2021 Medical Assistance Bulletin- Dental Benefit Limit Exception Process Update- Provider Handbook, 837 Dental/ADA 2012 Claim form. “This policy governs only the services cited. For guidance in prior authorization processes refer to PA Health & Wellness Provider Manual and Pennsylvania PROMISE™ 837 Dental/ADA form.”		07/2021	
Annual review. References reviewed, updated, and reformatted. Changed “review date” in the header to “date of last revision” and “date” in the revision log header to “revision date.”		6/24/2022	
Annual Review. Changed title from “Dental Anesthesia” to “IV Moderate Sedation, IV Deep Sedation, and General Anesthesia for Dental Procedures” and adopted the Envolve Dental Policy criteria from ENVD.UM.CP.0009, approved 11/21. Removed HCPCS code D9230. Minor rewording without clinical significance. Specified that general criteria in I. must be met in addition to POS-specific criteria in II or III. Clarified in I. that A (indications), B (lack of contraindications), and C (documentation) must all be met. Specified in II that absence of comorbidities applies to all indications. Moved professional society guidelines/recommendations, list of sedation/anesthesia drugs, and definitions to the background. In I.A. replaced “any” with “one or more” with no impact to criteria. Combined I.C.1. and I.C.2. Background: added “pre-operative preparation” and restructured wording with no impact to criteria. Added CDT disclaimer under coding implications. References reviewed, updated, and reformatted. Reviewed by internal specialist.		10/2023	

References

1. Andropoulos DB, Greene MF. Anesthesia and Developing Brains - Implications of the FDA Warning. *N Engl J Med.* 2017;376(10):905 to 907. doi:10.1056/NEJMp1700196

IV Moderate Sedation, IV Deep Sedation and General Anesthesia for Dental Procedures

2. American Academy on Pediatric Dentistry Ad Hoc Committee on Sedation and Anesthesia; American Academy on Pediatric Dentistry Council on Clinical Affairs. Policy on the use of deep sedation and general anesthesia in the pediatric dental office. *Pediatr Dent*. 2008;30(7 Suppl):66 to 67.
3. Coté CJ, Wilson S; AMERICAN ACADEMY OF PEDIATRICS; AMERICAN ACADEMY OF PEDIATRIC DENTISTRY. Guidelines for Monitoring and Management of Pediatric Patients Before, During, and After Sedation for Diagnostic and Therapeutic Procedures. *Pediatrics*. 2019;143(6):e20191000. doi:10.1542/peds.2019-1000
4. American Dental Association. *CDT2020 Current Dental Procedures*. 2023
5. American Dental Association. Guidelines for Use of Sedation and General Anesthesia by Dentists (2007). https://www.ada.org/-/media/project/ada-organization/ada/ada-org/files/publications/cdt/anesthesia_guidelines.pdf. Accessed August 30, 2023.
6. American Society of Anesthesiologists. Statement on Sedation & Anesthesia Administration in Dental Office-Based Settings. <https://www.asahq.org/standards-and-practice-parameters/statement-on-sedation--anesthesia-administration-in-dental-officebased-settings>. Accessed August 30, 2023.
7. Berkowitz R, Barinholtz D. *Anesthesia for Office-Based Pediatric Anesthesia. Smith's Anesthesia for Infants and Children*. 8th Edition. Elsevier Inc.; 2011
8. Collado V, Faulks D, Nicolas E, Hennequin M. Conscious sedation procedures using intravenous midazolam for dental care in patients with different cognitive profiles: a prospective study of effectiveness and safety. *PLoS One*. 2013;8(8):e71240. Published 2013 Aug 5. doi:10.1371/journal.pone.0071240
9. de Nova García MJ, Gallardo López NE, Martín Sanjuán C, Mourelle Martínez MR, Alonso García Y, Carracedo Cabaleiro E. Criteria for selecting children with special needs for dental treatment under general anaesthesia. *Med Oral Patol Oral Cir Bucal*. 2007;12(7):E496 to E503. Published 2007 Nov 1.
10. El Batawi HY. Effect of preoperative oral midazolam sedation on separation anxiety and emergence delirium among children undergoing dental treatment under general anesthesia. *J Int Soc Prev Community Dent*. 2015;5(2):88 to 94. doi:10.4103/2231-0762.155728
11. Heaton LJ, McNeil DW, Milgrom P. Propranolol and D-cycloserine as adjunctive medications in reducing dental fear in sedation practice. *SAAD Dig*. 2010;26:27 to 35.
12. Lyratzopoulos G, Blain KM. Inhalation sedation with nitrous oxide as an alternative to dental general anaesthesia for children. *J Public Health Med*. 2003;25(4):303 to 312. doi:10.1093/pubmed/fdg068
13. Malamed, SF. *Sedation: A Guide to Patient Management*. 5th Edition. Elsevier - Health Sciences Division; 2009.
14. Nelson T, Nelson G. The role of sedation in contemporary pediatric dentistry. *Dent Clin North Am*. 2013;57(1):145 to 161. doi:10.1016/j.cden.2012.09.007
15. Nelson TM, Xu Z. Pediatric dental sedation: challenges and opportunities. *Clin Cosmet Investig Dent*. 2015;7:97 to 106. Published 2015 Aug 26. doi:10.2147/CCIDE.S64250
16. Southerland JH, Brown LR. Conscious Intravenous Sedation in Dentistry: A Review of Current Therapy. *Dent Clin North Am*. 2016;60(2):309 to 346. doi:10.1016/j.cden.2015.11.009