

Clinical Policy: Vagus Nerve Stimulation

Reference Number: PA.CP.MP.12

Effective Date: 01/18

Last Review Date: 06/28/2021

[Coding Implications](#)

[Revision Log](#)

Description

Vagus nerve stimulation (VNS) has been used in the treatment of epilepsy and has been studied for the treatment of refractory depression and other indications. Electrical pulses are delivered to the cervical portion of the vagus nerve by an implantable device called a neurocybernetic prosthesis. Chronic intermittent electrical stimulation of the left vagus nerve is designed to treat medically refractory epilepsy. It has recently been introduced and approved by the FDA as an adjunctive therapy for treatment-resistant major depression.

Policy/Criteria

- I. It is the policy of Pennsylvania Health and Wellness[®] (PHW), that VNS is **medically necessary** in patients with medically refractory seizures who meet all of the following:
 - A. Diagnosis of local onset (formerly partial onset) seizures or generalized onset seizures;
 - B. Intractable epilepsy (both):
 1. Failure of at least 1 year of adherent therapy of at least two anti-seizure drugs, *and*
 2. Continued seizures which have a major impact on activities of daily living; *and*
 - C. Not a suitable candidate for or has failed resective epilepsy surgery;
 - D. Request is for an FDA-approved device.
- II. It is the policy of PHW that the safety and efficacy of VNS therapy has not been proven for any other conditions, including but not limited to the following:
 - A. Refractory (treatment resistant) major depression or bipolar disorder;
 - B. Obesity;
 - C. Headaches;
 - D. Cognitive impairment associated with Alzheimer's disease;
 - E. Addiction;
 - F. Anxiety Disorders;
 - G. Autism;
 - H. Eating Disorders;
 - I. Cancer;
 - J. Crohn's Disease;
 - K. Essential trauma;
 - L. Fibromyalgia;
 - M. Heart failure;
 - N. Impaired glucose tolerance/pre-diabetes;
 - O. Inflammation;
 - P. Overweight and obesity;
 - Q. Obsessive-compulsive disorder;
 - R. Panic disorder;
 - S. Post-traumatic stress disorder;
 - T. Prader-Willi Syndrome;
 - U. Sjogren's Syndrome;

CLINICAL POLICY

Vagus Nerve Stimulation

- V. Rheumatoid arthritis;
- W. Schizophrenia;
- X. Sleep disorders;
- Y. Stroke;
- Z. Tinnitus;
- AA. Tourette's syndrome;
- BB. Traumatic brain injury

III. It is the policy of PHW that the current research does not support the use of following types of VNS therapy over other currently available alternatives, due to the lack of large, high-quality studies supporting their use:

- A. Aspire SR Model 106 (Cyberonics) for vagus nerve stimulation;
- B. Transcutaneous VNS or active auricular transcutaneous electrical nerve stimulation.

Removal of Implant

Less than 0.5 percent of all patients have had the device removed. It can be turned off in the physician's office if the patient feels it is not helping or if the patient cannot tolerate the stimulation. If the device needs to be removed, only the pulse generator is removed, as attempting to remove the electrodes from around the nerve can cause damage and is not recommended.

Background

The vagus nerve stimulator is a pacemaker-like device implanted under the skin in the left side of the chest through a small incision, with a second small incision made at the base of the neck. The surgery is performed under local, regional, or general anesthesia and lasts 45 minutes to two hours. Most often it is performed as an outpatient surgery but some patients need to stay in the hospital overnight following surgery.

Partial (focal) seizures

Several studies have been done evaluating the safety and effectiveness of vagus nerve stimulation for treatment of epilepsy. A randomized active-control trial known as the E05 study found that 94 patients (of the total 254 patients in the study) receiving high stimulation showed an average reduction in seizure frequency, compared to baseline, of 28% versus 15% reduction in the 102 patients receiving low stimulation. A total of 310 patients completed the E03 and E05 double-blinded trials. Mean decline of seizure frequency overall was about 25-30% compared to baseline. Clinical experience has shown that improvement in seizures is maintained, or may even increase over time, but these data are based on uncontrolled observations. Side effects in both studies were similar and included hoarseness and occasional shortness of breath.

Although questions regarding patient selection criteria, optimal stimulation parameters, and cost-effectiveness in the United States remain under investigation, there is sufficient evidence regarding the benefit and safety of VNS to conclude that VNS may improve health outcomes in patients with medically refractory partial-onset seizures who are not suitable candidates for surgery or in whom surgical treatment has failed.

Generalized seizures

CLINICAL POLICY

Vagus Nerve Stimulation

Study results suggest VNS may be effective for generalized epilepsy (Karczeski & Schachter, 2015). However, case series and observational studies constitute the majority of available evidence. Although VNS is not currently FDA approved for the treatment of generalized seizures, it is often used in children and other patients, and in Europe is approved as adjunct therapy for epileptic disorders predominantly characterized by generalized or focal seizures that are refractory to antiseizure medications. In addition, the National Institute for Health and Care Excellence (NICE) recommends VNS for focal and generalized seizures, and the Scottish Intercollegiate Guidelines Network (SIGN) guidelines recommend VNS for epilepsy in patients unsuitable for respective surgery without stipulating seizure type.

Depression

VNS was FDA-approved for treatment resistant depression in 2005. However, VNS has no rigorous research data proving it is efficacious for treatment-resistant, unipolar major depression. Open-label studies suggest VNS may be effective; however, these are at risk for bias due to placebo effects. The one randomized trial of VNS for depression found no benefit, with outcomes comparable for active and sham treatment (response rates of 15 vs. 10 percent). In addition, there is a lack of thorough safety data for the use of VNS in depression.

Other Investigational Indications

Ongoing research efforts continue to investigate the role of vagus nerve stimulation (VNS) for the treatment of a variety of indications, including but not limited to cognitive deficits in Alzheimer's disease, resistant obesity, and headaches. Data supporting the long-term safety and efficacy from large clinical trials of VNS for the treatment of these indications, however, continue to be lacking.

AspireSR Model 106 (Cyberonics) for Vagus Nerve Stimulation

The AspireSR Model 106 (Cyberonics Inc.) received FDA Premarket Approval (PMA) in February 2014. The newest modification to the vagus nerve stimulation (VNS) implant detects tachycardia heart rates, which may be associated with an impending seizure, and automatically delivers stimulation to the vagus nerve. Like its predecessors, the AspireSR can also deliver stimulation in the normal and magnet modes. However, when programmed for AutoStim mode, the AspireSR requires no patient interaction to trigger the delivery of electrical stimulation. The AutoStim mode should not be used in patients with significant arrhythmias being treated with pacemakers and/or an implantable defibrillator, beta-blockers, or any other treatment that may impact the intrinsic heart rate.

A few small, preliminary studies and case reports have evaluated the AspireSR Model 106, and have shown positive results (Boon et al., 2015; Fisher et al., 2015; Schneider et al., 2015; Hampel et al, 2015). However, there is insufficient evidence to establish the safety and efficacy of the AspireSR Model 106 in reducing seizures until further, high quality trials establish its clinical value.

Transcutaneous Vagus Nerve Stimulation

Transcutaneous vagus nerve stimulation (tVNS) has been proposed as a noninvasive alternative to implantable VNS for a variety of indications, including, but not limited to epilepsy, major depression, chronic tinnitus and headaches. Currently, there are two main ways to apply tVNS.

One is to apply stimulation on the ear and the other is cervical noninvasive VNS, superficially applying stimulation in the vicinity of the vagus nerve using a specially designed device, (e.g. gammaCore). Noninvasive auricular tVNS stimulates the afferent auricular branch of the vagus nerve located medial of the tragus at the entry of the acoustic meatus. Given that the right vagal nerve has efferent fibers to the heart, tVNS is safe to be performed only in the left ear. tVNS has been proposed to study cognitive functioning in patients with epilepsy and major depression. The rationale is that direct stimulation of the afferent nerve fibers on the ear area with afferent vagus nerve distribution should produce a similar effect as classic VNS in reducing depressive symptoms without the burden of surgical intervention. A noninvasive, transcutaneous vagal nerve stimulator has been in use in Europe. Although no randomized studies have been done in patients with epilepsy, it appears promising in one pilot study.¹⁸ Small studies have shown positive results with tVNS for the treatment of depression (Hein et al, 2013; Fang et al, 2016). Additional, larger, peer-reviewed studies, with longer follow-up are necessary to determine the long-term safety and efficacy of transcutaneous VNS for depression.

gammaCore Sapphire™ (ElectroCore, LLC), is a hand-held prescription device that is placed externally on the side of the neck in the vicinity of the vagus nerve to deliver a low voltage electric signal to the nerve's afferent fibers. gammaCore has received FDA approval for the treatment of both episodic cluster and migraine headaches and more recently for the prevention of cluster headaches (CH). gammaCore delivers up to 30 stimulations in a 24-hour period, each lasting 2 minutes. The patient controls the intensity level. Once the maximum daily number of treatments has been reached, the device will not deliver any more treatments until the following 24-hour period. gammaCore is rechargeable and includes a charging case to charge the device. A gammaCore refill card is used to load the device with days of therapy based on a healthcare provider's prescription.

In the randomized PRESTO study, noninvasive vagus nerve stimulation (nVNS.) was superior to sham in the treatment of episodic migraine for pain freedom at 30 minutes and 60 minutes after the first treated attack.²⁹ In both the ACT1 and ACT2 trials, nVNS was superior to sham therapy in episodic CH but not in chronic CH.^{31,32} Preliminary clinical trials of nVNS in various primary headache disorders are encouraging, however, well-designed randomized controlled trials with larger sample size and long-term follow-up regarding safety and benefit is warranted. In addition, patient selection criteria needs to be defined.

The American Headache Society position statement on integrating new migraine treatments into clinical practice note that empirically validated behavioral treatments with Grade A evidence for the prevention of migraine, including cognitive behavioral therapy, biofeedback, and relaxation therapies, should be considered in the management of migraine. These modalities may also be used alone or in addition to pharmacologic treatment. They note further that several noninvasive devices have been developed and approved by the FDA for the treatment of patients with migraine.(i.e., single-pulse transcranial magnetic stimulation, electrical trigeminal nerve stimulation and nVNS.) Patients who prefer nondrug therapies and those who have failed to respond to, have contraindications to, or poor tolerability with pharmacotherapy may be candidates for neuromodulation.³³

CLINICAL POLICY

Vagus Nerve Stimulation



Per UpToDate, “There are several promising but unproven methods using neurostimulation to treat medically refractory cluster headache, including sphenopalatine ganglion stimulation, occipital nerve stimulation, noninvasive VNS, and deep brain stimulation. All are investigational and require further study to confirm long-term benefit and safety.”

Coding Implications

This clinical policy references Current Procedural Terminology (CPT®). CPT® is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2021, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

CPT®* Codes	Description
61885	Insertion or replacement of cranial neurostimulator pulse generator or receiver, direct or inductive coupling; with connection to a single electrode array
61886	Insertion or replacement of cranial neurostimulator pulse generator or receiver, direct or inductive coupling; with connection to two or more electrode arrays
61888	Revision or removal of cranial neurostimulator pulse generator or receiver
64553	Percutaneous implantation of neurostimulator electrodes; cranial nerve
64568	Incision for implantation of cranial nerve (eg, vagus nerve) neurostimulator electrode array and pulse generator
64569	Revision or replacement of cranial nerve (eg, vagus nerve) neurostimulator electrode array, including connection to existing pulse generator
64570	Removal of cranial nerve (eg, vagus nerve) neurostimulator electrode array and pulse generator

HCPCS Codes that Support Coverage Criteria

HCPCS Codes	Description
C1767	Generator, neurostimulator (implantable), nonrechargeable
C1778	Lead, neurostimulator (implantable)
C1816	Receiver and/or transmitter, neurostimulator (implantable)
C1883	Adaptor/extension, pacing lead or neurostimulator lead (implantable)
L8680	Implantable neurostimulator electrode, each
L8681	Patient programmer (external) for use with implantable programmable neurostimulator pulse generator, replacement only
L8682	Implantable neurostimulator radiofrequency receiver
L8683	Radiofrequency transmitter (external) for use with implantable neurostimulator radiofrequency receiver
L8685	Implantable neurostimulator pulse generator, single array, rechargeable, includes extension

CLINICAL POLICY
Vagus Nerve Stimulation



HCPCS Codes	Description
L8686	Implantable neurostimulator pulse generator, single array, nonrechargeable, includes extension
L8687	Implantable neurostimulator pulse generator, dual array, rechargeable, includes extension
L8688	Implantable neurostimulator pulse generator, dual array, nonrechargeable, includes extension
L8689	External recharging system for battery (internal) for use with implanted neurostimulator, replacement only

HCPCS Codes that Do Not Support Coverage Criteria

HCPCS Codes	Description
K1020	Noninvasive vagus nerve stimulator

ICD-10-CM Diagnosis Codes that Support Coverage Criteria

ICD-10-CM Code	Description
G40.011	Localization-related (focal) (partial) idiopathic epilepsy and epileptic syndromes with seizures of localized onset, intractable, with status epilepticus
G40.019	Localization-related (focal) (partial) idiopathic epilepsy and epileptic syndromes with seizures of localized onset, intractable, without status epilepticus
G40.111	Localization-related (focal) (partial) symptomatic epilepsy and epileptic syndromes with simple partial seizures, intractable, with status epilepticus
G40.119	Localization-related (focal) (partial) symptomatic epilepsy and epileptic syndromes with simple partial seizures, intractable, without status epilepticus
G40.211	Localization-related (focal) (partial) symptomatic epilepsy and epileptic syndromes with complex partial seizures, intractable, with status epilepticus
G40.219	Localization-related (focal) (partial) symptomatic epilepsy and epileptic syndromes with complex partial seizures, intractable, without status epilepticus
G40.311	Generalized idiopathic epilepsy and epileptic syndromes, intractable, with status epilepticus
G40.319	Generalized idiopathic epilepsy and epileptic syndromes, intractable, without status epilepticus
G40.A11	Absence epileptic syndrome, intractable, with status epilepticus
G40.A19	Absence epileptic syndrome, intractable, without status epilepticus
G40.411	Other generalized epilepsy and epileptic syndromes, intractable, with status epilepticus
G40.419	Other generalized epilepsy and epileptic syndromes, intractable, without status epilepticus
G40.803	Other epilepsy, intractable, with status epilepticus
G40.804	Other epilepsy, intractable, without status epilepticus
G40.813	Lennox-Gastaut syndrome, intractable, with status epilepticus
G40.814	Lennox-Gastaut syndrome, intractable, without status epilepticus

Reviews, Revisions, and Approvals	Date	Approval Date
Developed PA Policy	11/17	1/18
Changed “partial onset” to “focal onset” throughout to reflect seizure classification changes made by the International League Against Epilepsy in 2017. References reviewed and updated.	08/18	
Updated background with additional information on non-implantable VNS. References reviewed and updated.	03/19	
References reviewed and updated. Added CPT code-61888. Added ICD-10 code, G40.311 Specialist review.	05/2020	
Added additional investigational indications for VNS to section II. References reviewed and updated. Removed ICD-10 Codes: G40.001, G40.009, G40.201, G40.209, G40.309, G40.A09, G40.409, G40.509, G40.802, G40.909, G40.911 and G40.919. Added ICD-10: G40.813, G40.814. Added new HCPCS code K1020 to a new table of HCPCS codes that do not support coverage criteria. “Experimental/investigational” verbiage replaced with descriptive language in policy statement II and III. Replaced “member” with “member/enrollee.”	06/2021	

References

1. 2017 Revised Classification of Seizures. (2016, December 22). Retrieved from <https://www.epilepsy.com/article/2016/12/2017-revised-classification-seizures>
2. Agency for Healthcare Research and Quality: Evidence report technology assessment, number 77, management of treatment related epilepsy, May 2003.
3. A.T. Berg, B.G. Vickrey and F.M. Testa et al., How long does it take for epilepsy to become intractable? A prospective investigation, Ann Neurol 60 (2006), pp. 73–9.
4. Ben-Menachem E, Revesz D, Simon BJ, Silberstein S. Surgically implanted and non-invasive vagus nerve stimulation: a review of efficacy, safety and tolerability. Eur J Neurol. 2015 Sep; 22(9): 1260–1268.
5. Bauer S, Baier H, Baumgartner C, et al. Transcutaneous Vagus Nerve Stimulation (tVNS) for Treatment of Drug-Resistant Epilepsy: A Randomized, Double-Blind Clinical Trial (cMPsE02). Brain Stimul. 2016 May-Jun;9(3):356-63. Epub 2016 Jan 20.
6. Bischel MD: Apollo’s Medical Review Criteria Guidelines for Managed Care: Vagus nerve stimulation (VNS) for the treatment of refractory seizures. Twelfth edition, 2013.
7. Boon P, Vonck K, van Rijckevorsel K, et al. A prospective, multicenter study of cardiac-based seizure detection to activate vagus nerve stimulation. Seizure. 2015 Nov;32:52-61. doi: 10.1016/j.seizure.2015.08.011. Epub 2015 Sep 21.
8. Center for Devices and Radiological Health (CDRH). P970003/S50. VNS Therapy System [premarket approval letter]. July 15, 2005b. Food and Drug Administration.
9. Fang J, Rong P, Hong Y, et al. Transcutaneous Vagus Nerve Stimulation Modulates Default Mode Network in Major Depressive Disorder. Biol Psychiatry. 2015 Apr 2. pii: S0006-3223(15)00274-7. doi: 10.1016/j.biopsych.2015.03.025.

10. Fisher RS, Afra P, Macken M, et al. Automatic Vagus Nerve Stimulation Triggered by Ictal Tachycardia: Clinical Outcomes and Device Performance-The U.S. E-37 Trial. *Neuromodulation*. 2016 Feb;19(2):188-95. doi: 10.1111/ner.12376. Epub 2015 Dec 13.
11. Guan G, Karsy M, Ducis K, Bollo RJ. Surgical strategies for pediatric epilepsy. *Translational Pediatrics*. 2016 Apr; 5(2): 55–66.
12. Hampel KG, Vatter H, Elger CE, et al. Cardiac-based vagus nerve stimulation reduced seizure duration in a patient with refractory epilepsy. *Seizure*. 2015 Mar;26:81-5. doi: 10.1016/j.seizure.2015.02.004. Epub 2015 Feb 14.
13. Hayes Medical Technology Directory: Vagus nerve stimulation for epilepsy. June 9, 2014. Archived July 9, 2019.
14. Hein E, Nowak M, Kiess O, et al. Auricular transcutaneous electrical nerve stimulation in depressed patients: a randomized controlled pilot study. *J Neural Transm (Vienna)*. 2013;120(5):821.
15. Holtzheimer PE. Unipolar depression in adults: Treatment with surgical approaches. In: UpToDate, Roy-Byrne PP (Ed), UpToDate, Waltham, MA. Accessed July 27, 2020.
16. Schachter SC. Vagus nerve stimulation therapy for the treatment of epilepsy. In: UpToDate, Garcia P, (Ed), UpToDate, Waltham, MA. Accessed July 28, 2020.
17. Klinkenberg S, et al. Vagus nerve stimulation in children with intractable epilepsy: a randomized controlled trial. *Developmental Medicine and Child Neurology* 2012;54(9):855-61.
18. Kwan P, Brodie MJ. Early identification of refractory epilepsy, *N Engl J Med* 342 (2000), pp. 314–39.
19. Medical News Today: New FDA approved device offers hope to depressed patients, July 2005.
20. Medicare Coverage Database: NCD for Vagus Nerve Stimulation (160.18). . Centers for Medicare and Medicaid. Effective 2/15/19.
http://www.cms.hhs.gov/mcd/viewncd.asp?ncd_id=160.18&ncd_version=2&basket=ncd%3A160%2E18%3A2%3AVagus+Nerve+Stimulation+for+Treatment+of+Seizures
21. Morris GL, et al. Evidence-based guideline update: Vagus nerve stimulation for the treatment of epilepsy. *American Academy of Neurology*. August 28, 2013.
22. Panebianco M, Rigby A, Weston J, Marson AG. Vagus nerve stimulation for partial seizures. *Cochrane Database Syst Rev*. 2015 Apr 3;(4):CD002896.
23. Schachter SC. Overview of the management of epilepsy in adults. In: UpToDate, Garcia P, Edlow JA (Eds), UpToDate, Waltham, MA. Accessed July 28, 2020..
24. Sirven, JI. Evaluation and management of drug-resistant epilepsy. In: UpToDate, Garcia P (Ed), UpToDate, Waltham, MA. Accessed July 28, 2020..
25. Wilfong A. Seizures and epilepsy in children: Refractory seizures and prognosis. In: UpToDate, Nordli DR (Ed), UpToDate, Waltham, MA. July 28, 2020.
26. Tassorelli C, Grazzi L, de Tommaso M, et al. Noninvasive vagus nerve stimulation as acute therapy for migraine: The randomized PRESTO study. *Neurology*. 2018 Jul 24;91(4):e364-e373. doi: 10.1212/WNL.0000000000005857. Epub 2018 Jun 15.
27. Silberstein SD, Calhoun AH, Lipton RB, et al. Chronic migraine headache prevention with noninvasive vagus nerve stimulation: The EVENT study. *Neurology*. 2016 Aug 2;87(5):529-38. doi: 10.1212/WNL.0000000000002918. Epub 2016 Jul 13.
28. Silberstein SD, Mechtler LL, Kudrow DB, et al. Non-invasive vagus nerve stimulation for the acute treatment of cluster headache: findings from the randomized, double-blind, sham-

- controlled ACT1 study. *Headache*. 2016;56(8):1317–1332. (150 participants- company involved in interpretation)
29. Goadsby PJ, de Coo IF, Silver N, et al. ACT2 Study Group. Non-invasive vagus nerve stimulation for the acute treatment of episodic and chronic cluster headache: a randomized, double-blind, sham-controlled ACT2 study. *Cephalalgia*. 2018;38(5):959–969. (92 participants)
 30. American Headache Society. The American Headache Society Position Statement On Integrating New Migraine Treatments Into Clinical Practice. *Headache*. 2019 Jan;59(1):1-18. doi: 10.1111/head.13456. Epub 2018 Dec 10.
 31. May A. Cluster headache: Treatment and prognosis. In; UpToDate. Swanson JW (Ed) UpToDate, Waltham MA. Accessed July 28, 2020.
 32. Hayes Health Technology Assessment. Vagus Nerve Stimulation For Treatment-Resistant Depression. February 21, 2019. Update May 18, 2020.
 33. Smith JH. Acute treatment of migraine in adults. In: UpToDate, Swanson J W (Ed), UpToDate, Waltham, MA. Accessed July 28, 2020.
 34. Hayes Health Technology Assessment. Noninvasive Vagus Nerve Stimulation with gammaCore for Prevention or Treatment of Cluster Headache. May 12, 2020.