

Clinical Policy: Vagus Nerve Stimulation

Reference Number: PA.CP.MP.12

Effective Date: 01/18

Last Review Date: 07/18/2022

[Coding Implications](#)

[Revision Log](#)

Description

Vagus nerve stimulation (VNS) has been used in the treatment of epilepsy and has been studied for the treatment of refractory depression and other indications. Electrical pulses are delivered to the cervical portion of the vagus nerve by an implantable device called a neurocybernetic prosthesis. Chronic intermittent electrical stimulation of the left vagus nerve is designed to treat medically refractory epilepsy. It has recently been introduced and approved by the FDA as an adjunctive therapy for treatment-resistant major depression.

Policy/Criteria

- I. It is the policy of Pennsylvania Health and Wellness® (PHW), that VNS is **medically necessary** in patients with medically refractory seizures who meet all of the following:
 - A. Diagnosis of local onset (formerly partial onset) seizures or generalized onset seizures;
 - B. Intractable epilepsy (both):
 1. Failure of at least 1 year of adherent therapy of at least two anti-seizure drugs, *and*
 2. Continued seizures which have a major impact on activities of daily living; *and*
 - C. Not a suitable candidate for or has failed resective epilepsy surgery;
 - D. Request is for an FDA-approved device.
- II. It is the policy of PHW that the safety and efficacy of VNS therapy has not been proven for any other conditions, including but not limited to the following:
 - A. Refractory (treatment resistant) major depression or bipolar disorder;
 - B. Obesity;
 - C. Headaches;
 - D. Cognitive impairment associated with Alzheimer's disease;
 - E. Addiction;
 - F. Anxiety Disorders;
 - G. Autism;
 - H. Eating Disorders;
 - I. Cancer;
 - J. Crohn's Disease;
 - K. Essential trauma;
 - L. Fibromyalgia;
 - M. Heart failure;
 - N. Impaired glucose tolerance/pre-diabetes;
 - O. Inflammation;
 - P. Overweight and obesity;
 - Q. Obsessive-compulsive disorder;
 - R. Panic disorder;
 - S. Post-traumatic stress disorder;
 - T. Prader-Willi Syndrome;
 - U. Sjogren's Syndrome;
 - V. Rheumatoid arthritis;

CLINICAL POLICY

Vagus Nerve Stimulation



- W. Schizophrenia;
- X. Sleep disorders;
- Y. Stroke;
- Z. Tinnitus;
- AA. Tourette's syndrome;
- BB. Traumatic brain injury

III. It is the policy of PHW that the current research does not support the use of following types of VNS therapy over other currently available alternatives, due to the lack of large, high-quality studies supporting their use:

- A. Aspire SR Model 106 (Cyberonics) for vagus nerve stimulation;
- B. Transcutaneous VNS or active auricular transcutaneous electrical nerve stimulation.

Removal of Implant

Less than 0.5 percent of all patients have had the device removed. It can be turned off in the physician's office if the patient feels it is not helping or if the patient cannot tolerate the stimulation. If the device needs to be removed, only the pulse generator is removed, as attempting to remove the electrodes from around the nerve can cause damage and is not recommended.

Background

The vagus nerve stimulator is a pacemaker-like device implanted under the skin in the left side of the chest through a small incision, with a second small incision made at the base of the neck. The surgery is performed under local, regional, or general anesthesia and lasts 45 minutes to two hours. Most often it is performed as an outpatient surgery but some patients need to stay in the hospital overnight following surgery.

Partial (focal) seizures

Several studies have been done evaluating the safety and effectiveness of vagus nerve stimulation for treatment of epilepsy. A randomized active-control trial known as the E05 study found that 94 patients (of the total 254 patients in the study) receiving high stimulation showed an average reduction in seizure frequency, compared to baseline, of 28% versus 15% reduction in the 102 patients receiving low stimulation. A total of 310 patients completed the E03 and E05 double-blinded trials. Mean decline of seizure frequency overall was about 25-30% compared to baseline. Clinical experience has shown that improvement in seizures is maintained, or may even increase over time, but these data are based on uncontrolled observations. Side effects in both studies were similar and included hoarseness and occasional shortness of breath.

Although questions regarding patient selection criteria, optimal stimulation parameters, and cost-effectiveness in the United States remain under investigation, there is sufficient evidence regarding the benefit and safety of VNS to conclude that VNS may improve health outcomes in patients with medically refractory partial-onset seizures who are not suitable candidates for surgery or in whom surgical treatment has failed.

Generalized seizures

Study results suggest VNS may be effective for generalized epilepsy (Karciski & Schachter, 2015). However, case series and observational studies constitute the majority of available evidence. Although VNS is not currently FDA approved for the treatment of generalized seizures, it is often used in children and other patients, and in Europe is approved as adjunct therapy for epileptic disorders predominantly characterized by generalized or focal seizures that are refractory to antiseizure medications. In addition, the National Institute for Health and Care Excellence (NICE) recommends VNS for focal and generalized seizures, and the Scottish Intercollegiate Guidelines Network (SIGN) guidelines recommend VNS for epilepsy in patients unsuitable for respective surgery without stipulating seizure type.

Depression

VNS was FDA-approved for treatment resistant depression in 2005. However, VNS has no rigorous research data proving it is efficacious for treatment-resistant, unipolar major depression. Open-label studies suggest VNS may be effective; however, these are at risk for bias due to placebo effects. The one randomized trial of VNS for depression found no benefit, with outcomes comparable for active and sham treatment (response rates of 15 vs. 10 percent). In addition, there is a lack of thorough safety data for the use of VNS in depression.

Other Investigational Indications

Ongoing research efforts continue to investigate the role of vagus nerve stimulation (VNS) for the treatment of a variety of indications, including but not limited to cognitive deficits in Alzheimer's disease, resistant obesity, and headaches. Data supporting the long-term safety and efficacy from large clinical trials of VNS for the treatment of these indications, however, continue to be lacking.

AspireSR Model 106 (Cyberonics) for Vagus Nerve Stimulation

The AspireSR Model 106 (Cyberonics Inc.) received FDA Premarket Approval (PMA) in February 2014. The newest modification to the vagus nerve stimulation (VNS) implant detects tachycardia heart rates, which may be associated with an impending seizure, and automatically delivers stimulation to the vagus nerve. Like its predecessors, the AspireSR can also deliver stimulation in the normal and magnet modes. However, when programmed for AutoStim mode, the AspireSR requires no patient interaction to trigger the delivery of electrical stimulation. The AutoStim mode should not be used in patients with significant arrhythmias being treated with pacemakers and/or an implantable defibrillator, beta-blockers, or any other treatment that may impact the intrinsic heart rate.

A few small, preliminary studies and case reports have evaluated the AspireSR Model 106, and have shown positive results (Boon et al., 2015; Fisher et al., 2015; Schneider et al., 2015; Hampel et al, 2015). However, there is insufficient evidence to establish the safety and efficacy of the AspireSR Model 106 in reducing seizures until further, high quality trials establish its clinical value.

Transcutaneous Vagus Nerve Stimulation

Transcutaneous vagus nerve stimulation (tVNS) has been proposed as a noninvasive alternative to implantable VNS for a variety of indications, including, but not limited to epilepsy, major depression, chronic tinnitus and headaches. Currently, there are two main ways to apply tVNS.

One is to apply stimulation on the ear and the other is cervical noninvasive VNS, superficially applying stimulation in the vicinity of the vagus nerve using a specially designed device, (e.g. gammaCore). Noninvasive auricular tVNS stimulates the afferent auricular branch of the vagus nerve located medial of the tragus at the entry of the acoustic meatus. Given that the right vagal nerve has efferent fibers to the heart, tVNS is safe to be performed only in the left ear. tVNS has been proposed to study cognitive functioning in patients with epilepsy and major depression. The rationale is that direct stimulation of the afferent nerve fibers on the ear area with afferent vagus nerve distribution should produce a similar effect as classic VNS in reducing depressive symptoms without the burden of surgical intervention. A noninvasive, transcutaneous vagal nerve stimulator has been in use in Europe. Although no randomized studies have been done in patients with epilepsy, it appears promising in one pilot study.¹⁸ Small studies have shown positive results with tVNS for the treatment of depression (Hein et al, 2013; Fang et al, 2016). Additional, larger, peer-reviewed studies, with longer follow-up are necessary to determine the long-term safety and efficacy of transcutaneous VNS for depression.

gammaCore Sapphire™ (ElectroCore, LLC), is a hand-held prescription device that is placed externally on the side of the neck in the vicinity of the vagus nerve to deliver a low voltage electric signal to the nerve's afferent fibers. gammaCore has received FDA approval for the treatment of both episodic cluster and migraine headaches and more recently for the prevention of cluster headaches (CH). gammaCore delivers up to 30 stimulations in a 24-hour period, each lasting 2 minutes. The patient controls the intensity level. Once the maximum daily number of treatments has been reached, the device will not deliver any more treatments until the following 24-hour period. gammaCore is rechargeable and includes a charging case to charge the device. A gammaCore refill card is used to load the device with days of therapy based on a healthcare provider's prescription.

In the randomized PRESTO study, noninvasive vagus nerve stimulation (nVNS.) was superior to sham in the treatment of episodic migraine for pain freedom at 30 minutes and 60 minutes after the first treated attack.²⁸ In both the ACT1 and ACT2 trials, nVNS was superior to sham therapy in episodic CH but not in chronic CH.^{28,29} Another 2020 randomized, double-blind, sham-controlled clinical trial showed when comparing nVNS with sham, no statistically significant differences were found with regards to the primary endpoint of pain freedom at 120 minutes, although differences were found with various secondary endpoints and post hoc analysis.³²

Preliminary clinical trials of nVNS in various primary headache disorders are encouraging, but, for future studies, it is important to conduct large, properly blinded and controlled trials by independent researchers. Additionally, most studies nVNS devices enrolled participants who did not respond sufficiently to oral drug treatment; thus, the role of neurostimulation in an average population of migraine patients remains unknown.³³

The American Headache Society position statement on integrating new migraine treatments into clinical practice note that empirically validated behavioral treatments with Grade A evidence for the prevention of migraine, including cognitive behavioral therapy, biofeedback, and relaxation therapies, should be considered in the management of migraine. These modalities may also be used alone or in addition to pharmacologic treatment. They note further that several noninvasive devices have been developed and approved by the FDA for the treatment of patients with

CLINICAL POLICY

Vagus Nerve Stimulation



migraine.(i.e., single-pulse transcranial magnetic stimulation, electrical trigeminal nerve stimulation and nVNS.) Patients who prefer nondrug therapies and those who have failed to respond to, have contraindications to, or poor tolerability with pharmacotherapy may be candidates for neuromodulation.³⁰

Per UpToDate, “There are several promising but unproven methods using neurostimulation to treat medically refractory cluster headache, including sphenopalatine ganglion stimulation, occipital nerve stimulation, noninvasive VNS, and deep brain stimulation. All are investigational and require further study to confirm long-term benefit and safety.”

Coding Implications

This clinical policy references Current Procedural Terminology (CPT®). CPT® is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2021, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

CPT®*	Description
61885	Insertion or replacement of cranial neurostimulator pulse generator or receiver, direct or inductive coupling; with connection to a single electrode array
61886	Insertion or replacement of cranial neurostimulator pulse generator or receiver, direct or inductive coupling; with connection to two or more electrode arrays
61888	Revision or removal of cranial neurostimulator pulse generator or receiver
64553	Percutaneous implantation of neurostimulator electrodes; cranial nerve
64568	Incision for implantation of cranial nerve (eg, vagus nerve) neurostimulator electrode array and pulse generator
64569	Revision or replacement of cranial nerve (eg, vagus nerve) neurostimulator electrode array, including connection to existing pulse generator
64570	Removal of cranial nerve (eg, vagus nerve) neurostimulator electrode array and pulse generator

HCPCS Codes that Support Coverage Criteria

HCPCS Codes	Description
C1767	Generator, neurostimulator (implantable), nonrechargeable
C1778	Lead, neurostimulator (implantable)
C1816	Receiver and/or transmitter, neurostimulator (implantable)
C1883	Adaptor/extension, pacing lead or neurostimulator lead (implantable)
L8680	Implantable neurostimulator electrode, each
L8681	Patient programmer (external) for use with implantable programmable neurostimulator pulse generator, replacement only

CLINICAL POLICY
Vagus Nerve Stimulation



HCPCS Codes	Description
L8682	Implantable neurostimulator radiofrequency receiver
L8683	Radiofrequency transmitter (external) for use with implantable neurostimulator radiofrequency receiver
L8685	Implantable neurostimulator pulse generator, single array , rechargeable, includes extension
L8686	Implantable neurostimulator pulse generator, single array, nonrechargeable, includes extension
L8687	Implantable neurostimulator pulse generator, dual array, rechargeable, includes extension
L8688	Implantable neurostimulator pulse generator, dual array, nonrechargeable, includes extension
L8689	External recharging system for battery (internal) for use with implanted neurostimulator, replacement only

HCPCS Codes that Do Not Support Coverage Criteria

HCPCS Codes	Description
K1020	Noninvasive vagus nerve stimulator

ICD-10-CM Diagnosis Codes that Support Coverage Criteria

ICD-10-CM Code	Description
G40.011	Localization-related (focal) (partial) idiopathic epilepsy and epileptic syndromes with seizures of localized onset, intractable, with status epilepticus
G40.019	Localization-related (focal) (partial) idiopathic epilepsy and epileptic syndromes with seizures of localized onset, intractable, without status epilepticus
G40.111	Localization-related (focal) (partial) symptomatic epilepsy and epileptic syndromes with simple partial seizures, intractable, with status epilepticus
G40.119	Localization-related (focal) (partial) symptomatic epilepsy and epileptic syndromes with simple partial seizures, intractable, without status epilepticus
G40.211	Localization-related (focal) (partial) symptomatic epilepsy and epileptic syndromes with complex partial seizures, intractable, with status epilepticus
G40.219	Localization-related (focal) (partial) symptomatic epilepsy and epileptic syndromes with complex partial seizures, intractable, without status epilepticus
G40.311	Generalized idiopathic epilepsy and epileptic syndromes, intractable, with status epilepticus
G40.319	Generalized idiopathic epilepsy and epileptic syndromes, intractable, without status epilepticus
G40.A11	Absence epileptic syndrome, intractable, with status epilepticus
G40.A19	Absence epileptic syndrome, intractable, without status epilepticus
G40.411	Other generalized epilepsy and epileptic syndromes, intractable, with status epilepticus
G40.419	Other generalized epilepsy and epileptic syndromes, intractable, without status epilepticus

ICD-10-CM Code	Description
G40.803	Other epilepsy, intractable, with status epilepticus
G40.804	Other epilepsy, intractable, without status epilepticus
G40.813	Lennox-Gastaut syndrome, intractable, with status epilepticus
G40.814	Lennox-Gastaut syndrome, intractable, without status epilepticus

Reviews, Revisions, and Approvals	Date	Approval Date
Developed PA Policy	11/17	1/18
Changed “partial onset” to “focal onset” throughout to reflect seizure classification changes made by the International League Against Epilepsy in 2017. References reviewed and updated.	08/18	
Updated background with additional information on non-implantable VNS. References reviewed and updated.	03/19	
References reviewed and updated. Added CPT code-61888. Added ICD-10 code, G40.311 Specialist review.	05/2020	
Added additional investigational indications for VNS to section II. References reviewed and updated. Removed ICD-10 Codes: G40.001, G40.009, G40.201, G40.209, G40.309, G40.A09, G40.409, G40.509, G40.802, G40.909, G40.911 and G40.919. Added ICD-10: G40.813, G40.814. Added new HCPCS code K1020 to a new table of HCPCS codes that do not support coverage criteria. “Experimental/investigational” verbiage replaced with descriptive language in policy statement II and III. Replaced “member” with “member/enrollee.”	06/2021	
Annual review. Changed “review date” in the header to “date of last revision” and “date” in the revision log header to “revision date.” Background updated with additional study on nVNS for migraine headaches. References reviewed and updated. Reviewed by specialist.	07/18/2022	

References

1. 2017 Revised Classification of Seizures. Epilepsy Foundation.
<https://www.epilepsy.com/article/2016/12/2017-revised-classification-seizures>. Published December 2020. Accessed July 27, 2021.
2. Chapell R, Reston J, Snyder D, Treadwell J, Treager S, Turkelson C. 77 Management of treatment related epilepsy. In: AHRQ Evidence Report Summaries. Rockville, MD: Agency for Healthcare Research and Quality; 1998-2005.
<https://www.ncbi.nlm.nih.gov/books/NBK11892/>. Accessed July 27, 2021.
3. Berg AT, Langfitt J, Shinnar S, et al. How long does it take for partial epilepsy to become intractable?. *Neurology*. 2003;60(2):186-190. doi:10.1212/01.wnl.0000031792.89992.ec
4. Ben-Menachem E, Revesz D, Simon BJ, Silberstein S. Surgically implanted and non-invasive vagus nerve stimulation: a review of efficacy, safety and tolerability. *Eur J Neurol*. 2015; 22(9): 1260–1268. doi:10.1111/ene.12629

5. Bauer S, Baier H, Baumgartner C, et al. Transcutaneous Vagus Nerve Stimulation (tVNS) for Treatment of Drug-Resistant Epilepsy: A Randomized, Double-Blind Clinical Trial (cMPsE02). *Brain Stimul*. 2016;9(3):356-63. doi: 10.1016/j.brs.2015.11.003
6. Boon P, Vonck K, van Rijckevorsel K, et al. A prospective, multicenter study of cardiac-based seizure detection to activate vagus nerve stimulation. *Seizure*. 2015;32:52-61. doi: 10.1016/j.seizure.2015.08.011
7. Center for Devices and Radiological Health (CDRH). P970003/S50. VNS Therapy System [premarket approval letter]. July 15, 2005. Food and Drug Administration.
8. Fang J, Rong P, Hong Y, et al. Transcutaneous Vagus Nerve Stimulation Modulates Default Mode Network in Major Depressive Disorder. *Biol Psychiatry*. 2016;79(4):266-273. doi:10.1016/j.biopsych.2015.03.025
9. Fisher RS, Afra P, Macken M, et al. Automatic Vagus Nerve Stimulation Triggered by Ictal Tachycardia: Clinical Outcomes and Device Performance-The U.S. E-37 Trial. *Neuromodulation*. 2016;19(2):188-195. doi:10.1111/ner.12376
10. Guan G, Karsy M, Ducis K, Bollo RJ. Surgical strategies for pediatric epilepsy. *Transl Pediatr*. 2016; 5(2): 55–66. doi: 10.21037/tp.2016.03.02
11. Hampel KG, Vatter H, Elger CE, et al. Cardiac-based vagus nerve stimulation reduced seizure duration in a patient with refractory epilepsy. *Seizure*. 2015;26:81-5. doi: 10.1016/j.seizure.2015.02.004
12. Hein E, Nowak M, Kiess O, et al. Auricular transcutaneous electrical nerve stimulation in depressed patients: a randomized controlled pilot study. *J Neural Transm (Vienna)*. 2013;120(5):821. doi: 10.1007/s00702-012-0908-6
13. Holtzheimer PE. Unipolar depression in adults: Treatment with surgical approaches. UpToDate. www.uptodate.com. Updated July 8, 2021. Accessed July 28, 2021.
14. Schachter SC. Vagus nerve stimulation therapy for the treatment of epilepsy. UpToDate. www.uptodate.com. Updated June 15, 2021. Accessed July 28, 2021.
15. Klinkenberg S, Aalbers MW, Vles JSH, et al. Vagus nerve stimulation in children with intractable epilepsy: a randomized controlled trial. *Dev Med Child Neurol*. 2012;54(9):855-61. doi: 10.1111/j.1469-8749.2012.04305.x
16. Kwan P, Brodie MJ. Early identification of refractory epilepsy. *N Engl J Med*. 2000;342(5):314-319. doi:10.1056/NEJM200002033420503
17. National coverage determination: vagus Nerve Stimulation (160.18). Centers for Medicare and Medicaid Services. <https://www.cms.gov/medicare-coverage-database/new-search/search.aspx>. Published February 15, 2019. Accessed July 28, 2021.
18. Morris GL 3rd, Gloss D, Buchhalter J, Mack KJ, Nickels K, Harden C. Evidence-based guideline update: Vagus nerve stimulation for the treatment of epilepsy. *Neurology*. 2013;81(16):1453-1459. doi:10.1212/WNL.0b013e3182a393d1
19. Panebianco M, Rigby A, Weston J, Marson AG. Vagus nerve stimulation for partial seizures. *Cochrane Database Syst Rev*. 2015;2015(4):CD002896. Published 2015 Apr 3. doi:10.1002/14651858.CD002896.pub2
20. Schachter SC. Overview of the management of epilepsy in adults. UpToDate. www.uptodate.com. Updated July 20, 2021. Accessed July 28, 2021.
21. Sirven, JI. Evaluation and management of drug-resistant epilepsy. UpToDate. www.uptodate.com. Updated June 7, 2021. Accessed July 28, 2020..
22. Wilfong A. Seizures and epilepsy in children: Refractory seizures. UpToDate. www.uptodate.com. Updated October 20, 2020. Accessed July 28, 2021.

23. Tassorelli C, Grazzi L, de Tommaso M, et al. Noninvasive vagus nerve stimulation as acute therapy for migraine: The randomized PRESTO study. *Neurology*. 2018;91(4):e364-e373. doi:10.1212/WNL.0000000000005857
24. Silberstein SD, Calhoun AH, Lipton RB, et al. Chronic migraine headache prevention with noninvasive vagus nerve stimulation: The EVENT study. *Neurology*. 2016;87(5):529-538. doi:10.1212/WNL.0000000000002918
25. Silberstein SD, Mechtler LL, Kudrow DB, et al. Non-invasive vagus nerve stimulation for the acute treatment of cluster headache: findings from the randomized, double-blind, sham-controlled ACT1 study. *Headache*. 2016;56(8):1317-1332. doi:10.1111/head.12896.
26. Goadsby PJ, de Coo IF, Silver N, et al. ACT2 Study Group. Non-invasive vagus nerve stimulation for the acute treatment of episodic and chronic cluster headache: a randomized, double-blind, sham-controlled ACT2 study. *Cephalalgia*. 2018;38(5):959-969. doi:10.1177/0333102417744362
27. American Headache Society. The American Headache Society Position Statement On Integrating New Migraine Treatments Into Clinical Practice [published correction appears in *Headache*. 2019 Apr;59(4):650-651]. *Headache*. 2019;59(1):1-18. doi:10.1111/head.13456
28. May A. Cluster headache: Treatment and prognosis. UpToDate. www.uptodate.com. Updated February 25, 2021. Accessed July 28, 2021.
29. Hayes Health Technology Assessment. Vagus nerve stimulation for treatment-resistant depression. Hayes. www.hayesinc.com. Published February 21, 2019 (updated May 18, 2020). Accessed July 28, 2021.
30. Smith JH. Acute treatment of migraine in adults. UpToDate. www.uptodate.com. Updated June 9, 2021. Accessed July 28, 2021.
31. Hayes Health Technology Assessment. Noninvasive vagus nerve stimulation with gammacore for prevention or treatment of cluster headache. Hayes. www.hayesinc.com. Published May 12, 2020. Accessed July 28, 2021.
32. Blech B, Starling AJ, Marks LA, Wingerchuk DM, O'Carroll CB. Is Noninvasive Vagus Nerve Stimulation a Safe and Effective Alternative to Medication for Acute Migraine Control?. *Neurologist*. 2020;25(4):97-100. doi:10.1097/NRL.0000000000000274
33. Evers S. Non-Invasive Neurostimulation Methods for Acute and Preventive Migraine Treatment-A Narrative Review. *J Clin Med*. 2021;10(15):3302. Published 2021 Jul 27. doi:10.3390/jcm10153302