

Clinical Policy: Sacroiliac Joint Interventions for Pain Management

Reference Number: PA.CP.MP.166 Effective Date: 09/18 Date of Last Review: 07/28/2022 Coding Implications <u>Revision Log</u>

Description

Treatment for sacroiliac joint (SIJ) dysfunction is usually conservative (non-surgical) and focuses on trying to restore normal motion in the joint. In patients who have failed to respond to conservative therapy, an SIJ injection can be helpful for both diagnostic and therapeutic purposes. SIJ injections into the synovial sac of the SIJ may provide immediate and significant pain relief.

Policy/Criteria

It is the policy of PA Health and Wellness[®] (PHW) that invasive pain management procedures performed by a physician are **medically necessary** when *the relevant criteria are met and the patient receives only one procedure per visit, with or without radiographic guidance.*

- I. Sacroiliac joint injections are medically necessary for the following indications:
 - A. One diagnostic sacroiliac joint (SIJ) injection for SIJ pain, all of the following:
 - 1. Somatic or nonradicular low back and lower extremity pain below the level of L5 vertebra that interferes with ADLs for at least 3 months;
 - 2. Tenderness by palpation present over SIJ;
 - 3. There is a positive response to at least three SIJ pain provocation tests (distraction, compression, thigh thrust, Gaenslen's, Patrick's test/FABER test, or sacral thrust);
 - 4. The member/enrollee has failed to respond to conservative therapy including all of the following:
 - a. ≥ 6 weeks chiropractic, physical therapy or prescribed home exercise program;
 - b. NSAID \geq 3 weeks or NSAID contraindicated or not tolerated;
 - c. ≥ 6 weeks activity modification;
 - 5. Clinical findings and imaging studies, when available, lack obvious evidence for discrelated or facet joint pain;
 - 6. No other possible diagnosis is more likely.
 - **B.** A second diagnostic or confirmatory sacroiliac joint injection when pain was improved by at least 75% after the first diagnostic SIJ injection and at least 2 weeks have passed since the initial injection.
 - C. Subsequent therapeutic SIJ injections for recurrence of pain, all of the following:
 - Initial therapeutic injection(s) led to ≥ 50% relief and functional improvement for at least 2 months;
 - 2. Request is for SIJ administered for temporary relief of lower back pain in conjunction with other noninvasive treatment methods (e.g., to participate in physical therapy), and not as a stand-alone therapy;
 - 3. SIJ injection is given at intervals at least 2 months apart;
 - 4. Less than 4 therapeutic SIJ injections have been given at the same site in the last 12 months.



II. It is the policy of PHW that if pain does not improve by $\geq 75\%$ after the second diagnostic SIJ injections, *subsequent SIJ injections* are **not medically necessary** because effectiveness has not been established.

III. It is the policy of PHW that continuation of injections beyond 12 months is considered

- **IV. not medically necessary** because effectiveness and safety has not been established. When more definitive therapies cannot be tolerated or provided, consideration will be made on a case by case basis.
- V. It is the policy of PHW that *sacroiliac nerve blocks* are considered **not medically necessary** because effectiveness has not been established.
- **VI.** It is the policy of PHW that *radiofrequency neurotomy (conventional, cooled, and pulsed)* of the SIJ is considered **not medically necessary** because effectiveness has not been established. High-quality studies are lacking for conventional and pulsed radiofrequency neurotomy of the SIJ. For cooled radiofrequency neurotomy, additional well-designed studies are needed to evaluate effectiveness.

*Note: requests for services considered **not medically necessary** are to be reviewed on a case-by-case basis. Medical Director/clinical reviewer must override criteria when, in his/her professional judgement, the requested item is medically necessary.

Background

Sacroiliac Joint Injections

Treatment for sacroiliac joint dysfunction is usually conservative (non-surgical) and focuses on trying to restore normal motion in the joint. In patients who have failed 4 to 6 weeks of a comprehensive exercise program, local icing, mobilization/manipulation and NSAIDs, an SIJ injection can be helpful for both diagnostic and therapeutic purposes. SIJ injections into the synovial sac of the SIJ may provide immediate and significant pain relief. Adding a steroid to the solution injected may help to reduce any inflammation that may exist within the joint(s) and result in a prolonged period of freedom from pain.

Several studies without control groups have concluded that SIJ injections improve pain in the short term.¹ However, the majority of studies have small sample sizes and most lack comparison to standard interventions such as physical therapy.

A study by Visser et al. evaluated the effect of manual therapy and physiotherapy versus SIJ injection for low back and leg pain using a single-blinded randomized trial of treatment for 51 patients with SIJ-related leg pain. The effect of the treatment was evaluated after 6 and 12 weeks. Manual therapy had a significantly better success rate than physiotherapy (p = 0.003). The authors concluded in the small single-blinded prospective study, manual therapy appeared to be the choice of treatment for patients with SIJ-related leg pain.¹ A second choice of treatment to be considered is an intra-articular injection.¹

SIJ Radiofrequency Neurotomy



A growing number of studies have assessed the effect of treatment with radiofrequency denervation on SIJ pain, with mixed results. One study found no difference between conventional radiofrequency ablation (RFA) and a sham treatment on pain relief.² A 2017 publication of 3 randomized controlled trials of 681 participants with chronic low back pain found no statistically significant improvement in pain from treatment with a standardized exercise program plus RFA, versus the standardized exercise program alone.³ A few fair to poor quality studiesfound positive results from conventional and cooled RFA.¹ The American Society of Interventional Pain Physicians 2013 guidelines rate the evidence for cooled RFA as fair, and limited for conventional and pulsed RFA.⁴ Due to varying anatomy, there is no standard approach to denervation of the sacroiliac joint, nor clearly defined criteria for patient selection.¹

Coding Implications

This clinical policy references Current Procedural Terminology (CPT[®]). CPT[®] is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2020, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

CPT [®] Codes	Description
27096	Injection procedure for sacroiliac joint, anesthetic/steroid, with image guidance (fluoroscopy or CT) including arthrography when performed

CPT code that does not support coverage criteria

CPT®	Description
Codes	
64451	Injection(s), anesthetic agent(s) and/or steroid; nerves innervating the sacroiliac
	joint, with image guidance (ie, fluoroscopy or computed tomography)
64625	Radiofrequency ablation, nerves innervating the sacroiliac joint, with image guidance
	(ie, fluoroscopy or computed tomography)

HCPCS code that supports coverage criteria

HCPCS	Description
Codes	
G0260	Injection procedure for sacroiliac joint; provision of anesthetic, steroid and/or other
	therapeutic agent, with or without arthrography

ICD-10-CM Diagnosis Codes that Support Coverage Criteria

+ Indicates a code requiring an additional character

ICD-10-CM	Description
Code	
M43.08	Spondylolysis, sacral and sacrococcygeal region
M46.1	Sacroiliitis, not elsewhere classified



ICD-10-CM	Description
Code	
M47.818	Spondylosis without myelopathy or radiculopathy, sacral and sacrococcygeal
	region
M53.3	Sacrococcygeal disorders, not elsewhere classified
M53.87	Other specified dorsopathies, lumbosacral region
M53.88	Other specified dorsopathies, sacral and sacrococcygeal region
M54.30-M54.32	Sciatica
M54.40-M54.42	Lumbago with sciatica
M54.5	Low back pain
M54.81	Other dorsalgia
M54.9	Dorsalgia, unspecified

Reviews, Revisions, and Approvals	Revision Date	Approva l Date
New policy developed. Split from retired CP.MP.118 Injections for Pain Management. No criteria change.	09/18	10/18
Annual review of policy. Minor wording changes to match language in other pain injection policies. Added New 2020 CPT code- 64625 as not medically necessary. Added criteria stating SIJ nerve blocks as not medically necessary, along with code 64451.References reviewed and updated, with two additional references added. Specialty review completed. Reworded II. for clarity.	06/2020	
Annual review completed. Updated I.A. to specify that the criteria apply to therapeutic injections as well as diagnostic. Updated I.B. to state "A second diagnostic or confirmatory sacroiliac joint injection when pain was improved by at least 75% after the first diagnostic SIJ injection", rather than that pain did not improve. I.C. updated to specify "therapeutic" SIJ injection. II was changed from 50% to 75%. Updated background. Replaced member with member/enrollee in all instances. Changed "review date" in the header to "date of last revision" and "date" in the revision log header to "revision date." References reviewed, updated, and reformatted.	7/28/2022	

References

- 1. Visser LH, Woudenberg NP, de Bont J, et al. Treatment of the sacroiliac joint in patients with leg pain: a randomized-controlled trial. *Eur Spine J*. 2013;22(10):2310-2317. doi:10.1007/s00586-013-2833-2
- van Tilburg CW, Schuurmans FA, Stronks DL, Groeneweg JG, Huygen FJ. Randomized Sham-controlled Double-Blind Multicenter Clinical Trial to Ascertain the Effect of Percutaneous Radiofrequency Treatment for Sacroiliac Joint Pain: Three-month Results. *Clin* J Pain. 2016;32(11):921-926. doi:10.1097/AJP.00000000000351
- 3. Juch JNS, Maas ET, Ostelo RWJG, et al. Effect of Radiofrequency Denervation on Pain Intensity Among Patients With Chronic Low Back Pain: The Mint Randomized Clinical



Trials [published correction appears in JAMA. 2017 Sep 26;318(12):1188]. *JAMA*. 2017;318(1):68-81. doi:10.1001/jama.2017.7918

- 4. Manchikanti L, Abdi S, Atluri S, et al. An update of comprehensive evidence-based guidelines for interventional techniques in chronic spinal pain. Part II: guidance and recommendations. *Pain Physician*. 2013;16(2 Suppl):S49-S283.
- 5. Radiofrequency ablation for sacroiliac joint denervation for chronic low back pain. Hayes. Published February 9, 2017 (annual review February 15, 2018). Accessed August 4, 2021.
- MacVicar J, Kreiner DS, Duszynski B, Kennedy DJ. Appropriate Use Criteria for Fluoroscopically Guided Diagnostic and Therapeutic Sacroiliac Interventions: Results from the Spine Intervention Society Convened Multispecialty Collaborative. *Pain Med.* 2017;18(11):2081-2095. doi:10.1093/pm/pnx253
- 7. Chou R, Hashimoto R, Friedly J, et al. *Pain Management Injection Therapies for Low Back Pain.* Rockville (MD): Agency for Healthcare Research and Quality (US); 2015.
- 8. Chou R. Subacute and chronic low back pain: nonsurgical interventional treatment. UpToDate. <u>www.uptodate.com</u>. Updated June 10, 2021. Accessed August 4, 2021.
- Chou R, Qaseem A, Snow V, et al. Diagnosis and treatment of low back pain: a joint clinical practice guideline from the American College of Physicians and the American Pain Society [published correction appears in Ann Intern Med. 2008 Feb 5;148(3):247-8]. Ann Intern Med. 2007;147(7):478-491. doi:10.7326/0003-4819-147-7-200710020-00006
- Chou R, Qaseem A, Owens DK, Shekelle P; Clinical Guidelines Committee of the American College of Physicians. Diagnostic imaging for low back pain: advice for high-value health care from the American College of Physicians [published correction appears in Ann Intern Med. 2012 Jan 3;156(1 Pt 1):71]. Ann Intern Med. 2011;154(3):181-189. doi:10.7326/0003-4819-154-3-201102010-00008
- 11. Chou R, Loeser JD, Owens DK, et al. Interventional therapies, surgery, and interdisciplinary rehabilitation for low back pain: an evidence-based clinical practice guideline from the American Pain Society. *Spine (Phila Pa 1976)*. 2009;34(10):1066-1077. doi:10.1097/BRS.0b013e3181a1390d
- 12. Heggeness MH. AAOS endorses back pain guidelines. *AAOS Now*. <u>https://www.mainegeneral.org/app/files/public/6460f387-09dc-4968-b162-</u> <u>eee6121a1497/aaosbackpainguidelines.pdf</u>. Published September 2010. Accessed August 4, 2021.
- 13. Laslett M. Evidence-based diagnosis and treatment of the painful sacroiliac joint. *J Man Manip Ther*. 2008;16(3):142-152. doi:10.1179/jmt.2008.16.3.142
- Maas ET, Ostelo RW, Niemisto L, et al. Radiofrequency denervation for chronic low back pain. *Cochrane Database Syst Rev.* 2015;(10):CD008572. Published 2015 Oct 23. doi:10.1002/14651858.CD008572.pub2.
- 15. Manchikanti L, Datta S, Derby R, et al. A critical review of the American Pain Society clinical practice guidelines for interventional techniques: part 1. Diagnostic interventions. *Pain Physician*. 2010;13(3):E141-E174.
- 16. Manchikanti L, Datta S, Gupta S, et al. A critical review of the American Pain Society clinical practice guidelines for interventional techniques: part 2. Therapeutic interventions. *Pain Physician*. 2010;13(4):E215-E264.
- Simopoulos TT, Manchikanti L, Gupta S, et al. Systematic Review of the Diagnostic Accuracy and Therapeutic Effectiveness of Sacroiliac Joint Interventions. *Pain Physician*. 2015;18(5):E713-E756.



- 18. Soloman M, Mekhail MN, Mekhail N. Radiofrequency treatment in chronic pain. *Expert Rev Neurother*. 2010;10(3):469-474. doi:10.1586/ern.09.153
- 19. Staal JB, de Bie R, de Vet HC, Hildebrandt J, Nelemans P. Injection therapy for subacute and chronic low-back pain. *Cochrane Database Syst Rev.* 2008;2008(3):CD001824. Published 2008 Jul 16. doi:10.1002/14651858.CD001824.pub3
- 20. Qaseem A, Wilt TJ, McLean RM, et al. Noninvasive Treatments for Acute, Subacute, and Chronic Low Back Pain: A Clinical Practice Guideline From the American College of Physicians. *Ann Intern Med.* 2017;166(7):514-530. doi:10.7326/M16-2367
- Chen CH, Weng PW, Wu LC, Chiang YF, Chiang CJ. Radiofrequency neurotomy in chronic lumbar and sacroiliac joint pain: A meta-analysis. *Medicine (Baltimore)*. 2019;98(26):e16230. doi:10.1097/MD.00000000016230
- 22. North American Spine Society. Evidence-based clinical guidelines for multidisciplinary spine care: diagnosis and treatment of low back pain. <u>https://www.spine.org/Portals/0/assets/downloads/ResearchClinicalCare/Guidelines/LowBackPain.pdf</u>. 2020. Accessed August 4, 2021.