

Clinical Policy: Biofeedback

Reference Number: PA.CP.MP.168

Effective Date: 01/18

Date of Last Review: 7/29/2022

[Coding Implications](#)

[Revision Log](#)

Description

Biofeedback therapy provides visual, auditory or other evidence of the status of certain body functions so that a person can exert voluntary control over the functions, and thereby alleviate an abnormal bodily condition. Biofeedback therapy often uses electrical devices to transform bodily signals indicative of such functions as heart rate, blood pressure, skin temperature, salivation, peripheral vasomotor activity, and gross muscle tone into a tone or light, the loudness or brightness of which shows the extent of activity in the function being measured.¹

Note: For neurofeedback for behavioral health conditions, refer to *CP.BH.300 Neurofeedback for Behavioral Health Conditions*.

Policy/Criteria

- I. It is the policy of PA Health and Wellness® (PHW) that biofeedback is **medically necessary** for the following indications when the basic and treatment-specific criteria in A and B are met.

Reconsideration of medical necessity should be made if more than 14 biofeedback treatments sessions in a 12 month period are necessary.

A. Basic Criteria- meets all of the following:

1. The individual is motivated to actively participate in the treatment plan and agrees to the plan of care requirements, (e.g., practice and follow-through at home);
 - a. If a child, support and guidance are available for fulfillment of the plan of care, (e.g., practice and follow-through at home);
2. The individual is capable of participating in the treatment plan (physically as well as intellectually);
3. There is a readily identifiable and measurable response;
4. Biofeedback training is performed by a physician or qualified non-physician practitioner which can include physical and occupational therapists, nurse practitioners, physician assistants, and clinical nurse specialists.

B. Treatment-Specific Criteria - meets any of the following:

1. Stress, urge, or mixed urinary incontinence in cognitively intact adult females who have failed a documented four week trial of Kegel pelvic muscle exercise training;
2. Dysfunctional voiding in children, when other alternative options have been unsuccessful (e.g., timed voiding, prophylactic antibacterial therapy for recurrent urinary tract infections, short term anticholinergic medications to assist developing a normal voiding pattern);
3. Fecal incontinence and all of the following:
 - a. One of the following:
 - i. Anorectal manometry demonstrates weakness of the external anal sphincter;

- ii. Decreased ability to perceive rectal distension because of nerve injury;
- b. None of the following contraindications:
 - i. Isolated internal anal sphincter weakness;
 - ii. Overflow incontinence associated with behavioral or psychiatric disorders;
 - iii. Neurological disorders associated with substantial loss of rectal sensation and/or the inability to contract the external anal sphincter;
 - iv. Decreased rectal storage capacity from resection, inflammation, or fibrosis;
 - v. Suspected or established major structural damage to continence mechanisms;
- 4. Chronic constipation in patients with organic neuromuscular impairment who have difficulty with outlet obstruction;
- 5. Thermal biofeedback combined with relaxation training or electromyography (EMG) biofeedback as treatment options in management of tension and migraine headaches;
- 6. Chronic pain as part of a rehabilitation program;
- 7. Muscle re-education of specific muscle groups or for treating pathological muscle abnormalities of spasticity, incapacitating muscle spasm (including pain due to spasm), or weakness when more conventional treatments (heat, cold, massage, exercise, support) have not been successful.

II. It is the policy of PHW that there is insufficient evidence to draw conclusions regarding the efficacy of biofeedback for any other circumstances than those specified above.

Background

The three most commonly used forms of biofeedback therapy are: (1) electromyography (EMG), which measures muscle tension; (2) thermal biofeedback, which measures skin temperature; and (3) neurofeedback or electroencephalography (EEG), which measures brain wave activity. Various forms of biofeedback appear to be effective for a narrow range of health problems.

First line treatment of urinary incontinence (stress, urgency, mixed) consists of behavioral treatments with an emphasis on improving quality of life. Initial treatment includes lifestyle modifications and pelvic floor muscle exercise (Kegel exercises). Biofeedback is used as an adjunct to pelvic floor muscle exercises. By providing individuals with concurrent feedback on muscle tone, biofeedback is intended to improve the patient's ability to perform pelvic muscle exercises. Augmented versions also use abdominal and perineal EMG recordings to demonstrate improper contraction of abdominal and gluteal muscles. A systematic review and meta-analysis of 17 randomized or quasi-randomized trials found that compared with women who received pelvic floor muscle exercises alone, those that also received biofeedback were more likely to report improvement or cure of urinary incontinence.¹

Dysfunctional voiding in children is a learned behavior of abnormal urination, which often evolves from attempts to suppress impending or active bladder contractions by inappropriately contracting the pelvic floor muscles, thereby tightening the urinary sphincter complex. Symptoms vary but daytime wetness and urinary tract infections are common. Other urinary symptoms include urgency, frequency, infrequency, and constipation. Usual care of dysfunctional voiding includes voiding on a schedule and keeping voiding diaries. Kegel or pelvic floor exercises may help children gain conscious control of pelvic floor musculature and urination.² Biofeedback teaches children how to identify and control the muscle groups involved

in voiding. It is reserved for children with dysfunctional voiding despite an adequate trial of conservative therapy and/or pharmacotherapy. Available studies suggest that biofeedback-directed pelvic floor exercises can improve urinary function in dysfunctional voiding, including those who have previously failed conservative treatment. Biofeedback therapy may result in a faster resolution of symptoms than traditional pelvic floor training without biofeedback.

Biofeedback therapy improves symptoms in more than 70% of patients with defecatory disorders. Biofeedback can be useful in the treatment of constipation to train patients to relax their pelvic floor muscles during straining and to correlate relaxation and pushing to achieve defecation. By the relearning process, the non-relaxing pelvic floor is gradually suppressed and normal coordination restored. Biofeedback has been shown to improve rectoanal coordination during defecation and symptoms of constipation despite reduced laxative use. Biofeedback is also used in the treatment of fecal incontinence.³

American Gastroenterological Association

Pelvic floor retraining by biofeedback therapy rather than laxatives is recommended for defecatory disorders (strong recommendation, high-quality evidence).³

Instrumented anorectal biofeedback therapy should be used to manage symptoms in defecatory disorders (strong recommendation; minimal risk of harm; quality of evidence: moderate.)¹⁷

American Society of Colon and Rectal Surgeons

Biofeedback may be considered as an initial treatment for patients with fecal incontinence and some preserved voluntary sphincter contraction when there is no response to simple dietary modification, medications, and other supportive measures. In their most recent guidelines on the treatment of fecal incontinence, the American Society of Colon and Rectal Surgeons assign a strong recommendations in favor of biofeedback.⁴

American Academy of Neurology

The American Academy of Neurology recommends relaxation training, thermal biofeedback combined with relaxation training, EMG biofeedback, and cognitive-behavioral therapy as treatment options for prevention of migraine (Grade A). Specific recommendations regarding which of these to use for specific patients cannot be made.⁵

Coding Implications

This clinical policy references Current Procedural Terminology (CPT®). CPT® is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2020, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

CPT® Codes	Description
90901	Biofeedback training by any modality

CPT® Codes	Description
90912	Biofeedback training, perineal muscles, anorectal or urethral sphincter, including EMG and/or manometry, when performed; initial 15 minutes of one-on-one physician or other qualified health care professional contact with the patient
90913	Biofeedback training, perineal muscles, anorectal or urethral sphincter, including EMG and/or manometry, when performed; each additional 15 minutes of one-on-one physician or other qualified health care professional contact with the patient (List separately in addition to code for primary procedure)

HCPCS Codes	Description
N/A	

ICD-10-CM Diagnosis Codes that Support Coverage Criteria

ICD-10-CM Code	Description
G43.001 - G43.719	Migraine headache
G44.201 - G44.209	Tension-type headache
K59.00 - K59.09	Constipation
K59.4	Anal spasm
M62.40 - M62.49	Contracture of muscle
M62.50 - M62.59	Muscle wasting and atrophy, not elsewhere classified
N39.3 - N39.498	Stress incontinence
R15.0 - R15.9	Fecal incontinence

Reviews, Revisions, and Approvals	Revision Date	Approval Date
References reviewed and updated.	06/18	
Removed information note that improvement of fecal/urinary incontinence should be noted in 4 sessions. Removed criteria point under I.A stating including being responsive to care plan requirements and the condition can be appropriately treated with biofeedback. Added child specific sub item point under I.A.1. Codes reviewed	10/19	
References reviewed and updated. Removed CPT 90911 – code deleted 1/1/2020. Replaced with 2020 CPT codes, 90912 and 90913. Removed I.B.5 “Anal muscle abnormalities of spasticity,	06/2020	

incapacitating muscle spasm, and/or muscle weakness” as duplicative and revised language in I.B.3. Added contraindications to I.B.3.b.		
Annual Review completed. In II, replaced “experimental /investigational” language with the statement that there is insufficient evidence to draw conclusions regarding the efficacy of biofeedback for any other circumstances than those specified above. Added note to refer to CP.BH.300 Neurofeedback for behavioral health conditions. Updated background with no impact on criteria. Changed “review date” in the header to “date of last revision” and “date” in the revision log header to “revision date.” References reviewed, updated and reformatted. Reviewed by specialist.	7/29/2022	

References

1. Herderschee R, Hay-Smith EJ, Herbison GP, Roovers JP, Heineman MJ. Feedback or biofeedback to augment pelvic floor muscle training for urinary incontinence in women. *Cochrane Database Syst Rev* 2011; (7) CD009252. Published 2011 Jul 6. doi:10.1002/14651858.CD009252
2. American Gastroenterological Association, Bharucha AE, Dorn SD, Lembo A, Pressman A. American Gastroenterological Association medical position statement on constipation. *Gastroenterology*. 2013;144(1):211-217. doi:10.1053/j.gastro.2012.10.029
3. Paquette IM, Varma MG, Kaiser AM, Steele SR, Rafferty JF. The American Society of Colon and Rectal Surgeons' Clinical Practice Guideline for the Treatment of Fecal Incontinence. *Dis Colon Rectum*. 2015;58(7):623-636. doi:10.1097/DCR.0000000000000397
4. Silberstein SD. Practice parameter: evidence-based guidelines for migraine headache (an evidence-based review): report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology* 2000 [published correction appears in *Neurology* 2000 Jan 9;56(1):142]. *Neurology* 2000;55(6):754-762. doi:10.1212/wnl.55.6.754
5. Bertotto A, Schvartzman R, Uchôa S, Wender MCO. Effect of electromyographic biofeedback as an add-on to pelvic floor muscle exercises on neuromuscular outcomes and quality of life in postmenopausal women with stress urinary incontinence: A randomized controlled trial. *Neurourol Urodyn*. 2017;36(8):2142-2147. doi:10.1002/nau.23258
6. Liu J, Zeng J, Wang H, Zhou Y, Zeng C. *Zhonghua Fu Chan Ke Za Zhi*. 2014;49(10):754-757.
7. Hirakawa T, Suzuki S, Kato K, Gotoh M, Yoshikawa Y. Randomized controlled trial of pelvic floor muscle training with or without biofeedback for urinary incontinence. *Int Urogynecol J*. 2013 24(8):1347-1354. doi:10.1007/s00192-012-2012-8
8. Fitz FF, Resende AP, Stüpp L, Sartori MG, Girão MJ, Castro RA. Biofeedback for the treatment of female pelvic floor muscle dysfunction: a systematic review and meta-analysis. *Int Urogynecol J*. 2012 ;23(11):1495-1516. doi:10.1007/s00192-012-1707-1
9. Tugtepe H, Thomas DT, Ergun R, et al. Comparison of biofeedback therapy in children with treatment-refractory dysfunctional voiding and overactive bladder. *Urology*. 2015;85(4):900-904. doi:10.1016/j.urology.2014.12.031

10. Krzemińska K, Maternik M, Drożyńska-Duklas M, et al. High efficacy of biofeedback therapy for treatment of dysfunctional voiding in children. *Cent European J Urol*. 2012;65(4):212-215. doi:10.5173/ceju.2012.04.art6
11. Kajbafzadeh AM, Sharifi-Rad L, Ghahestani SM, Ahmadi H, Kajbafzadeh M, Mahboubi AH. Animated biofeedback: an ideal treatment for children with dysfunctional elimination syndrome. *J Urol*. 2011;186(6):2379-2384. doi:10.1016/j.juro.2011.07.118
12. Desantis DJ, Leonard MP, Preston MA, Barrowman NJ, Guerra LA. Effectiveness of biofeedback for dysfunctional elimination syndrome in pediatrics: a systematic review. *Pediatr Urol*. 2011;7(3):342-348. doi:10.1016/j.jpuro.2011.02.019
13. Robson KM, Lembo A.J. Fecal incontinence in adults: Management. UpToDate. www.uptodate.com. Published November 2, 2021. Accessed November 18, 2021.
14. MacIntosh A, Lam E, Vigneron V, Vignais N, Biddiss E. Biofeedback interventions for individuals with cerebral palsy: a systematic review. *Disabil Rehabil*. 2019;41(20):2369-2391. doi:10.1080/09638288.2018.1468933
15. Dinces EA. Treatment of tinnitus. UpToDate. www.uptodate.com. Published October 6, 2021. Accessed November 18, 2021.
16. Bharucha AE, Rao SSC, Shin AS. Surgical Interventions and the Use of Device-Aided Therapy for the Treatment of Fecal Incontinence and Defecatory Disorders. *Clin Gastroenterol Hepatol*. 2017;15(12):1844-1854. doi:10.1016/j.cgh.2017.08.023.
17. Wald A, Bharucha AE, Limketkai B, et al. ACG Clinical Guidelines: Management of Benign Anorectal Disorders. *Am J Gastroenterol*. 2021;116(10):1987-2008. doi:10.14309/ajg.0000000000001507
https://journals.lww.com/ajg/fulltext/2021/10000/acg_clinical_guidelines_management_of_benign.13.aspx
18. National Coverage Determination. Biofeedback Therapy (30.1) Centers for Medicare and Medicaid Services Web site. <http://www.cms.hhs.gov/mcd/search.asp>. Published July 1, 2001. Accessed December 2, 2021.
19. Clemens JQ. Urinary incontinence in men. UpToDate. www.uptodate.com. Published December 2, 2019. Accessed December 2, 2021.
20. Comiter CV, Speed J. Urinary incontinence after prostate treatment. UpToDate. www.uptodate.com. Published May 21, 2021. Accessed December 2, 2021.