

# Clinical Policy: Intestinal and Multivisceral Transplant

Reference Number: PA.CP.MP.58

Effective Date: 01/18

Date of Last Review: 5/17/2022

[Coding Implications](#)

[Revision Log](#)

## Description

Medical necessity criteria for the review of intestinal and multivisceral transplant requests

## Policy/Criteria

It is the policy of Pennsylvania Health & Wellness® (PHW) that any of the intestinal and/or multivisceral transplantation procedures listed in **I** are **medically necessary** for pediatric and adult members/enrollees to restore function in those with irreversible intestinal failure when meeting the criteria in section **II**:

### I. Transplantation Procedures

- A. Isolated *intestinal transplantation* is indicated for members/enrollees who have only isolated intestinal failure and no liver disease.
- B. Combined *intestinal and liver transplant* is indicated in those with intestinal failure and end stage liver disease.
- C. *Multivisceral transplant* is indicated in those with intestinal failure and gastrointestinal motility disorders (e.g., chronic idiopathic intestinal pseudo-obstruction, visceral myopathy, visceral neuropathy, total intestinal aganglionosis, and some forms of mitochondrial respiratory chain disorders that affect gastrointestinal motor function), or extensive mesenteric thrombosis.

### II. Procedure Criteria: Members must have one of the indications in **A** and none of the contraindications in **B**:

#### A. Indications, any one of the following:

- 1. Failure of total parenteral nutrition (TPN) as indicated by one of the following:
  - a. Impending or overt liver failure due to TPN, indicated by elevated serum bilirubin and/or liver enzymes, splenomegaly, thrombocytopenia, gastro-esophageal varices, coagulopathy, peristomal bleeding, or hepatic fibrosis/cirrhosis;
  - b. Thrombosis of  $\geq 2$  central veins, including jugular, subclavian, and femoral veins;
  - c. Two or more episodes of systemic sepsis due to line infection, per year, or one episode of septic shock, acute respiratory distress syndrome, and/or line related fungemia;
  - d. Frequent episodes of dehydration despite IV fluid supplementation;
  - e. Other complications leading to loss of vascular access;
- 2. High risk of death if transplant is not performed;
- 3. Severe short bowel syndrome (gastrostomy, duodenostomy, residual small bowel <10 cm in infants and <20 cm in adults);
- 4. Frequent hospitalizations for complications directly related to intestinal failure;
- 5. Significant hepatic cirrhosis associated with diffuse post-mesenteric thrombosis;

#### B. Does not have ANY of the following contraindications:

- 1. Malignancy with high risk of recurrence or death related to cancer;
- 2. Other severe uncontrolled medical condition expected to limit survival after transplant;

## Intestinal and Multivisceral Transplant

3. Glomerular filtration rate < 40 mL/min/1.73m<sup>2</sup> unless being considered for multi-organ transplant;
4. HIV infection with detectable viral load;
5. Presence of other GI diseases;
6. Acute liver failure, or cirrhosis with portal hypertension or synthetic dysfunction unless being considered for multi-organ transplant;
7. Septic shock;
8. Progressive cognitive impairment;
9. Stroke, acute coronary syndrome, or myocardial infarction (excluding demand ischemia) within 30 days;
10. Chronic infection with highly virulent and/or resistant microbes that are poorly controlled pre-transplant;
11. Inability to adhere to the regimen necessary to preserve the transplant, even with caregiver support;
12. Absence of an adequate or reliable social support system;
13. Active substance use or dependence including current tobacco use, vaping, marijuana smoking, or IV drug use without convincing evidence of risk reduction behaviors, such as meaningful and/or long-term participation in therapy for substance abuse and/or dependence. Serial blood and urine testing may be used to verify abstinence from substances that are of concern.

### Background

Intestinal transplantation is a therapeutic option for patients with intestinal failure. Intestinal failure is the loss of absorptive capacity of the small bowel secondary to severe primary gastrointestinal disease or surgically induced short bowel syndrome (SBS). The normal small intestine length varies widely, ranging from 3 to 8 meters. SBS occurs when there is approximately < 200 cm of small bowel remaining.

Multi-visceral transplantation includes the stomach, duodenum, pancreas, liver, and small intestine. A modified version excludes the liver if the recipient's liver is normal. A kidney transplant is occasionally included if the recipient has end-stage renal disease.

Common indications for intestinal transplantation in children include:

- |  |  |
|--|--|
| • Small bowel atresia  | • Microvillus inclusion disease        |
| • Gastroschisis  | • Short gut syndrome                   |
| • Aganglionosis (Hirschsprung's disease)                               | • Trauma                               |
| • Infections such as necrotizing enterocolitis and mesenteric ischemia | • Crohn's disease                      |
| • Intestinal pseudo-obstruction  | • Midgut volvulus                      |
|  | • Massive resection secondary to tumor |

Common indications for intestinal transplantation in adults include:

- |   |   |
|---|---|
| • Short gut syndrome  | • Small bowel secretory disorders               |
| • Mesenteric ischemia following thrombosis, embolism, volvulus, or trauma | • Tumors of mesenteric root and retroperitoneum |
| • Crohn's disease   | • Trauma  |
| • Small bowel tumors  | • Volvulus                                      |

- Pseudo-obstruction
- Radiation enteritis

### *Guideline Recommendations*

The British Society of Gastroenterology (2006) recommends: patients with SBS, including irreversible intestinal failure, expected to die prematurely on TPN, should be referred for consideration of short bowel transplant where appropriate.

The American Society of Transplantation (AST, 2001) issued a position paper on indications for pediatric intestinal transplantation. The AST recommends intestinal transplantation only for TPN-dependent children with intestinal failure who have or are likely to develop life-threatening TPN-related complications such as liver disease, recurrent sepsis, and threatened loss of central venous access. The AST stated that intestinal transplantation should not be performed solely because of continued dependence on TPN.

### **Coding Implications**

This clinical policy references Current Procedural Terminology (CPT®). CPT® is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2020, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

CPT® Codes	Description
44135	Intestinal allotransplantation; from cadaver donor
44136	Intestinal allotransplantation; from living donor
44715	Backbench standard preparation of cadaver or living donor intestine allograft prior to transplantation, including mobilization and fashioning of the superior mesenteric artery and vein
44720	Backbench reconstruction of cadaver or living donor intestine allograft prior to transplantation; venous anastomosis, each
44721	Backbench reconstruction of cadaver or living donor intestine allograft prior to transplantation; arterial anastomosis, each
47135	Liver allotransplantation, orthotopic, partial or whole, from cadaver or living donor, any age
47143	Backbench standard preparation of cadaver donor whole liver graft prior to allotransplantation, including cholecystectomy, if necessary, and dissection and removal of surrounding soft tissues to prepare the vena cava, portal vein, hepatic artery, and common bile duct for implantation; without trisegment or lobe split
47144	Backbench standard preparation of cadaver donor whole liver graft prior to allotransplantation, including cholecystectomy, if necessary, and dissection and removal of surrounding soft tissues to prepare the vena cava, portal vein, hepatic artery, and common bile duct for implantation; with trisegment split of whole liver

**CLINICAL POLICY**  
**Intestinal and Multivisceral Transplant**



CPT® Codes	Description
	graft into two partial liver grafts (i.e., left lateral segment (segments II and III) and right trisegment (segments I and IV through VIII))
47145	Backbench standard preparation of cadaver donor whole liver graft prior to allotransplantation, including cholecystectomy, if necessary, and dissection and removal of surrounding soft tissues to prepare the vena cava, portal vein, hepatic artery, and common bile duct for implantation; with lobe split of whole liver graft into two partial liver grafts (i.e., left lobe (segments II, III, and IV) and right lobe (segments I and V through VIII))
47146	Backbench reconstruction of cadaver or living donor liver graft prior to allotransplantation; venous anastomosis, each
47147	Backbench reconstruction of cadaver or living donor liver graft prior to allotransplantation; arterial anastomosis, each

HCPCS Codes	Description
S2053	Transplantation of small intestine and liver allografts
S2054	Transplantation of multivisceral organs
S2055	Harvesting of donor multivisceral organs, with preparation and maintenance of allografts; from cadaver donor
S2152	Solid organs(s), complete or segmental, single organ or combination of organs; deceased or living donor(s), procurement, transplantation, and related complications; including: drugs; supplies; hospitalization with outpatient follow-up; medical/surgical, diagnostic, emergency, and rehabilitative services; and the number of days of pre- and post-transplant care in the global definition

**ICD-10-CM Diagnosis Codes that Support Coverage Criteria**

ICD-10-CM Code	Description
A41.89	Other specified sepsis
A41.9	Sepsis, unspecified organism
K50.00-K52.9	Non-infective colitis and enteritis
K55.011-K55.9	Vascular disorders of intestine
K56.0-K56.7	Paralytic ileus and intestinal obstruction without hernia
K70.0-K77	Diseases of liver
P76.8	Other specified intestinal obstruction of newborn
P77.1-P77.9	Necrotizing enterocolitis of newborn
Q41.0-Q41.9	Congenital absence, atresia and stenosis of small intestine
Q43.1	Hirschsprung's disease
R65.20-R65.21	Severe sepsis
S35.299(A/D/S)	Unspecified injury of branches of celiac and mesenteric artery, initial, subsequent encounter and sequela
T86.850-T86.859	Complication of intestine transplant

**CLINICAL POLICY**  
**Intestinal and Multivisceral Transplant**



ICD-10-CM Code	Description
Z94.82	Intestine transplant status

Reviews, Revisions, and Approvals	Date	Approval Date
Policy developed	11/17	1/1/18
Specialist review (Surgical Transplant)		
References reviewed and updated	09/18	10/18
References reviewed and updated. Added CPT-47135	12/19	1/30/2020
Edited malignancy contraindication to not specify within 2 years, and added exceptions early stage prostate cancer, cancer that has been completely resected, or that has been treated and poses acceptable future risk. Clarified in I.C that multivisceral transplants are indicated in gastrointestinal motility disorders, along with examples of such. Added ICD 10 Q43.1 References reviewed and updated. Specialist reviewed. Annual Review performed.	2/26/2021	
Annual Review performed. Replaced contraindications of “severely limited functional status with poor rehabilitation potential” and those regarding past or current nonadherence to medical therapy, and psychological condition associated with the inability to comply with medical therapy with “Inability to adhere to the regimen necessary to preserve the transplant, even with caregiver support.” Changed “review date” in header to “Date of Last Revision” and “Date” in the revision log header to “Revision Date.” Edited contraindications: Replaced “non-hepatic malignancy...” with malignancy with high risk of recurrence or death...”; added GFR restriction, added HIV infection with detectable viral load, added stroke, acute coronary syndrome, or MI; added acute renal failure...; added septic shock; added progressive cognitive impairment; replaced “untreatable significant dysfunction of another major organ system...” with “Other severe uncontrolled medical condition expected to limit survival after transplant;” slightly reworded substance use contraindication; removed “acute medical instability...”; removed “uncorrectable bleeding diathesis.” Specialist reviewed.		

**References**

1. American Gastroenterological Association Clinical Practice Committee. AGA technical review on short bowel syndrome and intestinal transplantation. *Gastroenterology* 2003;124:1111-1134.
2. Bischel MD. Medical review criteria guidelines for managed care: Intestinal/multivisceral transplants. Apollo Managed Care Inc. Twelfth Edition, 2013.
3. Buchman, Alan L., et al. Parenteral Nutrition-Associated Liver Disease and the Role for Isolated Intestine and Intestine/Liver Transplantation. *Hepatology*, 2006;43-1: pp. 9-19., doi:10.1002/hep.20997.

4. Centers for Medicare & Medicaid Services. National Coverage Determination (NCD) for intestinal and multi-visceral transplantation (260.5). Effective May 11, 2006. Accessed April 29, 2021.
5. Khan FA Selvaggi G. Overview of intestinal and multivisceral transplantation. In: UpToDate, Brown RS (Ed). Published Sept 16, 2020. Accessed April 29, 2021.
6. Gilroy RK. Intestinal and multivisceral transplantation. Medscape Reference, Shaprio R (Ed), Apr 6, 2015. Updated May 24, 2018.
7. Greenstein SM. Intestinal transplantation. Medscape Reference, Mancini MC (Ed). Updated March 16, 2021.
8. Kato T, et al. Intestinal and multivisceral transplantation in children. *Ann Surg*. 2006; 243(6):756-64; discussion 764-6.
9. Kaufman SS, Atkinson JB, Bianchi A, et al. American Society of Transplantation. Indications for pediatric intestinal transplantation: a position paper of the American Society of Transplantation. *Pediatr Transplant*. 2001;5(2):80-87.
10. Kubal CA, Mangus RS, Tector AJ. Intestine and multivisceral transplantation: Current status and future directions. *Curr Gastroenterol Rep*. 2015;17(1):427
11. Levitsky J, Doucette K. Viral Hepatitis in Solid Organ Transplantation. *Am J Transplant*. 2013; 13: 147–168.
12. Llad L, et al. Management of portal vein thrombosis in liver transplantation: influence on morbidity and mortality. *Clin Transplant*. 2007;21: 716721.
13. Mangus RS, et al. Multivisceral transplantation: expanding indications and improving outcomes. *J Gastrointest Surg*. 2013 Jan;17(1):179-86.
14. Nightingale J, Woodward JM. Small Bowel and Nutrition Committee of the British Society of Gastroenterology. Guidelines for management of patients with a short bowel. *Gut*. 2006;55(Suppl IV):iv1-iv12.
15. Nishida S. Pediatric intestinal and multivisceral transplantation. Medscape Reference, Greenstein SM (Ed). Published February 22, 2021.  
<http://emedicine.medscape.com/article/1013915-overview>
16. Troppmann C, Gruessner RW. Intestinal transplantation. In: Surgical Treatment: Evidence-Based and Problem-Oriented, Holzheimer RG, Mannick JA (Ed), Munich: Zuckschwerdt; 2001.
17. Tzakis AG, et al. 100 multivisceral transplants at a single center. *Ann Surg*. 2005 October; 242(4): 480–493.
18. U.S. Department of Veteran Affairs. HIV/ AIDS. Laboratory Tests and HIV: Entire Lesson. Available at: <http://www.hiv.va.gov/patient/diagnosis/labtests-single-page.asp>
19. Vianna RM, et al. Multivisceral transplantation for diffuse portomesenteric thrombosis. *Ann Surg*. 2012 Jun;255(6):1144-50.
20. Wu G, Cruz RJ. Liver inclusion improves outcomes of intestinal retransplantation in adults. *Transplantation* 2015; 99:1265.
21. Dunn CP. Intestinal Transplantation. Medscape Reference. Mancini MC (Ed), January 18, 2017.
22. Bharadwaj S, Tandon P, Gohel TD, et al. Current status of intestinal and multivisceral transplantation. *Gastroenterol Rep (Oxf)*. 2017 Feb;5(1):20-28.
23. Pironi L, Arends J, Bozzetti F, et al. ESPEN guidelines on chronic intestinal failure in adults. *Clin Nutr*. 2016 Apr;35(2):247-307

## CLINICAL POLICY

### Intestinal and Multivisceral Transplant



24. Hawksworth JS, Desaid CS, Khan KM, et al. Visceral transplantation in patients with intestine-failure associated liver disease: evolving indications, graft selection, and outcomes. *Am J Transplant*. 2018 Jun; 18(6): 1312–1320.
25. Camilleri M. Chronic intestinal pseudo-obstruction. In: UpToDate, Friedman LS (Ed.), Published February 4, 2021. Accessed April 29, 2021.