

Clinical Policy: Adefovir (Hepsera)

Reference Number: PA.CP.PHAR.142

Effective Date: 10.17.18

Last Review Date: 10.17.18

[Revision Log](#)

Description

Adefovir (Hepsera®) is a nucleotide analogue and reverse transcriptase inhibitor with activity against human hepatitis B virus.

FDA Approved Indication(s)

Hepsera is indicated for the treatment of chronic hepatitis B in patients 12 years of age and older with evidence of active viral replication and either evidence of persistent elevations in serum aminotransferases (ALT or AST) or histologically active disease.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with PA Health & Wellness® that Hepsera is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Chronic Hepatitis B Infection (must meet all):

1. Diagnosis of chronic hepatitis B virus infection;
2. Prescribed by or in consultation with a gastroenterologist, hepatologist, infectious disease specialist, or transplant physician;
3. Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: Pegasys®, entecavir, or tenofovir;
**Prior authorization may be required for Pegasys and entecavir*
4. Hepsera is not prescribed concurrently with tenofovir;
5. Dose does not exceed 10 mg per day.

Approval duration: 6 months

B. Other diagnoses/indications

1. Refer to the off-label use policy if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): PA.CP.PMN.53.

II. Continued Therapy

A. Chronic Hepatitis B Infection (must meet all):

1. Currently receiving medication via Pennsylvania Health and Wellness benefit or member has previously met all initial approval criteria or the Continuity of Care Policy (PA.LTSS.PHAR.01) applies;
2. Member is responding positively to therapy;
3. Hepsera is not prescribed concurrently with tenofovir;
4. If request is for a dose increase, new dose does not exceed 10 mg per day.

Approval duration: 12 months

B. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via Pennsylvania Health and Wellness benefit or member has previously met all initial approval criteria or the Continuity of Care Policy (PA.LTSS.PHAR.01) applies.

Approval duration: Duration of request or 6 months (whichever is less); or

2. Refer to the off-label use policy if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): PA.CP.PMN.53.

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policy – PA.CP.PMN.53 or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

ALT: alamine aminotransferase

AST: aspartate aminotransferase

FDA: Food and Drug Administration

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
entecavir (Baraclude®)	0.5 to 1 mg PO QD	1 mg/day
Pegasys® (peginterferon alfa-2a)	180 mcg SC once weekly for 48 weeks	180 mcg/day
tenofovir disproxil fumarate (Viread®)	300 mg PO QD	300 mg/day
Vemlidy® (tenofovir alafenamide)	25 mg PO QD	25 mg/day

Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): hypersensitivity
- Boxed warning(s): severe acute exacerbations of hepatitis, nephrotoxicity, HIV resistance, lactic acidosis, and severe hepatomegaly with steatosis

Appendix D: General Information

- Hepsera labeling warns against coadministration of Hepsera with tenofovir-containing products. Hepsera may increase serum concentrations of tenofovir-containing products and vice versa, resulting in additive nephrotoxicity and diminishing therapeutic effect. In

the treatment of chronic hepatitis B, tenofovir should not be administered with Hepsera to avoid multi-drug resistance. In patients with concomitant HIV and chronic hepatitis B, treatment with tenofovir is sufficient.

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
Chronic hepatitis B	CrCl \geq 50 mL/min: 10 mg PO QD CrCl 30 to 49 mL/min: 10 mg PO Q48H CrCl 10 to 29 mL/min: 10 mg PO Q72H Hemodialysis: 10 mg every 7 days following dialysis	10 mg/day

VI. Product Availability

Tablet: 10 mg

VII. References

1. Hepsera Prescribing Information. Foster City, CA: Gilead Sciences, Inc.; November 2012. Available at: www.gilead.com. Accessed August 14, 2018.
2. American Association for the Study of Liver Diseases (AASLD). Update on prevention, diagnosis, and treatment of chronic hepatitis B: AASLD 2018 Hepatitis B Guidance. Hepatology 2018; 67(4):1560-1599. DOI 10.1002/hep.29800
3. World Health Organization. Guidelines for the prevention, care and treatment of persons with chronic hepatitis B infection. March 2015. Available at: http://apps.who.int/iris/bitstream/handle/10665/154590/9789241549059_eng.pdf;jsessionid=F33AA940563ABBB8DF1570D876EC494B?sequence=1. Accessed August 14, 2018.
4. Micromedex[®] Healthcare Series [Internet database]. Greenwood Village, Colo: Thomson Healthcare. Updated periodically. Accessed August 14, 2018.

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created	10/18	