


## Prior Authorization Review Panel

### Prior Authorization Review Panel

#### CHC-MCO Policy Submission

A separate copy of this form must accompany each policy submitted for review.  
Policies submitted without this form will not be considered for review.

<b>Plan: PA Health &amp; Wellness</b>	<b>Submission Date: 05/01/2020</b>
<b>Policy Number: PA.CP.PMN.218</b>	<b>Effective Date: 05/2020</b> <b>Revision Date: 05/2020</b>
<b>Policy Name: Lasmiditan (Reyvow)</b>	
<p><b>Type of Submission – <u>Check all that apply:</u></b></p> <p> <input checked="" type="checkbox"/> <b>New Policy</b>  <input type="checkbox"/> <b>Revised Policy*</b>  <input type="checkbox"/> <b>Annual Review - No Revisions</b>  <input type="checkbox"/> <b>Statewide PDL - Select this box when submitting policies for Statewide PDL implementation and when submitting policies for drug classes included on the Statewide PDL.</b> </p>	
<p><b>*All revisions to the policy <u>must</u> be highlighted using track changes throughout the document.</b></p> <p><b>Please provide any changes or clarifying information for the policy below:</b></p> <p style="text-align: center;"><b>New Policy Created</b></p>	
<b>Name of Authorized Individual (Please type or print):</b>  <b>Francis G. Grillo, MD</b>	<b>Signature of Authorized Individual:</b>  

## **Clinical Policy: Lasmiditan (Reyvow)**

Reference Number: PA.CP.PMN.218

Effective Date: 05/2020

Last Review Date: 05/2020

[Revision Log](#)

### **Description**

Lasmiditan (Reyvow<sup>TM</sup>) is a serotonin (5-HT) 1F agonist.

### **FDA Approved Indication(s)**

Reyvow is indicated for the acute treatment of migraine with or without aura in adults.

Limitation(s) of use: Reyvow is not indicated for the preventive treatment of migraine.

### **Policy/Criteria**

*Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.*

It is the policy of health plans affiliated with PA Health & Wellness that Reyvow is **medically necessary** when the following criteria are met:

#### **I. Initial Approval Criteria**

##### **A. Migraines (must meet all):**

1. Diagnosis of migraine headaches;
2. Age  $\geq$  18 years;
3. Failure of at least TWO formulary 5HT<sub>1B/1D</sub>-agonist migraine medications (see <https://papdl.com/preferred-drug-list> for list of preferred agents) at up to maximally indicated doses, unless contraindicated or clinically significant adverse effects are experienced;
4. For requests for quantities greater than 4 doses per month, member meets one of the following (a or b):
  - a. Failure of TWO prophylactic migraine medications, unless contraindicated or clinically significant adverse effects are experienced (*see Appendix B*);
  - b. Member is being treated by or in consultation with a neurologist or a headache specialist;
5. Dose does not exceed 200 mg (2 tablets) per day and 4 days per month.

**Approval duration:** 12 months

##### **B. Other diagnoses/indications**

1. Refer to the off-label use policy if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): PA.CP.PMN.53

#### **II. Continued Therapy**

##### **A. Migraines (must meet all):**

1. Currently receiving medication via PA Health & Wellness benefit and documentation supports positive response to therapy or the Continuity of Care policy (PA.LTSS.PHAR.01) applies;

2. Member is responding positively to therapy;
3. If request is for a dose increase, new dose does not exceed 200 mg (2 tablets) per day and 4 days per month.

**Approval duration:** 12 months

**B. Other diagnoses/indications** (must meet 1 or 2):

1. Currently receiving medication via PA Health & Wellness benefit and documentation supports positive response to therapy or the Continuity of Care policy (PA.LTSS.PHAR.01) applies.

**Approval duration:** Duration of request or 12 months (whichever is less); or

2. Refer to the off-label use policy if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): PA.CP.PMN.53

**III. Diagnoses/Indications for which coverage is NOT authorized:**

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – PA.CP.PMN.53

**IV. Appendices/General Information**

*Appendix A: Abbreviation/Acronym Key*

5-HT: serotonin

AAN: American Academy of Neurology

FDA: Food and Drug Administration

*Appendix B: Therapeutic Alternatives*

*This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.*

Abortive Migraine Therapy		
Drug Name	Dosing Regimen	Dose Limit/Maximum Dose
<b><i>Triptans</i></b>		
naratriptan (Amerge <sup>®</sup> )	One tablet (1 or 2.5 mg) PO at onset; can be repeated in 4 hours	5 mg/day
almotriptan (Axert <sup>®</sup> )	6.25 to 12.5 mg PO QD May repeat dose in 2 hours	25 mg/day
frovatriptan (Frova <sup>®</sup> )	2.5 mg PO QD May repeat dose in 2 hours	7.5 mg/day
sumatriptan (Imitrex <sup>®</sup> nasal spray)	One spray (5 to 20 mg) at onset into one nostril; can be repeated in 2 hours	40 mg/day
sumatriptan (Imitrex <sup>®</sup> )	One tablet (25 to 100 mg) PO at onset; can be repeated in two hours	200 mg/day
rizatriptan (Maxalt <sup>®</sup> /Maxalt MLT <sup>®</sup> )	One tablet (5 or 10 mg) PO at onset of migraine headache; can be repeated in two hours	30 mg/day

Abortive Migraine Therapy		
eletriptan (Relpax <sup>®</sup> )	20 or 40 mg PO QD May repeat dose in 2 hours	40 mg/dose 80 mg/day
zolmitriptan (Zomig <sup>®</sup> /Zomig <sup>®</sup> ZMT)	1.25 or 2.5 mg PO QD May repeat dose in 2 hours	5 mg/dose 10 mg/day
Prophylactic Migraine Therapy		
Drug Name	Dosing Regimen	Level of Evidence*
<b>Antiepileptic Drugs**</b>		
divalproex sodium (Depakote <sup>®</sup> )	500 to 1,000 mg/day PO	Level A (AAN; AHS)
divalproex sodium ER (Depakote <sup>®</sup> ER)	500 to 1,000 mg/day PO	Level A (AAN; AHS)
topiramate (Topamax <sup>®</sup> )	100 mg/day PO	Level A (AAN; AHS)
<b>Beta-Blockers</b>		
metoprolol (Lopressor <sup>®</sup> )	200 mg/day PO	Level A (AAN; AHS)
propranolol (Inderal <sup>®</sup> )	80 to 240 mg/day PO	Level A (AAN; AHS)
timolol (Blocadren <sup>®</sup> )	20 to 30 mg/day PO	Level A (AAN; AHS)
atenolol (Tenormin <sup>®</sup> )	100 mg/day PO	Level B (AAN; AHS)
nadolol (Corgard <sup>®</sup> )	80 to 240 mg/day PO	Level B (AAN; AHS)
<b>Serotonin Reuptake Inhibitors</b>		
venlafaxine XR (Effexor XR <sup>®</sup> )	150 mg/day PO	Level B (AAN; AHS)
<b>Tricyclic Antidepressants</b>		
amitriptyline (Elavil <sup>®</sup> )	30 to 150 mg/day PO	Level B (AAN; AHS)
<b>CGRP Inhibitors**</b>		
Aimovig (erenumab)	70 mg SC once a month; may be increased to 140 mg SC once a month	140 mg/month
Ajovy (fremanezumab)	225 mg SC once a month or 675 mg SC every 3 months	225 mg/month or 675 mg/3 months
Emgality (galcanezumab)	240 mg SC as a single loading dose, followed by 120 mg SC once a month	120 mg/month

Therapeutic alternatives are listed as Brand name<sup>®</sup> (generic) when the drug is available by brand name only and generic (Brand name<sup>®</sup>) when the drug is available by both brand and generic.

\*American Headache Society (AHS) 2018, American Academy of Neurology (AAN) 2012: Level A: established efficacy, Level B: probably effective, Level C: possibly effective.

\*\*FDA approved.

#### Appendix C: Contraindications/Boxed Warnings

None reported

*Appendix D: General Information*

- The AAN recommends that prophylactic migraine medications should be considered if the patient experiences 2 or more attacks per month that produce aggregate disability of 3 or more days/month.
- The AAN and the National Headache Foundation recommend that prophylactic migraine medications should be considered if one or more of the following are present: greater than 2 migraine headaches per week; migraines cause significant impairment in daily routine even with abortive treatment; contraindication to, adverse effects, overuse or failure of abortive migraine medications, presence of uncommon migraine condition (e.g., basilar migraine); or patient requesting prophylactic therapy.

**V. Dosage and Administration**

Indication	Dosing Regimen	Maximum Dose
Migraines	50 mg, 100 mg, or 200 mg taken orally, as needed	200 mg/dose

**VI. Product Availability**

Tablets: 50 mg, 100 mg

**VII. References**

1. Reyvow Prescribing Information. Indianapolis, IN: Lilly USA, LLC; October 2019. Available at: <http://pi.lilly.com/us/reyvow-uspi.pdf>. Accessed November 11, 2019.
2. Kuca B, Silberstein SD, Wietecha L, et al. Lasmiditan is an effective acute treatment for migraine. *Neurology*. 2018;91:e2222-32.
3. Goadsby PJ, Wietecha LA, Dennehy EB, et al. Phase 3 randomized, placebo-controlled, double-blind study of lasmiditan for acute treatment of migraine. *Brain*. 2019;142:1894-1904.
4. American Headache Society. The American Headache Society position statement on integrating new migraine treatments into clinical practice. *Headache*. 2019;59:1-18.
5. MICROMEDEX<sup>®</sup> Healthcare Series [Internet database]. Greenwood Village, Colo: Thomson Healthcare. Updated periodically. Accessed November 11, 2019.

Reviews, Revisions, and Approvals	Revision Date
Policy created	05/2020