


Prior Authorization Review Panel

CHC-MCO Policy Submission

A separate copy of this form must accompany each policy submitted for review.
Policies submitted without this form will not be considered for review.

Plan: PA Health & Wellness	Submission Date: 11/01/2020
Policy Number: PA.CP.PHAR.358	Effective Date: 01/2020 Revision Date: 10/2020
Policy Name: Gemtuzumab Ozogamicin (Mylotarg)	
<p>Type of Submission – <u>Check all that apply</u>:</p> <p> <input type="checkbox"/> New Policy <input checked="" type="checkbox"/> Revised Policy* <input type="checkbox"/> Annual Review - No Revisions <input type="checkbox"/> Statewide PDL - <i>Select this box when submitting policies for Statewide PDL implementation and when submitting policies for drug classes included on the Statewide PDL.</i> </p>	
<p>*All revisions to the policy <u>must</u> be highlighted using track changes throughout the document.</p> <p>Please provide any changes or clarifying information for the policy below:</p> <p>for acute promyelocytic leukemia, added age limit and removed requirement for use only for relapse as NCCN compendia include use for induction and consolidation; FDA/NCCN dosage limitations added; updated age limit to 1 month from 18 years for new diagnosed AML as per FDA label; references reviewed and updated</p>	
Name of Authorized Individual (Please type or print): Auren Weinberg, MD	Signature of Authorized Individual: 

Clinical Policy: Gemtuzumab Ozogamicin (Mylotarg)

Reference Number: PA.CP.PHAR.358

Effective Date: 10.03.17

Last Review Date: 11.20

[Revision Log](#)

Description

Gemtuzumab ozogamicin (Mylotarg™) is a CD33 directed antibody-drug conjugate.

FDA Approved Indication(s)

Mylotarg is indicated for the treatment of:

- Newly-diagnosed CD33-positive acute myeloid leukemia (AML) in adults and pediatric patients 1 month and older
- Relapsed or refractory CD33-positive AML in adults and in pediatric patients 2 years and older

Policy/Criteria

Provider must submit documentation (which may include office chart notes and lab results) supporting that member has met all approval criteria

It is the policy of health plans affiliated with PA Health & Wellness that Mylotarg is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Acute Myeloid Leukemia (must meet all):

1. Diagnosis of CD33-positive AML;
2. Prescribed by or in consultation with an oncologist or hematologist;
3. Member meets (a or b):
 - a. Age \geq 1 month with newly diagnosed disease;
 - b. Age \geq 2 years with relapsed or refractory disease;
4. Request meets one of the following (a, b, c, d, e or f):*
 - a. Age 1 month to $<$ 18 years: Newly diagnosed disease as combination therapy with standard chemotherapy (i and ii):
 - i. Induction - 1 cycle (1 vial): dose does not exceed 0.1 mg/kg (body surface area [BSA] $<$ 0.6 m²) or 3 mg/m² (BSA \geq 0.6 m²) given once;
 - ii. Intensification - 1 cycle (1 vial): dose does not exceed 0.1 mg/kg (BSA $<$ 0.6 m²) or 3 mg/m² (BSA \geq 0.6 m²) given once;
 - b. Age \geq 18 years: Newly diagnosed disease as combination therapy with daunorubicin and cytarabine (i and ii):
 - i. Induction - 1 cycle (1 vial): dose does not exceed 3 mg/m² on Days 1, 4, and 7;
 - ii. Consolidation - 2 cycles (2 vials): dose does not exceed 3 mg/m² on Day 1 of each cycle;
 - c. Age \geq 18 years: Newly diagnosed disease as single-agent therapy (i and ii):

- i. Induction - 1 cycle (1 vial): dose does not exceed 6 mg/m² on Day 1, and 3 mg/m² on Day 8;
- ii. Continuation therapy - 8 cycles (8 vials): dose does not exceed 2 mg/m² on Day 1 of each cycle;
- d. Age ≥ 2 years: Relapsed or refractory disease (single-agent regimen): single course: dose does not exceed 3 mg/m² on Days 1, 4, and 7 (3 vials);
- e. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

Approval duration: 12 months (Up to a total of 10 doses)

B. Acute Promyelocytic Leukemia (off-label) (must meet all):

- 1. Diagnosis of acute promyelocytic leukemia;
- 2. Prescribed by or in consultation with an oncologist or hematologist;
- 3. Age ≥ 2 years;
- 4. Dose is within FDA maximum limit for any FDA-approved indication or is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

Approval duration: 12 months (Up to 10 doses)

C. Other diagnoses/indications

- 1. Refer to PA.CP.PHAR.53 if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized).

II. Continued Therapy

A. All Indications in Section I (must meet all):

- 1. Currently receiving medication via Pennsylvania Health and Wellness benefit or member has previously met all initial approval criteria or the Continuity of Care Policy (PA.LTSS.PHAR.01) applies;
- 2. Member is responding positively to therapy;
- 3. For AML, member has NOT received the maximum recommended doses as described below (a, b or c):
 - a. As combination therapy with daunorubicin and cytarabine for newly diagnosed disease: up to 5 doses;
 - b. As single-agent therapy for newly diagnosed disease: up to 10 doses;
 - c. As single-agent therapy for relapsed or refractory disease: up to 3 doses;
- 4. If request is for a dose increase, request meets one of the following (a, b, c, d, e, or f):*
 - a. Age 1 month to < 18 years: Newly diagnosed disease as combination therapy with standard chemotherapy (i and ii):
 - i. Induction - 1 cycle (1 vial): dose does not exceed 0.1 mg/kg (body surface area [BSA] < 0.6 m²) or 3 mg/m² (BSA ≥ 0.6 m²) given once;
 - ii. Intensification - 1 cycle (1 vial): dose does not exceed 0.1 mg/kg (BSA < 0.6 m²) or 3 mg/m² (BSA ≥ 0.6 m²) given once;
 - b. Age ≥ 18 years: Newly diagnosed disease as combination therapy with daunorubicin and cytarabine (i and ii):
 - i. Induction - 1 cycle (1 vial): dose does not exceed 3 mg/m² on Days 1, 4, and 7;

- ii. Consolidation - 2 cycles (2 vials): dose does not exceed 3 mg/m² on Day 1 of each cycle;
- c. Age ≥ 18 years: Newly diagnosed disease as single-agent therapy (i and ii):
 - i. Induction - 1 cycle (1 vial): dose does not exceed 6 mg/m² on Day 1, and 3 mg/m² on Day 8;
 - ii. Continuation therapy - 8 cycles (8 vials): dose does not exceed 2 mg/m² on Day 1 of each cycle;
- d. Age ≥ 2 years: Relapsed or refractory disease (single-agent regimen): single course: dose does not exceed 3 mg/m² on Days 1, 4, and 7 (3 vials);
- e. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

Approval duration: 12 months (*Approve requested number of doses required to complete therapy and not to exceed a total of 10 doses*)

B. Other diagnoses/indications (must meet 1 or 2):

- 1. Currently receiving medication via PA Health & Wellness benefit or member has met all initial approval criteria or the Continuity of Care Policy (PA.LTSS.PHAR.01) applies;

Approval duration: Duration of request or 6 months (whichever is less); or

- 2. Refer to PA.CP.PHAR.53 if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized).

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policy – PA.CP.PHAR.53 or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

AML: acute myeloid leukemia

FDA: Food and Drug Administration

NCCN: National Comprehensive Cancer Center

Appendix B: General Information

Not applicable

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): hypersensitivity
- Boxed warning(s): hepatotoxicity

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
AML newly-diagnosed (combination regimen)	Adults: <i>Induction:</i> 3 mg/m ² IV (up to one 4.5 mg vial) on Days 1, 4, and 7 in combination with daunorubicin and cytarabine. If a second	<i>Induction:</i> 4.5 mg/dose (1 cycle)

	<p>induction cycle is required, do NOT administer Mylotarg.</p> <p><i>Consolidation:</i> 3 mg/m² IV on Day 1 (up to one 4.5 mg vial) in combination with daunorubicin and cytarabine for 2 cycles.</p> <p>Pediatric patients > 1 month: 3 mg/m² IV with body surface area (BSA) > 0.6 m². 0.1 mg/kg with body surface area (BSA) < 0.6 m²</p>	<p><i>Consolidation:</i> 4.5 mg/dose (2 cycles)</p> <p><i>Induction pediatric:</i> 1 cycle</p> <p><i>Consolidation pediatric:</i> 1 cycle</p>
AML newly-diagnosed (single-agent regimen)	<p><i>Induction:</i> 6 mg/m² IV on Day 1 and 3 mg/m² on Day 8 for 1 cycle</p> <p><i>Continuation:</i> 2 mg/m² IV on Day 1 every 4 weeks for up to 8 cycles</p>	<p><i>Induction:</i> 6 mg/m²/dose (1 cycle)</p> <p><i>Maintenance:</i> 2 mg/m²/dose every 4 weeks (8 cycles)</p>
AML relapsed or refractory (single-agent regimen)	3 mg/m ² IV (up to one 4.5 mg vial) on Days 1, 4, and 7 for 1 cycle	4.5 mg/dose (1 cycle)

VI. Product Availability

Injection: 4.5 mg as a lyophilized cake or powder in a single-dose vial

VII. References

1. Mylotarg Prescribing Information. Wyeth Pharmaceuticals Inc.; Philadelphia, PA. June 2020. Available at: www.mylotarg.com. Accessed August 15, 2020.
2. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at: http://www.nccn.org/professionals/drug_compendium. Accessed August 15, 2020.
3. National Comprehensive Cancer Network. Acute Myeloid Leukemia Version 3.2020. Available at: https://www.nccn.org/professionals/physician_gls/pdf/aml.pdf. Accessed August 15, 2020.

Reviews, Revisions, and Approvals	Date	P&T Approval Date
New policy created	07/18/18	
3Q 2019 annual review: No changes per Statewide PDL implementation 01/01/2020	07/17/19	
4Q 2019 annual review: No changes per Statewide PDL implementation 01-01-2020	10/30/19	
4Q 2020 annual review: for acute promyelocytic leukemia, added age limit and removed requirement for use only for relapse as NCCN compendia include use for induction and consolidation; FDA/NCCN dosage	08/20	11/20

Reviews, Revisions, and Approvals	Date	P&T Approval Date
limitations added; updated age limit to 1 month from 18 years for new diagnosed AML as per FDA label; references reviewed and updated		