


**Prior Authorization Review Panel**

**Prior Authorization Review Panel**

**CHC-MCO Policy Submission**

A separate copy of this form must accompany each policy submitted for review.  
Policies submitted without this form will not be considered for review.

<b>Plan: PA Health &amp; Wellness</b>	<b>Submission Date: 08/01/2021</b>
<b>Policy Number: PA.CP.PHAR.424</b>	<b>Effective Date: 01/15/2020</b> <b>Revision Date: 07/2021</b>
<b>Policy Name: Fulvestrant (Faslodex Injection)</b>	
<p><b>Type of Submission – <u>Check all that apply:</u></b></p> <p> <input type="checkbox"/> New Policy  <input checked="" type="checkbox"/> Revised Policy*  <input type="checkbox"/> Annual Review - No Revisions  <input type="checkbox"/> Statewide PDL - <i>Select this box when submitting policies for Statewide PDL implementation and when submitting policies for drug classes included on the Statewide PDL.</i> </p>	
<p><b>*All revisions to the policy <u>must</u> be highlighted using track changes throughout the document.</b></p> <p><b>Please provide any changes or clarifying information for the policy below:</b></p> <p>3Q 2021 annual review: no significant changes; references reviewed and updated.</p>	
<p><b>Name of Authorized Individual (Please type or print):</b></p> <p>Venkateswara R. Davuluri, MD</p>	<p><b>Signature of Authorized Individual:</b></p> 

## Clinical Policy: Fulvestrant (Faslodex Injection)

Reference Number: PA.CP.PHAR.424

Effective Date: 01/2020

Last Review Date: 07/2021

[Coding Implications](#)  
[Revision Log](#)

### Description

Fulvestrant (Faslodex<sup>®</sup> Injection) is an estrogen receptor antagonist.

### FDA Approved Indication(s)

Faslodex is indicated for the treatment of:

#### Monotherapy

- Hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative advanced breast cancer in postmenopausal women not previously treated with endocrine therapy.
- HR-positive advanced breast cancer in postmenopausal women with disease progression following endocrine therapy.

#### Combination Therapy

- HR-positive, HER2-negative advanced or metastatic breast cancer in postmenopausal women in combination with ribociclib, as initial endocrine based therapy or following disease progression on endocrine therapy.
- HR-positive, HER2-negative advanced or metastatic breast cancer in combination with palbociclib or abemaciclib in women with disease progression after endocrine therapy.

### Policy/Criteria

*Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.*

It is the policy of health plans affiliated with PA Health & Wellness<sup>®</sup> that Faslodex Injection is **medically necessary** when the following criteria are met:

#### I. Initial Approval Criteria

##### A. Breast Cancer (must meet all):

1. Diagnosis of advanced breast cancer (i.e., recurrent, stage III, or stage IV [metastatic]);
2. Prescribed by or in consultation with an oncologist;
3. Age  $\geq$  18 years;
4. Disease is HR-positive (i.e., estrogen or progesterone receptor [ER/PR]-positive);
5. Request meets one of the following (a or b):
  - a. Dose does not exceed 500 mg three times for the first month then once monthly;
  - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Approval duration:** 6 months

##### B. Ovarian, Fallopian Tube, and Primary Peritoneal Cancer (off-label) (must meet all):

1. Diagnosis of ovarian, fallopian tube, or primary peritoneal cancer;
2. Prescribed by or in consultation with an oncologist;
3. Disease is classified as low-grade serous carcinoma;
4. Dose is within FDA maximum limit for any FDA-approved indication or is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Approval duration:** 6 months

**C. Endometrial Carcinoma (off-label) (must meet all):**

1. Diagnosis of endometrial carcinoma;
2. Prescribed by or in consultation with an oncologist;
3. Disease is classified as grade 1 or 2 endometrioid carcinoma;
4. Faslodex is prescribed in one of the following ways (a, b, c, or d):
  - a. For recurrent or metastatic disease;
  - b. For stage II disease, in combination with sequential external beam radiation therapy;
  - c. For stage IIIA or higher disease;
  - d. For disease not suitable for primary surgery;
5. Dose is within FDA maximum limit for any FDA-approved indication or is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Approval duration:** 6 months

**D. Uterine Sarcoma (off-label) (must meet all):**

1. Diagnosis of uterine sarcoma;
2. Prescribed by or in consultation with an oncologist;
3. Disease is classified in one of the following ways (a or b):
  - a. Low-grade endometrial stromal sarcoma;
  - b. HR-positive (i.e., ER/PR-positive) uterine leiomyosarcoma;
4. Faslodex is prescribed in one of the following ways (a, b, c, or d):
  - a. Following total hysterectomy;
  - b. For vaginal or pelvic recurrence;
  - c. For metastatic disease;
  - d. For disease not suitable for primary surgery;
5. Dose is within FDA maximum limit for any FDA-approved indication or is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Approval duration:** 6 months

**E. Other diagnoses/indications**

1. Refer to the off-label use policy if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): PA.CP.PMN.53

**II. Continued Therapy**

**A. All Indications in Section I (must meet all):**

1. Currently receiving medication via PA Health & Wellness benefit and documentation supports positive response to therapy or the Continuity of Care policy (PA.LTSS.PHAR.01) applies;
2. Member is responding positively to therapy;
3. If request is for a dose increase, request meets one of the following (a or b):
  - a. New dose does not exceed 500 mg once monthly;
  - b. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Approval duration:** 12 months

**B. Other diagnoses/indications (must meet 1 or 2):**

1. Currently receiving medication via PA Health & Wellness benefit and documentation supports positive response to therapy or the Continuity of Care policy (PA.LTSS.PHAR.01) applies.

**Approval duration: Duration of request or 6 months (whichever is less);** or

2. Refer to the off-label use policy if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): PA.CP.PMN.53

**III. Diagnoses/Indications for which coverage is NOT authorized:**

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – PA.CP.PMN.53

**IV. Appendices/General Information**

*Appendix A: Abbreviation/Acronym Key*

ER: estrogen receptor

HR: hormone receptor

FDA: Food and Drug Administration

PR: progesterone receptor

HER2: human epidermal growth factor receptor 2

*Appendix B: Therapeutic Alternatives*

Not applicable

*Appendix C: Contraindications/Boxed Warnings*

- Contraindication(s): hypersensitivity
- Boxed warning(s): none reported

**V. Dosage and Administration**

Indication	Dosing Regimen	Maximum Dose
<u>Monotherapy</u> <ul style="list-style-type: none"> <li>• HR-positive, HER2-negative advanced breast cancer in postmenopausal women not previously</li> </ul>	<u>Faslodex:</u> 500 mg IM into buttocks (gluteal area) slowly (1 - 2 minutes per injection) as two 5 mL injections, one in each buttock, on Days 1, 15, 29 and once monthly thereafter.	<u>Faslodex:</u> 500 mg three times for first month then once monthly

Indication	Dosing Regimen	Maximum Dose
<p>treated with endocrine therapy.</p> <ul style="list-style-type: none"> <li>HR-positive advanced breast cancer in postmenopausal women with disease progression following endocrine therapy.</li> </ul>		
<p><u>Combination Therapy</u></p> <ul style="list-style-type: none"> <li>HR-positive, HER2-negative advanced or metastatic breast cancer in postmenopausal women in combination with ribociclib, as initial endocrine based therapy or following disease progression on endocrine therapy.</li> <li>HR-positive, HER2-negative advanced or metastatic breast cancer in combination with palbociclib or abemaciclib in women with disease progression after endocrine therapy.</li> </ul>	<p><u>Faslodex</u>: 500 mg IM into buttocks (gluteal area) slowly (1 - 2 minutes per injection) as two 5 mL injections, one in each buttock, on Days 1, 15, 29 and once monthly thereafter.</p> <p><u>Ribociclib</u>: 600 mg PO QD for 21 consecutive days followed by 7 days off treatment resulting in a complete cycle of 28 days.</p> <p><u>Palbociclib</u>: 125 mg PO QD for 21 consecutive days followed by 7 days off treatment to comprise a complete cycle of 28 days.</p> <p><u>Abemaciclib</u>: 150 mg PO BID.</p> <p><i>Pre/perimenopausal women treated with the combination of Faslodex plus palbociclib, abemaciclib, or ribociclib, should be treated with luteinizing hormone-releasing hormone (LHRH) agonists according to current clinical practice standards.</i></p>	<p><u>Faslodex</u>: 500 mg three times for first month then once monthly</p> <p><u>Ribociclib</u>: 600 mg/day</p> <p><u>Palbociclib</u>: 125 mg/day</p> <p><u>Abemaciclib</u>: 300 mg/day</p>

**VI. Product Availability**

Two 5 mL glass barrels (syringes), each containing 250 mg/5 mL of Faslodex solution for IM injection. The syringes are presented in a tray with polystyrene plunger rod and safety needles (SafetyGlide™) for connection to the barrel.

**VII. References**

1. Faslodex Prescribing Information. Wilmington, DE: AstraZeneca; September 2020. Available at <https://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=83d7a440-e904-4e36-afb5-cb02b1c919f7>. Accessed May 6, 2021.
2. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at: [http://www.nccn.org/professionals/drug\\_compendium](http://www.nccn.org/professionals/drug_compendium). Accessed May 6, 2021.
3. National Comprehensive Cancer Network. Breast Cancer Version 4.2021. Available at: [https://www.nccn.org/professionals/physician\\_gls/pdf/breast.pdf](https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf). May 6, 2021.
4. National Comprehensive Cancer Network. Ovarian Cancer Version 1.2021. Available at: [https://www.nccn.org/professionals/physician\\_gls/pdf/ovarian.pdf](https://www.nccn.org/professionals/physician_gls/pdf/ovarian.pdf). Accessed May 6, 2021.
5. National Comprehensive Cancer Network. Uterine Neoplasms Version 1.2021. Available at: [https://www.nccn.org/professionals/physician\\_gls/pdf/uterine.pdf](https://www.nccn.org/professionals/physician_gls/pdf/uterine.pdf). Accessed May 6, 2021.
6. Clinical Pharmacology [database online]. Tampa, FL: Gold Standard, Inc.; 2021. Available at: <http://www.clinicalpharmacology-ip.com/>.

**Coding Implications**

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J9395	Injection, fulvestrant, 25 mg

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created.	05.14.19	
3Q 2020 annual review: for endometrial carcinoma, added option for us in stage II disease, in combination with sequential external beam radiation therapy; references reviewed and updated.	08/2020	
3Q 2021 annual review: no significant changes; references reviewed and updated.	07/2021	