

Clinical Policy: Beremagene geperpavec-svdt (Vyjuvek)

Reference Number: PA.CP.PHAR.592

Effective Date: 08/2023

Last Review Date: 07/2024

Description

Beremagene Geperpavec (Vyjuvek™) is a herpes-simplex virus type 1 (HSV-1) vector-based gene therapy.

FDA Approved Indication(s)

Vyjuvek is indicated for the treatment of wounds in patients 6 months of age and older with dystrophic epidermolysis bullosa (DEB) with mutations(s) in the *collagen type VII alpha 1 chain (COL7A1)* gene.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of PA Health & Wellness® that Vyjuvek is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Dystrophic Epidermolysis Bullosa (must meet all):

1. Diagnosis of DEB as evidence by COL7A1 gene mutation confirmed by genetic testing (see Appendix E);
2. Prescribed by or in consultation with a geneticist, dermatologist, or histopathologist;
3. Age \geq 6 months;
4. Provider attestation that target wounds are clean in appearance with adequate granulation tissue, has excellent vascularization, and does not appear infected;
5. Documentation of size of target wounds at baseline (see Appendix F);
6. Provider attestation that member is concomitantly receiving standard of care preventative or treatment therapies for wound care (e.g., polymeric membrane, super-absorbent dressings, soft-silicone foam, enzyme alginogel, protease; see Appendix G);
7. Member does not have current evidence or history of squamous cell carcinoma in the area that will undergo treatment;
8. Vyjuvek is not prescribed concurrently with Filsuvez®;
9. Dose does not exceed one of the following (a or b):
 - a. Age 6 months to < 3 years: 1.6×10^9 plaque forming units (PFU) (0.8 mL) weekly;
 - b. Age \geq 3 years: 3.2×10^9 PFU (1.6 mL) weekly.

Approval duration: 6 months

B. Other diagnoses/indications

1. Refer to the off-label use policy if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): PA.CP.PMN.53

II. Continued Therapy

A. Dystrophic Epidermolysis Bullosa (must meet all):

1. Currently receiving medication via PA Health & Wellness benefit and documentation supports positive response to therapy or the Continuity of Care policy (PA.PHARM.01) applies;
2. Member is responding positively to therapy as evidenced by, including but not limited to, improvement in any of the following parameters (a or b):
 - a. Decrease in wound size;
 - b. Decrease in pain severity for target wound sites associated with dressing changes;
3. Provider attestation that member meets both of the following (a and b):
 - a. Continues to have incomplete wound closures that are clean in appearance with adequate granulation tissue, have excellent vascularization, and do not appear infected;
 - b. Vyjuvek is not applied on target wounds that have completely healed;
4. Vyjuvek is not prescribed concurrently with Filsuvez[®];
5. If request is for a dose increase, new dose does not exceed one of the following (a or b):
 - a. Age 6 months to < 3 years: 1.6×10^9 PFU (0.8 mL) weekly;
 - b. Age \geq 3 years: 3.2×10^9 PFU (1.6 mL) weekly.

Approval duration: 6 months

B. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via PA Health & Wellness benefit and documentation supports positive response to therapy or the Continuity of Care policy (PA.PHARM.01) applies.
Approval duration: Duration of request or 12 months (whichever is less); or
2. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): PA.CP.PMN.53

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – PA.CP.PMN.53

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

COL7A1: collagen type VII alpha 1 chain

DEB: dystrophic epidermolysis bullosa

EB: epidermolysis bullosa

FDA: Food and Drug Administration

HSV-1: herpes simplex virus type 1

IFM: immunofluorescence mapping

PFU: plaque forming units

TEM: transmission electron microscopy

Appendix B: Therapeutic Alternatives
Not applicable

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): none
- Boxed warning(s): none

Appendix D: General Information

- DEB is a serious, ultra-rare epidermolysis bullosa (EB) subtype caused by mutations in the *COL7A1* gene.
- Per 2017 Best Practice Guidelines for Skin and Wound Care in EB, the most recent classification for EB names four categories of the condition defined by the level of cleavage at the dermal and epidermal junction:
 - EB simplex (EBS)
 - Junctional EB (JEB)
 - Dystrophic EB (DEB)
 - Kindler syndrome

Appendix E: Diagnosis Information

- Per 2020 Clinical Practice Guidelines for Laboratory Diagnosis of EB, genetic testing is always recommended for the diagnosis of EB. Methods for clinical diagnosis in EB include immunofluorescence mapping (IFM), transmission electron microscopy (TEM), or genetic testing (e.g. next-generation sequencing, whole-exome sequencing, and Sanger sequencing).
 - IFM is recommended to obtain a rapid diagnosis and prognosis, and to prioritize genetic testing and facilitate interpretation of genetic results.
 - TEM is useful in a limited number of cases, and should be performed when IFM and genetic testing do not deliver conclusive results.
- Per 2017 Best Practice Guidelines for Skin and Wound Care in EB, definitive diagnosis is most commonly made from analysis of a skin biopsy using positive immunofluorescence, antigenic mapping, and TEM. Due to rarity of expertise and facilities, diagnosis is generally made using immunofluorescence and antigen mapping.
- No-charge Genetic Testing for Patients with Suspected DEB:
- The Krystal Decode DEB program (Krystal Biotech and Prevention Genetics collaboration) is open to all US residents, including residents of Puerto Rico, with suspected DEB. More information on the Decode DEB program can be found here: <https://www.preventiongenetics.com/sponsoredTesting/decode-deb>. Invitae Epidermolysis Bullosa and Palmoplantar Keratoderma Panel analyzes genes associated with EB. More information can be found on the Invitae website: <https://www.invitae.com/en/providers/test-catalog/test-434344>.

Appendix F: Dose by Wound Size

| Wound Area (cm ²) | Dose (PFU) | Volume (mL) |
|-------------------------------|-----------------------|-------------|
| < 20 | 4 x 10 ⁸ | 0.2 |
| 20 to < 40 | 8 x 10 ⁸ | 0.4 |
| 40 to 60 | 1.2 x 10 ⁹ | 0.6 |

**For wound area over 60 cm², recommended calculating the total dose based on table above until the maximum weekly dose is reached*

Appendix G: Recommended Wound Care for DEB

Per 2017 Best Practice Guidelines for Skin and Wound Care in EB:

- Wounds should be dressed with nonadherent silicone dressings, foam dressings that absorb exudates, and nonadherent silicone-based tape. Diluted bleach baths or compresses, topical antiseptics, and topic antibiotics are used as preventative measures against bacterial infections.
- Standard of Care for EB skin and wound care:
 - First choice of dressing for general EB wounds (when available): PolyMemb, Cutimed Siltec (super-absorbent)
 - First choice of dressing for chronic EB wounds (when available): PolyMem, Flaminal Hydro/Forte
- Recommended dressings for general EB skin and wound care:

| Dressing Type | Brand | Indication/ Function | Contraindication/ Comments | Wear Time |
|---------------------------|--|--|--|---|
| Polymeric membrane | PolyMem | <ul style="list-style-type: none"> • Where cleansing is required • Chronic wounds | <ul style="list-style-type: none"> • Stimulates high levels of exudate • Distinct smell does not necessarily indicate infection • Can still be difficult to retain on vertical surfaces | <ul style="list-style-type: none"> • Change frequently until exudate reduces |
| Super-absorbent dressings | <ul style="list-style-type: none"> • Cutimed Siltec • Sorbion Sachet S • Filvasorb/Vilwasorb Pro • Kerramax Care | <ul style="list-style-type: none"> • High exudate levels | <ul style="list-style-type: none"> • Can be cut between super-absorbent crystals, which appear in rows (as opposed to cutting across the crystal lattice) | |
| Soft silicone mesh | <ul style="list-style-type: none"> • Mepitel • Mepitel One • Adaptic Touch • Cuticell Contact | <ul style="list-style-type: none"> • Moist wound • Contact layer | | |
| Lipido-colloid | <ul style="list-style-type: none"> • Urgo Tul | <ul style="list-style-type: none"> • Moist wound, drier wounds and protection of vulnerable healed areas • Used as an alternative to | <ul style="list-style-type: none"> • Where retention is difficult (e.g., vertical surfaces) | |

| Dressing Type | Brand | Indication/ Function | Contraindication/ Comments | Wear Time |
|-------------------------|---|---|--|---|
| | | soft silicon (see above) in the presence of over-granulation | | |
| Soft silicone foam | <ul style="list-style-type: none"> • Mepilex • Mepilex Lite • Mepilex Transfer | <ul style="list-style-type: none"> • Absorption of exudate • Protection • Lightly exuding wounds • To transfer exudate to absorbent dressing • Where conformability is required (e.g. digits, axillae) | <ul style="list-style-type: none"> • Over-heating • May need to apply over recommended atraumatic primary dressing | |
| Foam | <ul style="list-style-type: none"> • Allevyn • UrgoTul Absorb • Aquacel Foam | <ul style="list-style-type: none"> • Absorption and protection | <ul style="list-style-type: none"> • May adhere if placed directly on wound bed, use alternative contact layer | |
| Bordered foam dressings | <ul style="list-style-type: none"> • Mepilex Border/ Mepliex Border Lite • Biatain Silicone Border/ Biatain Border Lite • Allevyn Gentle Border • Allevyn Border Lite • Kerrafoam • UrgoTul Absorb Border | <ul style="list-style-type: none"> • Isolated wounds • DDEB and mild RDEB | <ul style="list-style-type: none"> • Bordered dressings may require removal with SMAR to avoid skin stripping • May require primary contact layer • Poor absorption of highly viscous exudate | <ul style="list-style-type: none"> • Up to 4 days depending on personal choice |

| Dressing Type | Brand | Indication/ Function | Contraindication/ Comments | Wear Time |
|---------------|---|--|--|---|
| Keratin | <ul style="list-style-type: none"> • Keragel | <ul style="list-style-type: none"> • Chronic wounds | <ul style="list-style-type: none"> • Dilute with blend emollient if stinging occurs | <ul style="list-style-type: none"> • Reapply with dressing changes |

- Recommended dressings for chronic EB wounds based on consensus opinion

| Dressing Type | Brand | Indications | Contraindication/ Comments | Wear Time |
|--------------------|--|--|--|--|
| Polymeric membrane | <ul style="list-style-type: none"> • PolyMem • PolyMem Max • PolyMem WIC (under a secondary dressing or further layer of PolyMem) | <ul style="list-style-type: none"> • Infected wounds • Recalcitrant wounds | <ul style="list-style-type: none"> • Can provide initial increase in exudate resulting in further skin damage if not properly controlled • Distinct smell does not necessarily indicate infection • Protect periwound skin | <ul style="list-style-type: none"> • Change when wet to avoid hypothermia |
| Enzyme alginogel | <ul style="list-style-type: none"> • Flaminal Hydro • Flaminal Forte | <ul style="list-style-type: none"> • Low exudate • High exudate | <ul style="list-style-type: none"> • Debrides, de-sloughs and antimicrobial • Has some action in modulating excess proteases • Can be used on all wounds apart from third degree burns • Do not use if patient has sensitivity to alginates or polyethylene glycol | <ul style="list-style-type: none"> • Re-apply at each dressing change at least 2 mm thick |
| Honey | | <ul style="list-style-type: none"> • Sensitive wounds | <ul style="list-style-type: none"> • Can cause transient stinging or pain due to its acidity and high osmotic 'pull' • In turn this will contribute to high levels of exudate | |
| Protease modulator | <ul style="list-style-type: none"> • UrgoTul Start range • Promogran | <ul style="list-style-type: none"> • When excess protease may be present | <ul style="list-style-type: none"> • Promogran/ Promogran Prisma may cause initial transient stinging | <ul style="list-style-type: none"> • Frequent dressing changes may be required to |

| Dressing Type | Brand | Indications | Contraindication/Comments | Wear Time |
|---------------|--|-------------|--|------------------|
| | <ul style="list-style-type: none"> Promogran Prisma (with silver) | | <ul style="list-style-type: none"> Excess product cannot be saved once opened as it degrades on contact with air A secondary dressing required and the product may provoke initial heavy exudate | avoid maceration |

V. Dosage and Administration

| Indication | Dosing Regimen | Maximum Dose |
|------------|--|--|
| DEB | <p>Age 6 months to < 3 years: 1.6 x 10⁹ PFU (0.8 mL) topically once weekly</p> <p>Age ≥ 3 years: 3.2 x 10⁹ PFU (1.6 mL) topically once weekly</p> | <p>Age 6 months to < 3 years: 1.6 x 10⁹ PFU/ weekly</p> <p>Age ≥ 3 years: 3.2 x 10⁹ PFU/ weekly</p> |

VI. Product Availability

Biological suspension in a single dose vial (1 mL extractable volume) mixed into excipient gel vial: 5 x 10⁹ PFU/mL

VII. References

- Vyjuvek Prescribing Information. Pittsburgh, PA: Krystal Biotech, Inc.; May 2023. Available at: <https://www.krystallabel.com/pdf/vyjuvek-us-pi.pdf>. Accessed May 13, 2024.
- ClinicalTrials.gov. The objective of this study is to compare the efficacy and safety of Beremagene Geperpavec (B-VEC) topical gel with that of placebo for the treatment of dystrophic epidermolysis bullosa (DEB). Available at: <https://www.clinicaltrials.gov/ct2/show/NCT04491604>. Accessed May 28, 2024.
- Guide S, Gonzalez ME, Bağcı IS, et al. Trial of beremagene geperpavec (B-VEC) for dystrophic epidermolysis bullosa. *N Engl J Med.* 2022;387(24):2211-2219. doi:10.1056/NEJMoa2206663.
- Denyer J, Pillay E, Clapham J, et al. Best practice guidelines for skin and wound care in epidermolysis bullosa. *An International Consensus.* Wounds International, 2017.
- Has C, Liu L, Bolling MC, Charlesworth AV, et al. Clinical practice guidelines for laboratory diagnosis of epidermolysis bullosa. *Br J Dermatol.* 2020 Mar;182(3):574-592. doi: 10.1111/bjd.18128.
- Mellerio JE, El Hachem M, Bellon N, et al. Emergency management in epidermolysis bullosa: consensus clinical recommendations from the European reference network for rare skin diseases. *Orphanet J Rare Dis.* 2020 Jun 6;15(1):142.

7. El Hachem M, Zambruno G, Bourdon-Lanoy E, et al. Multicentre consensus recommendations for skin care in inherited epidermolysis bullosa. *Orphanet J Rare Dis.* 2014 May 20;9:76.

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

| HCPCS Codes | Description |
|-------------|---|
| J3401 | Beremagene geperpavec-svdt for topical administration, containing nominal 5 x 10 ⁹ pfu/mL vector genomes, per 0.1 mL |

| Reviews, Revisions, and Approvals | Date |
|--|---------|
| Policy created | 07/2023 |
| Added HCPCS code [J3401]; added exclusion of concomitant use with Filsuvez. | 04/2024 |
| 3Q 2024 annual review: for Appendix E, updated Decode DEB testing program information and website link; references reviewed and updated. | 07/2024 |