

## Clinical Policy: Bortezomib (Velcade)

Reference Number: PA.CP.PHAR.410

Effective Date: 01.2019

Last Review Date: 01.2023

[Coding Implications](#)  
[Revision Log](#)

### Description

Bortezomib (Velcade®) is a proteasome inhibitor.

FDA Approved Indication(s) Velcade is indicated for treatment of patients with:

- multiple myeloma (MM)
- mantle cell lymphoma (MCL)

### Policy/Criteria

*Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.*

It is the policy of PA Health & Wellness® that Velcade is **medically necessary** when the following criteria are met:

#### I. Initial Approval Criteria

##### A. Multiple Myeloma and Mantle Cell Lymphoma (must meet all):

1. Diagnosis of one of the following (a or b):
  - a. MM;
  - b. MCL (B-cell lymphoma subtype);
2. Prescribed by or in consultation with an oncologist or hematologist;
3. Age  $\geq$  18 years;
4. For Velcade requests, member must use bortezomib, if available, unless contraindicated or clinically significant adverse effects are experienced;
5. Request meets one of the following (a or b):
  - a. Dose does not exceed 1.3 mg/m<sup>2</sup>;
  - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Approval duration:** 6 months

##### B. NCCN Recommended Uses (off-label) (must meet all):

1. Diagnosis of one of the following (a, b, c, d, e, f, g, h or i):
  - a. Kaposi sarcoma - as subsequent systemic therapy, for relapsed/refractory advanced cutaneous, oral, visceral, or nodal disease that has progressed on or not responded to first-line systemic therapy, and progressed on alternate first-line systemic therapy;
  - b. Multicentric Castleman's disease (B-cell lymphoma subtype) - as subsequent therapy;
  - c. Systemic light chain amyloidosis;
  - d. Adult T-cell leukemia/lymphoma - as subsequent therapy;
  - e. Waldenström macroglobulinemia/lymphoplasmacytic lymphoma;
  - f. T-cell acute lymphoblastic leukemia (T-ALL)-for relapsed or refractory disease;

- g. Pediatric acute lymphoblastic leukemia (ALL) - as subsequent therapy;
- h. Pediatric Hodgkin lymphoma (HL) - as subsequent therapy in combination with ifosafamide and vinorelbine;
- i. Other category 1, 2A, or 2B NCCN-recommended uses not listed;
- 2. Prescribed by or in consultation with an oncologist or hematologist;
- 3. Age  $\geq$  18 years (all indications except pediatric ALL and HL);
- 4. For Velcade requests, member must use bortezomib, if available, unless contraindicated or clinically significant adverse effects are experienced;
- 5. Dose is within FDA maximum limit for any FDA-approved indication or is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Approval duration:** 6 months

**C. Other diagnoses/indications**

- 1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): PA.CP.PMN.53.

**II. Continued Therapy**

**A. All Indications in Section I (must meet all):**

- 1. Currently receiving medication via PA Health & Wellness benefit or member has previously met all initial approval criteria or the Continuity of Care policy (PA.LTSS.PHAR.01) applies;
- 2. Member is responding positively to therapy;
- 3. For Velcade requests, member must use bortezomib, if available, unless contraindicated or clinically significant adverse effects are experienced;
- 4. If request is for a dose increase, request meets one of the following (a or b):
  - a. New dose does not exceed 1.3 mg/m<sup>2</sup>;
  - b. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Approval duration:** 12 months

**B. Other diagnoses/indications (must meet 1 or 2):**

- 1. Currently receiving medication via PA Health & Wellness benefit and documentation supports positive response to therapy or the Continuity of Care policy (PA.LTSS.PHAR.01) applies.

**Approval duration: Duration of request or 6 months (whichever is less); or**

- 2. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): PA.CP.PMN.53.

**III. Diagnoses/Indications for which coverage is NOT authorized:**

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policy – PA.CP.PMN.53 or evidence of coverage documents.

#### IV. Appendices/General Information

##### *Appendix A: Abbreviation/Acronym Key*

ALL: acute lymphoblastic leukemia

NCCN: National Comprehensive Cancer  
Network

FDA: Food and Drug Administration

HL: Hodgkin lymphoma

MCL: mantle cell lymphoma

MM: multiple myeloma

T-ALL: T-cell acute lymphoblastic leukemia

##### *Appendix B: Therapeutic Alternatives*

Not applicable

##### *Appendix C: Contraindications/Boxed Warnings*

- Contraindication(s):
  - Patients with hypersensitivity (not including local reactions) to bortezomib, boron, or mannitol, including anaphylactic reactions
  - Contraindicated for intrathecal administration
- Boxed warning(s): none reported

#### V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
MM	<ul style="list-style-type: none"> <li>• <u>First-line therapy</u>: 1.3 mg/m<sup>2</sup> as a 3 to 5 second bolus IV injection or SC injection in combination with PO melphalan and PO prednisone for nine 6-week treatment cycles.</li> <li>• <u>Relapse*</u>: 1.3 mg/m<sup>2</sup> as a 3 to 5 second bolus IV injection or SC injection as a single agent or in combination with dexamethasone for up to eight 3-week cycles. For therapy beyond eight cycles, see PI for additional dosing options.</li> </ul> <p><small>*If relapse occurs ≥ 6 months after a previous response to Velcade, treatment may be restarted at the last tolerated dose.</small></p>	1.3 mg/m <sup>2</sup>
MCL	<ul style="list-style-type: none"> <li>• <u>First-line therapy</u>: 1.3 mg/m<sup>2</sup> as a 3 to 5 second bolus IV injection or SC injection in combination with IV rituximab, cyclophosphamide, doxorubicin and PO prednisone (VcR-CAP) for up to six 3-week treatment cycles, plus two additional cycles if a positive response.</li> <li>• <u>Relapse</u>: 1.3 mg/m<sup>2</sup> as a 3 to 5 second bolus IV injection or SC injection for up to eight 3-week treatment cycles. Therapy may extend beyond eight cycles.</li> </ul>	1.3 mg/m <sup>2</sup>

#### VI. Product Availability

Single-dose vials for injection:

- Sterile lyophilized powder for reconstitution: 1 mg, 2.5 mg, 3.5 mg

- Solution: 2.5 mg/mL, 3.5 mg/1.4 mL

*\*The branded product, Velcade, is only available as 3.5 mg sterile lyophilized powder*

## **VII. References**

1. Velcade Prescribing Information. Cambridge, MA: Millennium Pharmaceuticals, Inc.; August 2022. Available at: [http://www.velcade.com/files/PDFs/VELCADE\\_PRESCRIBING\\_INFORMATION.pdf](http://www.velcade.com/files/PDFs/VELCADE_PRESCRIBING_INFORMATION.pdf). Accessed November 11, 2022.
2. Bortezomib Prescribing Information. Lake Forest, IL: Hospira, Inc.; May 2022. Available at [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2022/209191s000lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/209191s000lbl.pdf). Accessed November 11, 2022.
3. Bortezomib Prescribing Information. Durham, NC: Accord Healthcare; July 2022. Available at: [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2022/215441s000lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/215441s000lbl.pdf). Accessed November 11, 2022.
4. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at: [http://www.nccn.org/professionals/drug\\_compendium](http://www.nccn.org/professionals/drug_compendium). Accessed November 14, 2021.
5. National Comprehensive Cancer Network. T-Cell Lymphomas Version 1.2021. Available at: [https://www.nccn.org/professionals/physician\\_gls/pdf/t-cell.pdf](https://www.nccn.org/professionals/physician_gls/pdf/t-cell.pdf). Accessed November 11, 2022.
6. National Comprehensive Cancer Network. Pediatric Acute Lymphoblastic Leukemia Version 1.2022. Available at: [https://www.nccn.org/professionals/physician\\_gls/pdf/ped\\_all.pdf](https://www.nccn.org/professionals/physician_gls/pdf/ped_all.pdf). Accessed November 11, 2022.
7. National Comprehensive Cancer Network. Acute Lymphoblastic Leukemia Version 2.2021. Available at: [https://www.nccn.org/professionals/physician\\_gls/pdf/all.pdf](https://www.nccn.org/professionals/physician_gls/pdf/all.pdf). Accessed November 11, 2022.
8. National Comprehensive Cancer Network. B-Cell Lymphomas Version 5.2022. Available at: [https://www.nccn.org/professionals/physician\\_gls/pdf/b-cell.pdf](https://www.nccn.org/professionals/physician_gls/pdf/b-cell.pdf). Accessed November 14, 2022.

## **Coding Implications**

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

<b>HCPCS Codes</b>	<b>Description</b>
J9041	Injection, bortezomib (Velcade), 0.1 mg
J9044	Injection, bortezomib (not otherwise specified), 0.1mg

<b>Reviews, Revisions, and Approvals</b>	<b>Date</b>	<b>P&amp;T Approval Date</b>
Policy created.	01/2019	
1Q 2020 annual review: references reviewed and updated.	01/2020	
1Q 2021 annual review: AIDS-related Kaposi sarcoma pediatric HL NCCN recommended uses added; references reviewed and updated.	01/2021	

Reviews, Revisions, and Approvals	Date	P&T Approval Date
1Q 2022 annual review: removed requirement for Velcade to be prescribed in combination with HIV therapy for Kaposi sarcoma indication per NCCN; added T-ALL indication per NCCN; references reviewed and updated.	01/2022	
1Q 2023 annual review; added new 1mg and 2.5mg strengths of bortezomib (available generically only from Hospira); added redirection to generic bortezomib for brand Velcade requests. Added also new 2.5 and 3.5mg formulations (available generically only) as solution for a single-dose injection. Added verbiage directing to the use of generic bortezomib unless contraindicated or clinically significant adverse effects are experienced	01/2023	