

Clinical Policy: Ivabradine (Corlanor)

Reference Number: PA.CP.PMN.70

Effective Date: 01/2018

Last Review Date: 01/2026

Description

Ivabradine (Corlanor[®]) is a hyperpolarization-activated cyclic nucleotide-gated channel blocker.

FDA approved indication(s)

Corlanor is indicated:

- To reduce the risk of hospitalization for worsening heart failure in patients with stable, symptomatic chronic heart failure with left ventricular ejection fraction $\leq 35\%$, who are in sinus rhythm with resting heart rate ≥ 70 beats per minute and either are on maximally tolerated doses of beta-blockers or have a contraindication to beta-blocker use;
- For the treatment of stable symptomatic heart failure due to dilated cardiomyopathy (DCM) in pediatric patients aged 6 months and older, who are in sinus rhythm with an elevated heart rate.

Policy/Criteria

** Provider must submit documentation (including office chart notes and lab results) supporting that member has met all approval criteria**

It is the policy of PA Health & Wellness[®] that Corlanor is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Heart Failure (must meet all):

1. Diagnosis of chronic heart failure;
2. Prescribed by or in consultation with a cardiologist;
3. Age ≥ 6 months;
4. LVEF $\leq 35\%$ for adults or $\leq 45\%$ for pediatrics;
5. Member is in sinus rhythm with a resting heart rate of one of the following (a, b, c, or d):
 - a. Age 6 to 12 months: ≥ 105 beats per minute;
 - b. Age 1 to 3 years: ≥ 95 beats per minute;
 - c. Age 3 to 5 years: ≥ 75 beats per minute;
 - d. Age 5 years and older: ≥ 70 beats per minute;
6. Failure of two of the following beta-blockers recommended for heart failure at up to maximally indicated doses, each used for ≥ 30 days, unless clinically significant adverse effects are experienced or all are contraindicated: bisoprolol, carvedilol (immediate- or extended-release), extended-release metoprolol succinate;
7. Member has used one of the aforementioned beta blockers for ≥ 30 days within the past 60 days, unless clinically significant adverse effects are experienced or all are contraindicated;
8. Dose does not exceed both of the following (a and b):
 - a. 15 mg per day;
 - b. 2 tablets per day or 15 mL per day

Approval duration: 12 months

- B.** Other diagnoses/indications – Refer to PA.CP.PMN.53 if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized)

II. Continued Therapy

A. Heart Failure (must meet all):

1. Currently receiving medication via PA Health & Wellness benefit, or documentation supports that member is currently receiving Corlanor for heart failure, has received this medication for at least 30 days, and is responding positively to therapy or Continuity of Care Policy (PA.PHARM.01) applies;
2. Member is responding positively to therapy;
3. If request is for a dose increase, new dose does not exceed 15 mg (2 tablets or 15 mL) per day.

Approval duration: 12 months

B. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via PA Health & Wellness benefit and documentation supports positive response to therapy or Continuity of Care Policy (PA.PHARM.01) applies;

Approval duration: 12 months

or

2. Refer to PA.CP.PMN.53 if requested indication is NOT listed under section III (Diagnoses/Indications for which coverage is NOT authorized)

III. Diagnoses/Indications for which coverage is NOT authorized:

- A.** Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off-label use policy – PA.CP.PMN.53 or evidence of coverage documents

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

DCM: dilated cardiomyopathy

FDA: Food and Drug Administration

LVEF: left ventricular ejection fraction

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
Beta-Blockers Recommended for Heart Failure		
bisoprolol	Heart Failure[†]	10 mg/day

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
	Initially, 1.25 mg PO QD for 48 hours, then 2.5 mg QD for the first month, then 5 mg QD.	
carvedilol (Coreg [®] , Coreg CR [®])	<p>Heart Failure <u>Immediate-release:</u> Initially, 3.125 mg PO BID for 2 weeks. Dosage may be subsequently increased to 6.25, 12.5, and then 25 mg PO BID over successive intervals of at least 2 weeks.</p> <p><u>Extended-release:</u> Initially, 10 mg PO QD for 2 weeks. Dosage may be subsequently increased to 20, 40, and then 80 mg PO QD over successive intervals of at least 2 weeks.</p>	<p>Immediate-release: 100 mg/day</p> <p>Extended-release: 80 mg/day</p>
metoprolol succinate extended release (Toprol XL [®])	<p>Heart Failure 25 mg PO QD for 2 weeks in patients with NYHA class II heart failure, or 12.5 mg PO QD in patients with more severe heart failure. Double the dose every 2 weeks as tolerated, up to the target dosage of 200 mg PO QD.</p>	200 mg/day

*Therapeutic alternatives are listed as Brand name[®] (generic) when the drug is available by brand name only and generic (Brand name[®]) when the drug is available by both brand and generic.
 †Off-label indication*

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s):
 - Acute decompensated heart failure
 - Clinically significant hypotension
 - Sick sinus syndrome, sinoatrial block, or 3rd degree AV block, unless a functioning demand pacemaker is present
 - Clinically significant bradycardia
 - Severe hepatic impairment
 - Pacemaker dependence (heart rate maintained exclusively by the pacemaker)
 - Concomitant use of strong cytochrome P450 3A4 (CYP3A4) inhibitors
- Boxed warning(s): none reported

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
Heart failure	Adult and pediatric patients ≥ 40 kg: Initially 2.5 mg (pediatrics and vulnerable adults) or 5 mg PO BID. After 2 weeks of treatment, adjust dose based on heart rate. The maximum dose is 7.5 mg BID.	15 mg/day

Indication	Dosing Regimen	Maximum Dose
	Pediatric patients < 40 kg: Initially 0.05 mg/kg PO BID. Adjust dose at 2-week intervals by 0.05 mg/kg based on heart rate. The maximum dose is 0.2 mg/kg (patients aged 6 months to < 1 year) or 0.3 mg/kg (patients aged ≥ 1 year), up to a total of 7.5 mg BID.	

VI. Product Availability

Tablets: 5 mg, 7.5 mg
Oral solution: 5 mg/5 mL

IV. References

1. Corlanor Prescribing Information. Thousand Oaks, CA: Amgen Inc.; August 2021. Available at: <https://www.corlanor.com>. Accessed October 17, 2025.
2. Yancy CW, Jessup M, Bozkurt B, Butler J, et al. American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. 2013 ACCF/AHA guideline for the management of heart failure: a report of the American College of Cardiology Foundation/American Heart Association Task Force on practice guidelines. *Circulation*. 2013 Oct 15;128(16):e240-327.
3. Yancy CW, Jessup M, Bozkurt B, et al. 2016 ACC/AHA/HFSA focused update on new pharmacological therapy for heart failure: an update of the 2013 ACCF/AHA guideline for the management of heart failure: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Failure Society of America. *Circulation*. 2016;134: 000-000.
4. Yancy CW, Jessup M, Bozkurt B, et al. 2017 ACC/AHA/HFSA focused update of the 2013 ACCF/AHA guideline for the management of heart failure: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Failure Society of America. *Circulation*. 2017;136:e137-e161.
5. Heidenreich PA, Bozkurt B, Aguilar D, et al. 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *J Am Coll Cardiol*. 2022 May, 79 (17) e263–e421. <https://doi.org/10.1016/j.jacc.2021.12.012>
6. Clinical Pharmacology [database online]. Philadelphia, PA: Elsevier; 2023. Available at: www.clinicalkeys.com/pharmacology.
7. Maddox TM, Januzzi JL, Allen LA, et al. 2024 ACC Expert Consensus Decision Pathway for treatment of heart failure with reduced ejection fraction: A report of the American College of Cardiology Solution Set Oversight Committee. *JACC*. 2024; 83(15): 1444-1488.

Reviews, Revisions, and Approvals	Date
Added age restriction and DDI contraindication as the interactions are severe per PI/safety approach; Modified max dose requirement to include specific quantity limit. Updated references.	03/2018
1Q 2019 annual review: references reviewed and updated.	01/2019
Aligned initiation approval duration and continued approval duration.	05/2019

Reviews, Revisions, and Approvals	Date
1Q 2020 annual review: added recently FDA-approved pediatric indication extension references reviewed and updated.	01/2020
1Q 2021 annual review: no significant changes; references reviewed and updated.	01/2021
1Q 2021 annual review: references reviewed and updated.	01/2022
1Q 2023 annual review: no significant changes; references reviewed and updated.	01/2023
1Q 2024 annual review: removed commercially unavailable branded therapeutic alternatives; references reviewed and updated.	01/2024
1Q 2025 annual review: no significant changes; updated Section V to include specific weight-based maximum doses for pediatric patients per PI; references reviewed and updated.	01/2025
1Q 2026 annual review: no significant changes; references reviewed and updated.	01/2026