

Clinical Policy: Tadalafil (Adcirca)

Reference Number: PA.CP.PHAR.198

Effective Date: 01/18

Last Review Date: 07/18

[Coding Implications](#)

[Revision Log](#)

Description

The intent of the criteria is to ensure that patients follow selection elements established by Pennsylvania Health and Wellness[®] clinical policy for tadalafil (Adcirca[®])*.

**Tadalafil brand, Adcirca, should be not confused with tadalafil brand, Cialis.*

FDA Approved Indication(s)

Adcirca is indicated for the treatment of pulmonary arterial hypertension (PAH) (World Health Organization [WHO] Group 1) to improve exercise ability.

Studies establishing effectiveness included predominately patients with New York Heart Association (NYHA) Functional Class II-III symptoms and etiologies of idiopathic or heritable PAH (61%) or PAH associated with connective tissue diseases (23%).

Policy/Criteria

It is the policy of health plans affiliated with Pennsylvania Health and Wellness that Adcirca is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Pulmonary Hypertension (must meet all):

1. Diagnosis of PAH;
2. Prescribed by or in consultation with a cardiologist or pulmonologist;
3. Failure of a trial of a calcium channel blocker, unless member meets one of the following (a or b):
 - a. Inadequate response or contraindication to acute vasodilator testing;
 - b. Contraindication or clinically significant adverse effects to a calcium channel blocker are experienced;
4. Prescribed dose of Adcirca does not exceed 40 mg once daily.

Approval duration: 6 months

B. Other diagnoses/indications: Refer to PA.CP.PMN.53

II. Continued Approval

A. Pulmonary Hypertension (must meet all):

1. Currently receiving medication via Pennsylvania Health and Wellness benefit or member has previously met all initial approval criteria; or the Continuity of Care policy (PA.LTSS.PHAR.01) applies;
2. Member is responding positively to therapy;
3. Prescribed dose of Adcirca does not exceed 40 mg once daily.

Approval duration: 12 months

B. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via Pennsylvania Health and Wellness benefit and documentation supports positive response to therapy; or the Continuity of Care policy (PA.LTSS.PHAR.01) applies;
2. Refer to PA.CP.PMN.53

Background

Description/Mechanism of Action:

Adcirca (tadalafil), an oral treatment for pulmonary arterial hypertension, is a selective inhibitor of cyclic guanosine monophosphate (cGMP)–specific phosphodiesterase type 5 (PDE5). Tadalafil is an inhibitor of PDE5, the enzyme responsible for the degradation of cGMP. Pulmonary arterial hypertension is associated with impaired release of nitric oxide by the vascular endothelium and consequent reduction of cGMP concentrations in the pulmonary vascular smooth muscle. PDE5 is the predominant phosphodiesterase in the pulmonary vasculature. Inhibition of PDE5 by tadalafil increases the concentrations of cGMP resulting in relaxation of pulmonary vascular smooth muscle cells and vasodilation of the pulmonary vascular bed.

Formulations:

Adcirca oral tablets: 20 mg

Appendices

Appendix A: Abbreviation Key

- FC: functional classification
- NYHA: New York Heart Association
- PAH: pulmonary arterial hypertension
- PH: pulmonary hypertension
- WHO: World Health Organization

Appendix B: Pulmonary Hypertension: WHO Classification

- Group 1: PAH (pulmonary arterial hypertension)
- Group 2: PH due to left heart disease
- Group 3: PH due to lung disease and/or hypoxemia
- Group 4: CTEPH (chronic thromboembolic pulmonary hypertension)
- Group 5: PH due to unclear multifactorial mechanisms

Appendix C: Pulmonary Hypertension: WHO/NYHA Functional Classes (FC)

Treatment Approach*	FC	Status at Rest	Tolerance of Physical Activity (PA)	PA Limitations	Heart Failure
Monitoring for progression of PH and treatment of co-existing conditions	I	Comfortable at rest	No limitation	Ordinary PA does not cause undue dyspnea or fatigue, chest pain, or near syncope.	

Treatment Approach*	FC	Status at Rest	Tolerance of Physical Activity (PA)	PA Limitations	Heart Failure
Advanced treatment of PH with PH-targeted therapy - see Appendix D**	II	Comfortable at rest	Slight limitation	Ordinary PA causes undue dyspnea or fatigue, chest pain, or near syncope.	
	III	Comfortable at rest	Marked limitation	Less than ordinary PA causes undue dyspnea or fatigue, chest pain, or near syncope.	
	IV	Dyspnea or fatigue may be present at rest	Inability to carry out any PA without symptoms	Discomfort is increased by any PA.	Signs of right heart failure

*PH supportive measures may include diuretics, oxygen therapy, anticoagulation, digoxin, exercise, pneumococcal vaccination. **Advanced treatment options also include calcium channel blockers.

Appendix D: Pulmonary Hypertension: Targeted Therapies

Mechanism of Action	Drug Class	Drug Subclass	Drug	Brand/Generic Formulations	
Reduction of pulmonary arterial pressure through vasodilation	Prostacyclin* pathway agonist	Prostacyclin	Epoprostenol	Velettri (IV) Flolan (IV) Flolan generic (IV)	
	<i>*Member of the prostanoid class of fatty acid derivatives.</i>	Synthetic prostacyclin analog	Treprostinil	Orenitram (oral tablet) Remodulin (IV) Tyvasco (inhalation)	
			Iloprost	Ventavis (inhalation)	
			Non-prostanoid prostacyclin receptor (IP receptor) agonist	Selexipag	Uptravi (oral tablet)
	Endothelin receptor antagonist	Selective receptor antagonist	Ambrisentan	Letairis (oral tablet)	
			Nonselective dual action receptor antagonist	Bosentan	Tracleer (oral tablet)
				Macitentan	Opsummit (oral tablet)
	Nitric oxide-cyclic guanosine monophosphate enhancer	Phosphodiesterase type 5 (PDE5) inhibitor	Sildenafil	Revatio (IV, oral tablet, oral suspension)	
			Tadalafil	Adcirca (oral tablet)	
		Guanylate cyclase stimulant	Riociguat	Adempas (oral tablet)	

Coding Implications

Tadalafil

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
N/A	

Reviews, Revisions, and Approvals	Date	Approval Date
Removed WHO/NYHA classifications from initial criteria since specialist is involved in care. References reviewed and updated	02/18	

References

- i. Adcirca Prescribing Information. Indianapolis, IN: Eli Lilly and Company; August 2017. Available at <http://pi.lilly.com/us/adcirca-pi.pdf>. Accessed November 21, 2017.
- ii. McLaughlin VV, Archer SL, Badesch DB, et al. ACCF/AHA 2009 expert consensus document on pulmonary hypertension: A report of the American College of Cardiology Foundation Task Force on Expert Consensus Documents and the American Heart Association - developed in collaboration with the American College of Chest Physicians, American Thoracic Society, Inc., and the Pulmonary Hypertension Association. *J Am Coll Cardiol*. 2009; 53(17): 1573-1619.
- iii. Taichman D, Ornelas J, Chung L, et. al. CHEST guideline and expert panel report: Pharmacologic therapy for pulmonary arterial hypertension in adults. *Chest*. 2014; 146 (2): 449-475.
- iv. Abman SH, Hansmann G, Archer SL, et al. Pediatric pulmonary hypertension: Guidelines from the American Heart Association and American Thoracic Society. *Circulation*. 2015 Nov 24; 132(21): 2037-99.
- v. Kim NH, Delcroix M, Jenkins DP, et al. Chronic thromboembolic pulmonary hypertension. *J Am Coll Cardiol* 2013; 62(25): Suppl D92-99.
- vi. Galiè N, Humbert M, Vachiary JL, et al. 2015 ESC/ERS Guidelines for the diagnosis and treatment of Pulmonary Hypertension. *European Heart Journal*. Doi:10.1093/eurheartj/ehv317.