

Clinical Policy: Carbamazepine ER (Equetro)

Reference Number: PA.CP.PMN.137

Effective Date: 03.13.18

Last Review Date: 04.18.18

[Revision Log](#)

Description

Carbamazepine extended release (Equetro[®]) is an antiepileptic drug and mood stabilizer.

FDA Approved Indication(s)

Equetro is indicated for the treatment of:

- Acute manic or mixed episodes associated with bipolar I disorder
- Pain associated with trigeminal neuralgia
- Partial seizures with complex symptomatology (e.g., psychomotor, temporal lobe), generalized tonic-clonic seizures (grand mal), and mixed seizure patterns, which include the seizure types listed here or other partial or generalized seizures.

Limitation(s) of use: Equetro is not indicated for the treatment of absence seizures (petit mal). Carbamazepine has been associated with increased frequency of generalized convulsions in these patients.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with PA Health & Wellness[®] that Equetro is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Bipolar Disorder, Trigeminal Neuralgia, or Epilepsy (must meet all):

1. One of the following diagnoses (a, b, or c):
 - a. Bipolar disorder;
 - b. Trigeminal neuralgia;
 - c. Epilepsy (partial seizures, generalized tonic-clonic seizures [grand mal], or mixed types)
2. If diagnosis is bipolar disorder or trigeminal neuralgia, age \geq 18 years;
3. Member has experienced clinically significant adverse effects to immediate release carbamazepine or has contraindication(s) to its excipients;
4. Member has experienced clinically significant adverse effects to extended release carbamazepine (e.g., Tegretol[®] XL) or has contraindication(s) to its excipients;
5. Dose does not exceed:
 - a. Bipolar disorder, epilepsy: 1600 mg/day;
 - b. Trigeminal neuralgia: 1200 mg/day.

Approval duration: 12 months

B. Other diagnoses/indications

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): HIM.PHAR.21 for health insurance marketplace and CP.PMN.53 for Medicaid.

II. Continued Therapy

A. Bipolar Disorder, Trigeminal Neuralgia, or Epilepsy (must meet all):

1. Currently receiving medication via PA Health & Wellness benefit or member has previously met all initial approval criteria or the Continuity of Care policy (PA.LTSS.PHAR.01) applies;
2. Member is responding positively to therapy;
3. If request is for a dose increase, new dose does not exceed:
 - a. Bipolar disorder, epilepsy: 1600 mg/day;
 - b. Trigeminal neuralgia: 1200 mg/day.

Approval duration: 12 months

B. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via PA Health and Wellness benefit and documentation supports positive response to therapy or the Continuity of Care policy (PA.LTSS.PHAR.01) applies; or
2. Refer to CP.PHAR.57 - Global Biopharm Policy.

III. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

FDA: Food and Drug Administration

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
carbamazepine (Carbatrol, Epitol [®] , Tegretol [®] , Tegretol-XR)	<p>Adults: Bipolar Disorder</p> <ul style="list-style-type: none"> • Immediate-release products: Initially, 200 mg PO twice daily. Usual daily dose range is 600 to 1600 mg/day in divided doses. <p>Adults: Trigeminal neuralgia</p> <ul style="list-style-type: none"> • Carbatrol extended-release capsule: Initial: 200 mg PO once on day 1; may increase by 200 mg/day given every 12 hours as needed for efficacy and tolerability (range, 200 to 1200 mg/day; most patients, 400 to 800 mg/day) to max 1200 mg/day • Immediate-release, chewable, or extended-release tablet: Initial: 100 mg 	<p><i>Oral formulations:</i> 1200 mg/day for trigeminal neuralgia. In rare instances, 1600 mg/day for epilepsy or bipolar disorder</p>

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
	<p>PO twice daily on day 1; may increase by 100 mg every 12 hours as needed for pain control (range, 200 to 1200 mg/day; most patients, 400 to 800 mg/day) to max 1200 mg/day</p> <ul style="list-style-type: none"> Suspension: Initial: 50 mg orally 4 times daily on day 1; may increase by 200 mg/day (50 mg 4 times daily) as needed for pain control (range, 200 to 1200 mg/day; most patients, 400 to 800 mg/day) to max 1200 mg/day. <p>Adults: Epilepsy, partial, generalized, and mixed types</p> <ul style="list-style-type: none"> Carbatrol extended-release capsule: Initial: 200 mg PO twice daily for the first week; may increase by adding up to 200 mg/day in 2 divided doses at weekly intervals to the minimum effective level (usually 800 to 1200 mg/day); generally, do not exceed 1200 mg/day and rarely, up to 1600 mg/day may be given Extended-release tablet: Initial: 200 mg PO twice daily for the first week; may increase by adding up to 200 mg/day in 2 divided doses at weekly intervals to the minimum effective level (usually 800 to 1200 mg/day); generally, do not exceed 1200 mg/day and rarely, up to 1600 mg/day may be given Immediate-release or chewable tablet: Initial: 200 mg PO twice daily for the first week; may increase by adding up to 200 mg/day in 3 or 4 divided doses at weekly intervals to the minimum effective level (usually 800 to 1200 mg/day); generally, do not exceed 1200 mg/day, and rarely, up to 1600 mg/day may be given Suspension: Initial, 100 mg PO 4 times daily for the first week; may increase by adding up to 200 mg/day in 3 or 4 divided doses at weekly intervals to the minimum effective level (usually 800 to 1200 mg/day); generally, do not exceed 	

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
	1200 mg/day, and rarely, up to 1600 mg/day may be given	

Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.

IV. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
Acute manic or mixed episodes associated with bipolar I disorder	200 mg PO twice daily; the dose may be increased by 200 mg per day to achieve optimal clinical response. Doses higher than 1600 mg per day have not been studied in mania associated with bipolar disorder.	1600 mg/day
Trigeminal neuralgia	Initial: On the first day, start with one 200 mg capsule PO once daily. This dose may be increased by up to 200 mg/day using increments of 100 mg every 12 hours only as needed to reach an effective and tolerated dose. Do not exceed a total daily dose of 1200 mg. Maintenance: Control of pain can be maintained in most patients with 400 mg to 800 mg daily. However, some patients may be maintained on as little as 200 mg daily, while others may require as much as 1200 mg daily.	1200 mg/day
Epilepsy	<p>Adults and children over 12 years of age: The recommended initial dose is 200 mg PO twice daily. Increase in weekly increments of 200 mg a day, administered as an equally divided, twice daily dose, until an optimal response is obtained. Dosage generally should not exceed 500 mg twice daily in children 12 to 15 years old; 600 mg twice daily in children 15 to 18 years old; and 800 mg twice daily in adults.</p> <p>Children under 12 years of age: Ordinarily, optimal clinical response is achieved at daily doses below 35 mg/kg</p>	1600 mg/day

V. Product Availability

Extended-release capsules: 100 mg, 200 mg, and 300 mg

VI. References

1. Equetro Prescribing Information. Parsippany, NJ: Validus Pharmaceuticals LLC; October 2016. Available at: <http://equetro.com/full-prescribing-information/>. Accessed March 5, 2018.
2. American Psychiatric Association Practice Guideline for the Treatment of Patients with Bipolar Disorder: Second Edition (2010). Available at: http://psychiatryonline.org/pb/assets/raw/sitewide/practice_guidelines/guidelines/bipolar.pdf. Accessed online March 5, 2018.
3. Clinical Pharmacology [database online]. Tampa, FL: Gold Standard, Inc.; 2018. Available at: <http://www.clinicalpharmacology-ip.com/>.
4. Micromedex® Healthcare Series [Internet database]. Greenwood Village, Colo: Thomson Healthcare. Updated periodically. Accessed March 5, 2018.

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created	03.13.18	04.18.18