

Clinical Policy: Sapropterin Dihydrochloride (Kuvan, Javygtor, Zelvysia)

Reference Number: PA.CP.PHAR.43

Effective Date: 01/2018

Last Review Date: 04/2026

Description

Sapropterin dihydrochloride (Kuvan[®]) is a synthetic form of tetrahydrobiopterin (BH4), the cofactor for the enzyme phenylalanine hydroxylase.

FDA Approved Indication(s)

Kuvan is indicated to reduce blood phenylalanine (Phe) levels in adult and pediatric patients one month of age and older with hyperphenylalaninemia (HPA) due to tetrahydrobiopterin- (BH4-) responsive phenylketonuria (PKU). Kuvan is to be used in conjunction with a Phe-restricted diet.

Policy/Criteria

It is the policy of PA Health & Wellness[®] that Kuvan and sapropterin formulations are **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Phenylketonuria (must meet all):

1. Diagnosis of hyperphenylalaninemia (HPA) due to phenylketonuria (PKU);
2. Prescribed by or in consultation with an endocrinologist, metabolic or genetic disease specialist;
3. Recent (within 90 days) phenylalanine (Phe) blood level is ≥ 360 $\mu\text{mol/L}$;
4. Member is currently adherent on a phenylalanine-restricted diet and will continue this diet during treatment with sapropterin;
5. Sapropterin is not prescribed concurrently with Sephience[™] or Palynziq, unless titrating with the plan to discontinue sapropterin;
6. If request is for brand Kuvan, Javygtor or Zelvysia, member must use generic sapropterin, unless contraindicated or clinically significant adverse effects are experienced;
7. Documentation of member's current weight (in kg);
8. Dose does not exceed 20 mg/kg per day.

Approval Duration: 12 months

B. Other diagnoses/indications: Refer to PA.CP.PMN.53

II. Continued Approval

A. Phenylketonuria (must meet all):

1. Currently receiving medication via PA Health & Wellness benefit or member has previously met all initial approval criteria; or the Continuity of Care policy (PA.PHARM.01) applies;
2. Member is responding positively to therapy as demonstrated by a reduction in Phe blood levels since initiation of therapy;

CLINICAL POLICY

Sapropterin Dihydrochloride

3. Member is currently adherent on a phenylalanine-restricted diet and will continue this diet during treatment with sapropterin;
4. If request is for brand Kuvan, Javygtor, or Zelvysia member must use generic sapropterin, unless contraindicated or clinically significant adverse effects are experienced;
5. Sapropterin is not prescribed concurrently with Sephience™ or Palynziq, unless titrating with the plan to discontinue sapropterin;
6. Documentation of member's current weight (in kg);
7. Dose does not exceed 20 mg/kg per day.

Approval Duration: 12 months

B. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via PA Health & Wellness benefit and documentation supports positive response to therapy; or the Continuity of Care policy (PA.PHARM.01) applies; or
2. Refer to PA.CP.PMN.53

III. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

BH4: tetrahydrobiopterin

HPA: hyperphenylalaninemia

FDA: Food and Drug Administration

Phe: phenylalanine

PKU: phenylketonuria

Appendix B: Therapeutic Alternatives

Not applicable

Appendix C: Contraindications/Boxed Warnings

None reported

Appendix D: General Information

- According to the prescribing information, if a 10 mg/kg per day starting dose is used, then response to therapy is determined by change in blood Phe following treatment with Kuvan at 10 mg/kg per day for a period of up to 1 month. Blood Phe levels should be checked after 1 week of Kuvan treatment and periodically for up to a month. If blood Phe does not decrease from baseline at 10 mg/kg per day, the dose may be increased to 20 mg/kg per day. Patients whose blood Phe does not decrease after 1 month of treatment at 20 mg/kg per day are non-responders and treatment with Kuvan should be discontinued in these patients.

IV. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
BH4-responsive PKU	Age 1 month to ≤ 6 years (starting dose) 10 mg/kg QD. Age ≥ 7 years (starting dose): 10 to 20 mg/kg QD	20 mg/kg/day

V. Product Availability

Tablets: 100 mg

Powder for oral solution: 100 mg, 500 mg

VII. References

1. Kuvan Prescribing Information. Novato, CA: BioMarin Pharmaceutical, Inc.; August 2024. Available at www.Kuvan.com. Accessed January 14, 2026.
2. Javygtor Prescribing Information. Princeton, NJ: Dr. Reddy's Laboratories Inc.; January 2022. Available at www.Javygtor.com. Accessed January 14, 2026.
3. Zelvyisia [package insert]. Piscataway, NJ: Aucta Pharmaceuticals, Inc.; April 2025. Accessed January 14, 2026.
4. Levy HL, Milanowski A, Chakrapani A, et. al. Efficacy of sapropterin dihydrochloride (tetrahydrobiopterin, 6R-BH4) for reduction of phenylalanine concentration in patients with phenylketonuria: a phase III randomized placebo-controlled study. *Lancet*. 2007;370(9586):504.
5. Vockly J, Andersson HC, Antshel KM, et al. ACMG practice guidelines: phenylalanine hydroxylase deficiency: diagnosis and management guideline. *Genet Med*. 2014;16(2):188-200.
6. Camp KM, Parisi MA, Acosta PB, et al. Phenylketonuria scientific review conference: state of the science and future research needs. *Mol Genet Metab*. June 2014;112(2):87-122.
7. van Spronsen FJ. Mild hyperphenylalaninemia: to treat or not to treat. *J Inherit Metab Dis*. 2011;34:651-656.
8. Smith WE, Berry SA, Bloom K, et al. Phenylalanine hydroxylase deficiency diagnosis and management: a 2023 evidence-based clinical guideline of the American College of Medical Genetics and Genomics (ACMG). *Genetics in Medicine*. 2024;27(1):101289. doi: <https://doi.org/10.1016/j.gim.2024.101289>.
9. van Spronsen FJ. Mild hyperphenylalaninemia: to treat or not to treat. *J Inherit Metab Dis*. 2011;34:651-656.
10. Smith WE, Berry SA, Bloom K, et al. Phenylalanine hydroxylase deficiency diagnosis and management: a 2023 evidence-based clinical guideline of the American College of Medical Genetics and Genomics (ACMG). *Genetics in Medicine*. 2024;27(1):101289. doi: <https://doi.org/10.1016/j.gim.2024.101289>.
11. Van Wegberg AMJ, MacDonald A, Ahring K, et al. European guidelines on diagnosis and treatment of phenylketonuria: first revision. *Molecular Genetics and Metabolism*. June 2025;145(2):109125. <https://doi.org/10.1016/j.ymgme.2025.109125>.

Reviews, Revisions, and Approvals	Date
1Q 2018 annual review: Use in conjunction with a Phe-restricted diet is removed. Initial approval duration increased from 2 to 3 months to allow adequate time for follow-up. Continuation criteria that refers to an increase in dietary Phe tolerance or improvement in neuropsychiatric symptoms is deleted leaving reduction of Phe levels per the PI. References reviewed and updated.	02/2018
1Q 2019 annual review: references reviewed and updated.	01/2019

CLINICAL POLICY
Sapropterin Dihydrochloride



Reviews, Revisions, and Approvals	Date
1Q 2020 annual review: references reviewed and updated.	01/2020
1Q 2021 annual review: references reviewed and updated.	01/2021
2Q 2021 annual review: added requirements for a Phe-restricted diet and excluded coverage of concurrent use of Kuvan and Palynziq; references reviewed and updated.	04/2021
2Q 2022 annual review: references reviewed and updated.	04/2022
2Q 2023 annual review: no significant changes; references reviewed and updated.	04/2023
2Q 2024 annual review: added redirection to generic product for brand requests; increased initial auth duration to align with those of other drugs for rare diseases; for Continued Therapy added exclusion for concomitant use with Palynziq to match with the Initial Approval Criteria; references reviewed and updated.	04/2024
2Q 2025 annual review: no significant changes; added requirement for a redirection from Javygtor (branded generic) to unbranded generic sapropterin per an SDC recommendation; added a requirement for documentation of member's current weight for dose calculation purposes; references reviewed and updated.	04/2025
Added Sephience (newly FDA-approved for PKU) as an agent that should not be used concomitantly with sapropterin; extended initial approval duration to 12 months; references reviewed and updated.	10/2025
2Q 2026 annual review: no significant changes; added requirement for a redirection from Zelvysia (another branded generic) to unbranded generic sapropterin; added endocrinologist as a possible specialist to align with the Palynziq and Sephience criteria; added adherence to Phe-restricted diet and align with Sephience criteria; references reviewed and updated.	04/2026