

Clinical Policy: Tesamorelin (Egrifta SV)

Reference Number: PA.CP.PHAR.109

Effective Date: 01/2018

Last Review Date: 07/2024

Description

Tesamorelin (Egrifta SV[®]) is a growth hormone releasing factor analog.

FDA Approved Indication(s)

Egrifta SV is indicated for the reduction of excess abdominal fat in human immunodeficiency virus (HIV)-infected adult patients with lipodystrophy.

Limitation(s) of use:

- Long-term cardiovascular safety of Egrifta SV treatment have not been established. Consider risk/benefit of continuation of treatment in patients who have not had a reduction in visceral adipose tissue.
- Egrifta SV is not indicated for weight loss management as it has a weight neutral effect.
- There are no data to support improved compliance with anti-retroviral therapies in HIV-positive patients taking Egrifta SV.

Policy/Criteria

It is the policy of PA Health & Wellness that Egrifta SV[®] is **medically necessary** when one of the following criteria are met:

I. Initial Approval Criteria

A. Human immunodeficiency virus (HIV) with Lipodystrophy (must meet all):

1. Diagnosis of HIV infection with lipodystrophy;
2. Age \geq 18 years;
3. Meets clinical indicators for abdominal lipodystrophy (a or b):
 - a. If female, waist circumference \geq 94 cm and waist-hip ratio \geq 0.88;
 - b. If male, waist circumference \geq 195cm and waist-hip ratio \geq 0.94;
4. Member is currently receiving and adherent to antiretroviral therapy;
5. Prescribed dose of Egrifta SV does not exceed 1.4 mg once daily.

Approval Duration: 6 months

B. Other diagnoses/indications : Refer to PA.CP.PMN.53

II. Continued Approval

A. HIV with Lipodystrophy (must meet all):

1. Currently receiving medication via PA Health & Wellness benefit or member has previously met all initial approval criteria or the Continuity of Care policy (PA.LTSS.PHAR.01) applies;
2. Member is responding positively to therapy;
3. If request is for a dose increase, new dose does not exceed 1.4 mg per day.

Approval Duration: 12 months

B. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via PA Health & Wellness benefit and documentation supports positive response to therapy or the Continuity of Care policy (PA.LTSS.PHAR.01) applies; or
2. Refer to PA.CP.PMN.53

III. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

FDA: Food and Drug Administration

HIV: human immunodeficiency virus

Appendix B: Therapeutic Alternatives

Not applicable

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s):
 - Disruption of the hypothalamic-pituitary axis due to hypophysectomy, hypopituitarism, pituitary tumor/surgery, head irradiation or head trauma
 - Active malignancy. Any preexisting malignancy should be inactive and its treatment complete prior to instituting therapy with Egrifta SV
 - Pregnancy.
 - Known hypersensitivity to tesamorelin or excipients in Egrifta SV
- Boxed warning(s): none reported

Appendix D: General Information

- On June 15, 2020, Theratechnologies discontinued Egrifta and permanently replaced it with Egrifta SV, a smaller volume injection able to be stored at room temperature.

IV. Dosage and Administration

| Indication | Dosing Regimen | Maximum Dose |
|----------------------------------|--|--------------|
| HIV infection with lipodystrophy | 1.4 mg (0.35 mL) SC QD After reconstitution and administration, any unused solution should be thrown away | 1.4 mg/day |

V. Product Availability

Single-use vial with powder for reconstitution: 2 mg

VI. References

1. Egrifta SV Prescribing Information. Montreal, Quebec, Canada: Theratechnologies Inc.; October 2019. Available at <http://www.egriftasv.com>. Accessed May 8, 2024.
2. Falutz J, Mamputu JC, Potvin D, et al. Effects of tesamorelin (TH9507), a growth hormone-releasing factor analog, in human immunodeficiency virus-infected patients with excess abdominal fat: a pooled analysis of two multicenter, double-blind placebo-controlled phase 3

trials with safety extension data. *J Clin Endocrinol Metab.* 2010 Sep;95(9):4291-304. doi: 10.1210/jc.2010-0490.

3. Falutz J, Allas S, Blot K, et al. Metabolic effects of a growth hormone-releasing factor in patients with HIV. *N Engl J Med.* 2007 Dec 6;357(23):2359-70. doi: 10.1056/NEJMoa072375.

| HCPCS Codes | Description |
|-------------|------------------------|
| J3590 | Unclassified biologics |

| Reviews, Revisions, and Approvals | Date |
|--|---------|
| Removed adherence to current antiretroviral therapy on re-auth; pregnancy contraindication added per safety guidance endorsed by Centene Medical Affairs; references reviewed and updated. | 05/2018 |
| 3Q 2019 annual review: No changes per Statewide PDL implementation 01-01-2020 | 07/2019 |
| 3Q 2020 annual review: replaced old formulation Egrifta with new formulation Egrifta SV and updated dose; removed pregnancy contraindication from criteria as separate edits are in place to address these risks; references reviewed and updated. | 07/2020 |
| 3Q 2021 annual review: no significant changes; references reviewed and updated. | 07/2021 |
| 3Q 2022 annual review: no significant changes; added quantity restriction (1 vial per day) to dosing requirement; updated HCPCS codes; references reviewed and updated. | 07/2022 |
| 3Q 2023 annual review: no significant changes; references reviewed and updated. | 07/2023 |
| 3Q 2024 annual review: revised clinical indicators for abdominal lipodystrophy criteria to require waist circumference and waist-hip ratio thresholds that reflect efficacy studies; per PI, revised FDA Approved Indications and contraindications, removed criteria allowing pediatric use in members with closed epiphyses; updated HCPCS codes; references reviewed and updated. | 07/2024 |